

AMSC/CMSC 460: Midterm 2

Prof. Doron Levy

April 17, 2018

**Read carefully the following instructions:**

- Write your name & student ID on the exam book and sign it.
- You may not use any books, notes, or calculators.
- Solve all problems. Answer all problems after carefully reading them. Start every problem on a new page.
- Show all your work and explain everything you write.
- Exam time: 75 minutes
- Good luck!

**Problems: (Each problem = 10 points)**

1. (a) Explain the advantages of interpolating at Chebyshev points.

**Solution:** When interpolating data that is sampled from a function  $f(x)$  at  $n + 1$  points  $x_0, \dots, x_n$ , the interpolation error is given by

$$\frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{i=0}^n (x - x_i).$$

When considering the interval  $[-1, 1]$ , choosing  $x_0, \dots, x_n$  as Chebyshev points (i.e. the  $n + 1$  roots of  $T_{n+1}(x)$ ), minimizes the product term in the error, i.e.,

$$\max_{x \in [-1, 1]} \prod_{i=0}^n (x - x_i),$$

has the smallest value out of all possible choices of interpolation points.

- (b) Compute the unique interpolating polynomial of degree  $\leq 2$  that interpolates data sampled from  $f(x) = x^2$  at an appropriate number of Chebyshev points on the interval  $[-1, 1]$ .

**Solution:** Since we are asked to compute a quadratic polynomial that interpolates data that is sampled from a quadratic function, we can immediately use the uniqueness of the interpolating polynomial and conclude that the answer must be the function itself, i.e.,  $P_2(x) = x^2$ . Any direct calculation must lead to this answer.

- (c) Repeat part (b) with  $f(x) = x^4$ .

**Solution:** Here we should compute the answer. We are seeking a quadratic interpolant, which means that we need 3 values. The 3 values should be the roots of the cubic Chebyshev polynomial  $T_3(x) = 4x^3 - 3x$ :

$$x_0 = -\frac{\sqrt{3}}{2}, \quad x_1 = 0, \quad x_2 = \frac{\sqrt{3}}{2}.$$

Computing the divided differences, we get

$$f(x_0) = \frac{9}{16}, \quad f[x_0, x_1] = -\frac{9}{8\sqrt{3}}, \quad f[x_0, x_1, x_2] = \frac{3}{4}.$$

Hence, the interpolating polynomial is

$$P_2(x) = \frac{9}{16} - \frac{9}{8\sqrt{3}} \left( x + \frac{\sqrt{3}}{2} \right) + \frac{3}{4} \left( x + \frac{\sqrt{3}}{2} \right) x = \dots = \frac{3}{4}x^2.$$

Note: Chebyshev polynomials are given by

$$T_0(x) = 1, \quad T_1(x) = x, \quad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) = 0, \forall n \geq 1.$$

2. Find a spline of degree 2,  $S(x)$ , on the interval  $[0, 2]$ , for which  $S(0) = 0$ ,  $S(1) = 2$ ,  $S(2) = 0$ , and  $S'(0) = 0$ . Use the points  $0, 1, 2$  as the knots.

**Solution:**

Consider a quadratic spline of the form

$$S(x) = \begin{cases} S_0(x), & 0 \leq x \leq 1, \\ S_1(x), & 1 \leq x \leq 2. \end{cases} = \begin{cases} a_0 + a_1x + a_2x^2, & 0 \leq x \leq 1, \\ b_0 + b_1x + b_2x^2, & 1 \leq x \leq 2. \end{cases}$$

We write 6 equations for the 6 unknowns. First we have the 4 interpolation conditions:

- $S_0(0) = 0 \implies a_0 = 0$ .
- $S_0(1) = 2 \implies a_0 + a_1 + a_2 = 2$ .
- $S_1(1) = 2 \implies b_0 + b_1 + b_2 = 2$ .
- $S_1(2) = 0 \implies b_0 + 2b_1 + 4b_2 = 0$ .

The continuity of the first derivative implies that

$$S'_0(1) = S'_1(1) \implies a_1 + 2a_2 = b_1 + 2b_2.$$

Finally, we have the additional condition at the first derivative:

$$S'(0) = 0 \implies a_1 = 0.$$

Solving for  $a_0, a_1, a_2, b_0, b_1, b_2$  we end up with

$$S(x) = \begin{cases} 2x^2, & 0 \leq x \leq 1, \\ -8 + 16x - 6x^2, & 1 \leq x \leq 2. \end{cases}$$

3. Use the Gram-Schmidt process to find orthogonal polynomials of degrees 0, 1, 2, on the interval  $[0, 1]$ , with respect to the weight  $w(x) = 1 + x$ .

Note: you do not need to normalize the polynomials. For the quadratic polynomial,  $P_2(x)$ , write the coefficients but do not explicitly calculate the integrals.

**Solution:** We note that we are only asked to find orthogonal polynomials without normalizing them. Set  $P_0(x) = 1$ . We then let  $P_1(x) = x - cP_0(x)$ . The orthogonality condition,  $\langle P_0, P_1 \rangle_w = 0$ , implies that  $\langle P_0, x - cP_0 \rangle_w = 0$ , i.e.,  $\langle P_0, x \rangle_w = c \langle P_0, P_0 \rangle_w$ , or

$$c = \frac{\langle P_0, x \rangle_w}{\|P_0\|_w^2} = \frac{\int_0^1 x \cdot 1 \cdot (1+x) dx}{\int_0^1 1^2 (1+x) dx} = \dots = \frac{5}{9}.$$

Hence  $P_1(x) = x - \frac{5}{9}$ . For the quadratic polynomial, we set

$$P_2(x) = x^2 - cP_0(x) - dP_1(x).$$

The orthogonality condition  $\langle P_0, P_2 \rangle_w = 0$  implies that

$$c = \frac{\langle x^2, P_0 \rangle_w}{\|P_0\|_w^2} = \frac{\int_0^1 x^2 \cdot 1 \cdot (1+x) dx}{\frac{3}{2}}$$

The orthogonality condition  $\langle P_1, P_2 \rangle_w = 0$  implies that

$$d = \frac{\langle x^2, P_1 \rangle_w}{\|P_1\|_w^2} = \frac{\int_0^1 x^2(x - \frac{5}{9})(x+1) dx}{\int_0^1 (x - \frac{5}{9})^2(x+1) dx}.$$

4. Let  $f(x) = x^2$ . Find the quadratic polynomial  $Q_2^*(x)$  that minimizes

$$\int_{-\infty}^{\infty} e^{-x^2} (f(x) - Q_2(x))^2 dx,$$

among all quadratic polynomials  $Q_2(x)$ .

Note: You may use:

$$H_0(x) = 1, \quad H_1(x) = 2x, \quad H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x), \quad \forall n \geq 1$$

$$\int_{-\infty}^{\infty} e^{-x^2} H_n(x)H_m(x) dx = \delta_{nm} 2^n n! \sqrt{\pi}$$

$$\int_{-\infty}^{\infty} x^m e^{-x^2} dx = \Gamma\left(\frac{m+1}{2}\right), \quad \text{for even } m$$

$$\Gamma(1/2) = \sqrt{\pi}, \quad \Gamma(3/2) = \frac{1}{2}\sqrt{\pi}, \quad \Gamma(5/2) = \frac{3}{4}\sqrt{\pi}.$$

**Solution:** We have the appropriate orthogonal polynomials:

$$H_0(x) = 1, \quad H_1(x) = 2x, \quad H_2(x) = 4x^2 - 2.$$

We note that the norms are

$$\|H_0\|^2 = \sqrt{\pi}, \quad \|H_1\|^2 = 2\sqrt{\pi}, \quad \|H_2\|^2 = 8\sqrt{\pi}.$$

Set

$$Q_2^*(x) = c_0H_0(x) + c_1H_1(x) + c_2H_2(x).$$

Then

$$c_0 = \frac{\langle f, H_0 \rangle_w}{\|H_0\|_w^2} = \frac{\int_{-\infty}^{\infty} e^{-x^2} x^2 dx}{\sqrt{\pi}} = \frac{\frac{\sqrt{\pi}}{2}}{\sqrt{\pi}} = \frac{1}{2}.$$

$$c_1 = \frac{\langle f, H_1 \rangle_w}{\|H_1\|_w^2} = \frac{\int_{-\infty}^{\infty} e^{-x^2} x^2 \cdot 2x dx}{2\sqrt{\pi}} = 0.$$

$$c_2 = \frac{\langle f, H_2 \rangle_w}{\|H_2\|_w^2} = \frac{\int_{-\infty}^{\infty} e^{-x^2} x^2 (4x^2 - 2) dx}{8\sqrt{\pi}} = \frac{4\Gamma(\frac{5}{2}) - 2\Gamma(\frac{3}{2})}{8\sqrt{\pi}} = \frac{4\frac{3}{4}\sqrt{\pi} - 2\frac{1}{2}\sqrt{\pi}}{8\sqrt{\pi}} = \frac{1}{4}.$$

Hence

$$Q_2^*(x) = \frac{1}{2}H_0(x) + \frac{1}{4}H_2(x) = \frac{1}{2} + \frac{1}{4}(4x^2 - 2) = x^2$$

as should be...