

Geometric structures on manifolds

William M. Goldman

December 11, 2021

MATHEMATICS DEPARTMENT, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742 USA

E-mail address: wmg@math.umd.edu

URL: <http://www.math.umd.edu/~wmg>

Key words and phrases. Euclidean geometry, affine geometry, projective geometry, manifold, coordinate atlas, convexity, connection, parallel transport, homogeneous coordinates, Lie groups, homogeneous spaces, metric space, Riemannian metric, geodesic, completeness, developing map, holonomy homomorphism, proper transformation group, Lie algebra, vector field

The author gratefully acknowledges research support from NSF Grants DMS1065965, DMS1406281, DMS1709791 as well as the Research Network in the Mathematical Sciences DMS1107367 (GEAR)..

ABSTRACT. The study of locally homogeneous geometric structures on manifolds was initiated by Charles Ehresmann in 1936, who first proposed the classification of putting a “classical geometry” on a topological manifold. In the late 1970’s, locally homogeneous *Riemannian* structures on 3-manifolds formed the context for Bill Thurston’s *Geometrization Conjecture*, later proved by Perelman. This book develops the theory of geometric structures modeled on a homogeneous space of a Lie group, which are not necessarily Riemannian. Drawing on a diverse collection of techniques, we hope to invite researchers at all levels to this fascinating and currently extremely active area of mathematics.

Contents

Introduction	1
Organization of the text	3
Part One: Affine and Projective Geometry	3
Part Two: Geometric Manifolds	5
Part Three: Affine and projective structures	8
Acknowledgements	10
Prerequisites	12
Notation, terminology and general background	12
Vectors and matrices	12
Projective equivalence	14
General Topology	14
Fundamental group and covering spaces	15
Smooth manifolds	16
Vector fields	18
Part 1. Affine and projective geometry	21
Chapter 1. Affine geometry	23
1.1. Euclidean space	23
1.2. Affine space	24
1.2.1. The geometry of parallelism	25
1.2.2. Affine transformations	26
1.3. The connection on affine space	28
1.3.1. The tangent bundle of an affine space	28
1.3.2. Parallel transport	29
1.3.3. Acceleration and geodesics	30
1.4. Parallel structures	31
1.4.1. Parallel Riemannian structures	31
1.4.2. Parallel tensor fields	32
1.4.3. Complex affine geometry	32
1.5. Affine vector fields	33
1.5.1. Translations and parallel vector fields	33
1.5.2. Homotheties and radiant vector fields	33
1.5.3. Affineness criteria	34

1.6.	Affine subspaces	36
1.7.	Volume in affine geometry	36
1.7.1.	Centers of gravity	37
1.7.2.	Divergence	38
1.8.	Linearizing affine geometry	38
Chapter 2.	Projective geometry	41
2.1.	Ideal points	41
2.2.	Projective subspaces	43
2.2.1.	Affine patches	44
2.3.	Projective mappings	46
2.3.1.	Embeddings and Collineations	46
2.3.2.	Singular projective mappings	47
2.3.3.	Locally projective maps	50
2.4.	Affine patches	51
2.4.1.	Projective vector fields	51
2.5.	Classical projective geometry	52
2.5.1.	One-dimensional reflections	53
2.5.2.	Fundamental theorem of projective geometry	56
2.5.3.	Distance via cross-ratios	59
2.5.4.	Products of reflections	59
2.6.	Asymptotics of projective transformations	60
2.6.1.	Some examples	62
2.6.2.	Higher dimensional projective maps	63
2.6.3.	Limits of similarity transformations	64
2.6.4.	Normality domains	65
Chapter 3.	Duality and non-Euclidean geometry	67
3.1.	Dual projective spaces	68
3.2.	Correlations and polarities	69
3.2.1.	Example: Elliptic geometry	69
3.2.2.	Elliptic polarities and elliptic geometry	70
3.2.3.	Polarities	71
3.2.4.	Quadrics	72
3.3.	Projective model of hyperbolic geometry	74
3.3.1.	The hyperbolic plane	75
3.3.2.	The upper halfspace model of hyperbolic 3-space	76
3.3.2.1.	Quaternions	76
3.3.2.2.	Hyperbolic 3-space	77
Chapter 4.	Convexity	81
4.1.	Convex domains and cones	81
4.1.1.	Halfspaces and supporting hyperplanes	83

4.1.2. Convexity in projective space	84
4.2. The Hilbert metric	84
4.2.1. Definition and basic properties	84
4.2.2. The Hilbert metric on a triangle	85
4.3. Vey's semisimplicity theorem	86
4.4. The Vinberg metric	90
4.4.1. The Vinberg characteristic function	91
4.4.2. The covector field	94
4.4.3. The metric tensor	95
4.5. Benzécri's Compactness Theorem	98
4.5.1. Convex bodies in projective space	100
4.5.2. How convex bodies can degenerate	102
4.5.2.1. Corners	102
4.5.2.2. Flats	102
4.5.2.3. Osculating conics	102
4.5.3. Reduction to the affine case	104
4.6. Quasi-homogeneous and divisible domains	108
Part 2. Geometric manifolds	111
Chapter 5. Locally homogeneous geometric structures	113
5.1. Geometric atlases	114
5.1.1. The pseudogroup of local mappings	115
5.1.2. (G, X) -automorphisms	116
5.2. Development, holonomy	117
5.2.1. Construction of the developing map	118
5.2.2. Role of the holonomy group	121
5.2.3. Extending geometries	122
5.2.4. Simple applications of the developing map	122
5.3. The graph of a geometric structure	123
5.3.1. The tangent (G, X) -bundle	123
5.3.2. Developing sections	125
5.3.3. The associated principal bundle	125
5.4. The classification of geometric 1-manifolds	131
5.4.1. Compact Euclidean 1-manifolds and flat tori	132
5.4.2. Compact affine 1-manifolds and Hopf circles	133
5.4.2.1. Geodesics	133
5.4.3. Classification of projective 1-manifolds	134
5.4.4. Homogeneous affine structures	136
5.4.5. Grafting	142
Chapter 6. Examples of Geometric Structures	143
6.1. The hierarchy of geometries	144

6.1.1.	Enlarging and refining	144
6.1.2.	Cartesian products	146
6.1.3.	Fibrations	146
6.1.4.	Suspensions	147
6.2.	Parallel structures in affine geometry	148
6.2.1.	Flat tori and Euclidean structures	148
6.2.2.	Parallel suspensions	148
6.2.3.	Closed Euclidean manifolds	149
6.2.3.1.	A Euclidean \mathbb{Q} -homology 3-sphere	149
6.3.	Homogeneous subdomains	150
6.3.1.	Projective structures from non-Euclidean geometry	150
6.4.	Hopf manifolds	151
6.4.1.	Geodesics on Hopf manifolds	152
6.4.2.	The sphere of directions	153
6.4.3.	Hopf tori	153
6.5.	Radiant manifolds	156
6.5.1.	Radiant vector fields	156
6.5.2.	Radiant suspensions	159
6.5.2.1.	Radiant similarity manifolds	160
6.5.2.2.	Radiant affine surfaces	160
6.5.2.3.	Cross-sections to the radiant flow	162
Chapter 7.	Classification	165
7.1.	Marking geometric structures	165
7.1.1.	Marked Riemann surfaces	165
7.1.2.	Moduli of flat tori	166
7.1.3.	Marked geometric manifolds	166
7.1.4.	The infinitesimal approach	167
7.2.	Deformation spaces of geometric structures	167
7.2.1.	Historical remarks	169
7.3.	Representation varieties	169
7.3.1.	Example: $\mathrm{SL}(2)$ -characters of \mathbb{F}_2	171
7.3.1.1.	Coxeter Extension	173
7.3.1.2.	Hyperbolic three-holed spheres	174
7.3.2.	Twist flows and Fenchel-Nielsen earthquakes	175
7.4.	Fenchel-Nielsen coordinates on Fricke space	176
7.5.	Open manifolds	179
Chapter 8.	Completeness	183
8.1.	Locally homogeneous Riemannian manifolds	183
8.1.1.	Complete locally homogeneous Riemannian manifolds	184
8.1.2.	Topological rigidity of complete structures	185

8.1.3. Euclidean manifolds	186
8.2. Affine structures and connections	186
8.3. Completeness and convexity of affine connections	187
8.3.1. Review of affine connections	187
8.3.2. Geodesic completeness and the developing map	190
8.4. Unipotent holonomy	192
8.5. Complete affine structures on the 2-torus	194
8.6. Complete affine manifolds	198
8.6.1. The Bieberbach theorems	198
8.6.2. Complete affine solvmanifolds	199
Part 3. Affine and projective structures	203
Chapter 9. Affine structures on surfaces and the Euler characteristic	205
9.1. Benzécri's theorem on affine 2-manifolds	205
9.1.1. The surface as an identification space.	205
9.1.2. The turning number	208
9.1.3. The Milnor-Wood inequality	212
9.2. Higher dimensions	213
9.2.1. The Chern-Gauss-Bonnet Theorem	213
9.2.2. Smillie's examples of flat tangent bundles	216
9.2.3. The Kostant-Sullivan Theorem	218
Chapter 10. Affine structures on Lie groups and algebras	221
10.1. Étale representations and the developing map	222
10.1.1. Reduction to simply-connected groups	222
10.1.2. The translational part of the étale representation	223
10.2. Two-dimensional commutative associative algebras	224
10.3. Left-invariant connections and left-symmetric algebras	226
10.3.1. Bi-invariance and associativity	228
10.3.2. Completeness and right-nilpotence	230
10.3.3. Radiant vector fields	230
10.3.4. Volume forms and the characteristic polynomial	231
10.4. Two-dimensional noncommutative associative algebras	231
10.4.0.1. Haar measures and the characteristic polynomial	235
10.4.1. The opposite structure	235
10.4.2. A complete structure	236
10.4.3. The first halfspace family	237
10.4.3.1. Lorentzian structure	239
10.4.4. Parabolic deformations	239
10.4.4.1. Parabolic domains	240

10.4.4.2.	Simplicity	241
10.4.4.3.	Clans and homogeneous cones	241
10.4.5.	Deformations of the opposite structure	242
10.4.5.1.	Nonradiant deformation	242
10.4.5.2.	Radiant deformation	243
10.4.5.3.	Radiance and parallel volume	244
10.4.5.4.	Deforming to the complete structure	244
10.5.	Complete affine structures on closed 3-manifolds	244
10.5.1.	Central translations	245
10.6.	Fried's counterexample to Auslander's conjecture	248
10.7.	Solvable 3-dimensional algebras	251
10.8.	Incomplete affine 3-manifolds	252
10.8.1.	Parabolic cylinders	252
10.8.2.	Nonradiant deformations of radiant halfspace quotients	253
Chapter 11.	Parallel volume and completeness	257
11.1.	The volume obstruction	257
11.2.	Nilpotent holonomy	258
11.3.	Smillie's nonexistence theorem	260
11.4.	Fried's classification of closed similarity manifolds	263
11.4.1.	Completeness versus radiance	264
11.4.2.	Canonical metrics and incompleteness	264
11.4.3.	Incomplete geodesics recur	267
11.4.4.	Existence of halfspace neighborhood	271
11.4.5.	Visible points on the boundary of the halfspace	272
11.4.6.	The invisible set is affine and locally constant	272
Chapter 12.	Hyperbolicity	275
12.1.	The Kobayashi metric	276
12.2.	Kobayashi hyperbolicity	279
12.2.1.	Complete hyperbolicity and convexity	280
12.2.2.	The infinitesimal form	281
12.2.3.	Completely incomplete manifolds	286
12.3.	Hessian manifolds	289
Chapter 13.	Projective structures on surfaces	293
13.1.	Classification in higher genus	293
13.2.	Convex \mathbb{RP}^2 -structures	294
13.2.1.	Triangle groups	295
13.2.2.	Generalized Fenchel-Nielsen earthquakes	297
13.2.3.	Bulging deformations	298
13.3.	Coordinates for convex structures	300
13.3.1.	Fock-Goncharov coordinates	301

13.3.2. Affine spheres and Labourie-Loftin parametrization	302
13.4. Pathological developing maps and grafting	303
13.4.0.1. Smillie-Thurston example	304
Chapter 14. Complex-projective structures	307
14.1. Schwarzian parametrization	308
14.1.1. Affine structures and the complex exponential map	309
14.1.2. Projective structures and quadratic differentials	310
14.1.3. Solving the Schwarzian equation	312
14.2. Fuchsian holonomy	313
Chapter 15. Geometric structures on 3-manifolds	317
15.1. Higher dimensions: flat conformal and spherical CR-structures	317
15.2. Complete affine 3-manifolds	319
15.3. Margulis spacetimes	322
15.3.1. Affine boosts and crooked planes	322
15.3.2. Marked length spectra	324
15.3.3. Deformations of hyperbolic surfaces	325
15.3.4. Classification	327
15.4. Dupont's classification of hyperbolic torus bundles	328
Appendix A. Transformation groups	331
A.1. Group actions	331
A.2. Proper and syndetic actions	332
A.3. Topological transformation groupoids	333
Appendix B. Affine connections in local coordinates	335
B.1. Affine connections in local coordinates	335
B.2. Projective Equivalence	337
B.3. The (pseudo-) Riemannian connection	338
B.4. The Levi-Civita connection for the Poincaré metric	339
B.4.1. The usual coordinate system on \mathbb{H}^2	339
B.4.2. Connection 1-form for orthonormal frame	341
B.4.3. Geodesic curvature of a hypercycle	342
Appendix C. Facts about metric spaces	343
C.1. Nonincreasing homeomorphisms are isometries	343
C.2. Compactness of distance nonincreasing maps	344
C.3. The Lebesgue number of an open covering	344
Appendix D. Semicontinuous functions	347
D.1. Definitions and elementary properties	347
D.2. Approximation by continuous functions	347

Appendix E. $SL(2, \mathbb{C})$ and $O(3, 1)$	351
Bibliography	353
List of Figures	377
List of Tables	381
Index	383

Introduction

Symmetry powerfully unifies the various notions of geometry. Based on ideas of Sophus Lie, Felix Klein's 1972 Erlanger program proposed that geometry is the study of properties of a space X invariant under a group G of transformations of X . For example Euclidean geometry is the geometry of n -dimensional Euclidean space \mathbb{R}^n invariant under its group of rigid motions. This is the group of transformations which transforms an object ξ into an object congruent to ξ . In Euclidean geometry one can speak of points, lines, parallelism of lines, angles between lines, distance between points, area, volume, and many other geometric concepts. All these concepts can be derived from the notion of distance, that is, from the metric structure of Euclidean geometry. Thus any distance-preserving transformation or *isometry* preserves all of these geometric entities.

Notions more primitive than that of distance are the *length* and *speed* of a smooth curve. Namely, the distance between points a, b is the infimum of the length of curves γ joining a and b . The length of γ is the integral of its speed $\|\gamma'(t)\|$. Thus Euclidean geometry admits an infinitesimal description in terms of the *Riemannian metric tensor*, which allows a measurement of the size of the velocity vector $\gamma'(t)$. In this way standard Riemannian geometry generalizes Euclidean geometry by imparting Euclidean geometry to each tangent space.

Other geometries "more general" than Euclidean geometry are obtained by removing the metric concepts, but retaining other geometric notions. *Similarity geometry* is the geometry of Euclidean space where the equivalence relation of congruence is replaced by the broader equivalence relation of similarity. It is the geometry invariant under similarity transformations. In similarity geometry does not involve distance, but rather involves angles, lines and parallelism. *Affine geometry* arises when one speaks only of points, lines and the relation of parallelism. And when one removes the notion of parallelism and only studies lines, points and the relation of incidence between them (for example, three points being *collinear* or three lines being *concurrent*) one arrives at *projective geometry*. However in projective geometry, one must enlarge the space to *projective space*, which is the space upon which all the projective transformations are defined.

Here is a basic example illustrating the differences among the various geometries. Consider a particle moving along a smooth path; it has a well-defined velocity vector field (this uses only the differentiable structure of \mathbb{R}^n). In Euclidean geometry, it makes sense to discuss its "speed," so "motion at unit speed" (that is, "arc-length-parametrized

geodesic”) is a meaningful concept there. But in affine geometry, the concept of “speed” or “arc-length” must be abandoned: yet “motion at constant speed” remains meaningful since the property of moving at constant speed can be characterized as parallelism of the velocity vector field (zero acceleration). In projective geometry this notion of “constant speed” (or “parallel velocity”) must be further weakened to the concept of “projective parameter” introduced by J. H. C. Whitehead [283].

Synthetic projective geometry was developed by the architect Desargues in 1636–1639 out of attempts to understand the geometry of perspective. Two hundred years later non-Euclidean (hyperbolic) geometry was developed independently and practically simultaneously by Bolyai in 1833 and Lobachevsky in 1826–1829. These geometries were unified in 1871 by Klein who noticed that Euclidean, affine, hyperbolic and elliptic geometry were all “present” in projective geometry. Later in the nineteenth century, mathematical crystallography developed, leading to the theory of *Euclidean crystallographic groups*. Answering Hilbert’s eighteenth problem on the finiteness of the number of space groups in any given dimension n , Bieberbach developed a structure theory in 1918??. For torsionfree groups, the quotient spaces identified with *flat Riemannian manifolds* of dimension n , that is, Riemannian n -manifolds having zero sectional curvature. Such Riemannian structures are locally isometric to Euclidean space E^n . In particular, every point has an open neighborhood isometric to an open subset of E^n . These local isometries define a local Euclidean geometry on the neighborhood. Furthermore on overlapping neighborhoods, the local Euclidean geometries “agree,” that is, they are related by restrictions of global isometries $E^n \rightarrow E^n$. The neighborhoods form *coordinate patches*, the local isometries from the patches to E^n are the *coordinate charts*, and the restrictions of isometries of E^n are the corresponding *coordinate changes*. In this way a flat Riemannian manifold is defined by a coordinate atlas for a *Euclidean structure*.

More generally, for any geometry one can define geometric structures on a manifold M modeled on the homogeneous space (G, X) . A geometric atlas consists of an open covering of M by patches $U \hookrightarrow M$, a system of charts $U \xrightarrow{\psi} X$ such that the coordinate changes are locally restrictions of transformations of X which lie in G .

The plethora of different geometries suggests that, at least at a superficial level, no general inclusive theory of locally homogeneous geometric structures exists. Each geometry has its own features and idiosyncrasies, and special techniques particular to each geometry are

used in each case. For example, a surface modeled on \mathbb{CP}^1 has the underlying structure of a Riemann surface, and viewing a \mathbb{CP}^1 -structure as a projective structure on a Riemann surface provides a satisfying classification of \mathbb{CP}^1 -structures. Namely, as was presumably understood by Poincaré, *the deformation space of \mathbb{CP}^1 -structures on a closed surface Σ with $\chi(\Sigma) < 0$ identifies with a holomorphic affine bundle over the Teichmüller space of Σ* . When X is a complex manifold upon which G acts biholomorphically, holomorphic mappings provide a powerful tool in the study, a class of local mappings more flexible than “constant” maps (maps which are “locally in G ”) but more rigid than general smooth maps. Another example occurs when X admits a G -invariant connection (such as an invariant (pseudo-)Riemannian structure). Then the geodesic flow provides a powerful tool for the study of (G, X) -manifolds.

We emphasize the interplay between different mathematical techniques as an attractive aspect of this general subject. See [129] for a recent historical account of this material.

Organization of the text

The book divides into three parts. Part One describes affine and projective geometry and provides some of the main background on these extensions of Euclidean geometry. As noted by Lie and Klein, most classical geometries can be modeled in projective geometry. We introduce projective geometry as an extension of affine geometry, so we begin with a detailed discussion of affine geometry as an extension of Euclidean geometry and projective geometry as an extension of affine geometry. Part Two describes how to put the geometry of a Klein geometry (G, X) on a manifold M , and gives the basic examples and constructions. One goal is to *classify* the (G, X) -structures on a fixed topology in terms of a *deformation space* whose points correspond to equivalence classes of *marked structures*, whereby a marking is an extra piece of information which fixes the topology as the geometry of M varies. Part Three describes recent developments in this general theory of locally homogeneous geometric structures.

Part One: Affine and Projective Geometry

The first chapter introduces affine geometry as the geometry of parallelism. Two objects are *parallel* if they are related by a *translation*. Translations form a vector space \mathbf{V} , and act *simply transitively* on affine space. That is, for two points $p, q \in \mathbf{A}$ there is a unique translation taking p to q . In this way, points in \mathbf{A} identify with the vector space \mathbf{V} ,

but this identification depends on the (arbitrary) choice of a basepoint, or *origin* which identifies with the zero vector in \mathbf{V} . One might say that an affine space is a vector space, where the origin is forgotten. More accurately, the special role of the zero vector is suppressed, so that all points are regarded equally.

The action by translations now allows the definition of *acceleration* of a smooth curve. A curve is a *geodesic* if its acceleration is zero, that is, if its velocity is parallel. In affine space itself, unparametrized geodesics are straight lines; a parametrized geodesic is a curve following a straight line at “constant speed”. Of course, the “speed” itself is undefined, but the notion of “constant speed” just means that the acceleration is zero.

This notion of parallelism is a special case of the notion of an *affine connection*, except the existence of *globally defined* translations effecting the notion of parallelism is a special feature to our setting — the setting of *flat connections*. Just as Euclidean geometry is affine geometry with a parallel Riemannian metric, other linear-algebraic notions enhance affine geometry with parallel tensor fields. The most notable (and best understood) are flat Lorentzian (and pseudo-Riemannian) structures.

Chapter Two develops the geometry of projective space, viewed as the compactification of affine space. *Ideal points* arise as “where parallel lines meet.” A more formal definition of an ideal point is an equivalence class of lines, where the equivalence relation is parallelism of lines. Linear families (or *pencils*) of lines form planes, and indeed the set of ideal points in a projective space form a *projective hyperplane*, that is, a projective space of one lower dimension. Projective geometry appears when the ideal points lose their special significance, just as affine geometry appears when the zero vector $\mathbf{0}$ in a vector space loses its special significance.

However, we prefer a more efficient (if less synthetic) approach to projective geometry in terms of linear algebra. Namely, the *projective space associated to a vector space* \mathbf{V} is the space $\mathbf{P}(\mathbf{V})$ of 1-dimensional linear subspaces of \mathbf{V} (that is, lines in \mathbf{V} passing through $\mathbf{0}$). *Homogeneous coordinates* are introduced on projective space as follows. Since a 1-dimensional linear subspace is determined by any nonzero element, its coordinates determine a point in projective space. Furthermore the homogeneous coordinates are uniquely defined up to *projective equivalence*, that is, the equivalence relation defined by multiplication by nonzero scalars. Projectivizing linear subspaces of \mathbf{V} produces projective subspaces of $\mathbf{P}(\mathbf{V})$, and projectivizing linear automorphisms of \mathbf{V} yield *projective automorphisms*, or *collineations* of $\mathbf{P}(\mathbf{V})$.

The equivalence of the geometry of incidence in $P(V)$ with the algebra of V is remarkable. Homogeneous coordinates provide the “dictionary” between projective geometry and linear algebra. The collineation group is compactified as a projective space of “projective endomorphisms;” this will be useful for studying limits of sequences of projective transformations. These “singular projective transformations” are important in controlling developing maps of geometric structures, as developed in the second part.

The third chapter discusses, first from the classical viewpoint of polarities, the Cayley-Beltrami-Klein model for hyperbolic geometry. Polarities are the geometric version of nondegenerate symmetric or skew-symmetric bilinear forms on vector spaces. They provide a natural context for hyperbolic geometry, which is one of the principal examples of geometry in this study.

The Hilbert metric on a properly convex domain in projective space is introduced and is shown to be equivalent to the categorically defined Kobayashi metric [177, 179]. Later this notion is extended to manifolds with projective structure.

The fourth chapter develops notions of convexity. The Cayley-Beltrami-Klein metric on hyperbolic space is a special case of the Hilbert metric on properly convex domains. These define natural metric structures on certain well-studied projective structures. As an application of the Hilbert metric is Vey’s semisimplicity theorem, which is later used to characterize closed hyperbolic projective manifolds as quotients of sharp convex cones. Then another metric (due to Vinberg) is introduced, and is used to give a new proof of *Benzécri’s Compactness Theorem* that the collineation group acts properly and cocompactly on the space of convex bodies in projective space — in particular the quotient is a compact (Hausdorff) manifold. This is used to characterize the boundary of convex domains which cover convex projective manifolds. Recently Benzécri’s theorem has been used by Cooper, Long and Tillman [78] in their study of cusps of \mathbb{RP}^n -manifolds.

Part Two: Geometric Manifolds

The second part globalizes these geometric notions to manifolds, introducing *locally homogeneous geometric structures* in the sense of Whitehead [282] and Ehresmann [96] in the fifth chapter. We associate to every transformation group (G, X) a category of geometric structures on manifolds locally modeled on the geometry of X invariant under the group G . Because of the “rigidity” of the local coordinate

changes of open sets in X which arise from transformations in G , these structures on M intimately relate to the fundamental group $\pi_1(M)$.

Chapter 5 discusses three viewpoints for these structures. First we describe the coordinate atlases for the pseudogroup arising from (G, X) . Using the aforementioned rigidity, these are globalized in terms of a *developing map*

$$\widetilde{M} \xrightarrow{\text{dev}} X,$$

defined on the universal covering space \widetilde{M} of the geometric manifold M . The developing map is equivariant with respect to the holonomy homomorphism

$$\pi_1(M) \xrightarrow{h} G$$

which represents the group $\pi_1(M)$ of deck transformations of $\widetilde{M} \rightarrow M$ in G . Each of these two viewpoints represent M as a quotient: in the coordinate atlas description, M is the quotient of the disjoint union

$$\mathcal{U} := \coprod_{\alpha \in A} U_\alpha$$

of the coordinate patches U_α ; in the second description, M is represented as the quotient of \widetilde{M} by the action of the group $\pi_1(M)$. While a map defined on a connected space \widetilde{M} may seem more tractable than a map defined on the disjoint union \mathcal{U} , the space \widetilde{M} can still be quite large. The third viewpoint replaces \widetilde{M} with M and replaces the developing map by a section of a bundle defined over M . The bundle is a *flat bundle*, (that is, has *discrete structure group* in the sense of Steenrod). The corresponding *developing section* is characterized by transversality with respect to the foliation arising from the flat structure. This replaces the coordinate charts (respectively the developing map) being local diffeomorphisms into X .

Chapter 6 discusses examples of geometric manifolds from these three points of view. Although the main interest in these notes are structures modeled on affine and projective geometry, we describe other interesting structures.

All these structures are inter-related, because some geometries "contain" or "refine others." For example, affine geometry *contains Euclidean geometry*, when the metric notions are abandoned, but notions of parallelism are retained. This corresponds to the inclusion of the Euclidean isometry group as a subgroup of the affine automorphism group. Other examples include the projective and conformal models for non-Euclidean geometry. In these examples, the model space of the refined geometry is an open subset of the larger model space, and the

transformations in the refined geometry are restrictions of transformations in the larger geometry.

This *hierarchy of geometries* plays a crucial role in the theory. This is simply the geometric interpretation of the inclusion relations between closed subgroups of Lie groups. This *algebraicization* of classical geometries in the nineteenth century by Lie and Klein organized the proliferation of classical geometries. We adopt that point of view here. Indeed, we use this as a cornerstone in the construction and classification of geometric structures. The classification of geometric manifolds often shows that a manifold modeled on one geometry may actually have a *stronger* geometry. For example, Fried's theorem (discussed in §11.4) shows that a closed manifold with a similarity structure is either a Euclidean manifold or a manifold on $\mathbf{E}^n \setminus \{\mathbf{0}\} \cong S^{n-1} \times \mathbb{R}$ with its invariant (product) Riemannian metric.

Chapter 7 deals with the general classification of (G, X) -structures from the point of view of developing sections. The main result is an important observation due to Thurston [265] that the *deformation space* of marked (G, X) -structures on a fixed topology Σ is itself “locally modeled” on the quotient of the space $\text{Hom}(\pi_1(\Sigma), G)$ by the group $\text{Inn}(G)$ of inner automorphisms of G . The description of $\mathbb{R}P^1$ -manifolds is described in this framework.

Chapter 8 deals with the important notion of *completeness*, for taming the developing map. In general, the developing map may be quite pathological — even for closed (G, X) -manifolds — but under various hypotheses, can be proved to be a covering space onto its image. However, the main techniques borrow from Riemannian geometry, and involves *geodesic completeness* of the Levi-Civita connection (the Hopf-Rinow theorem). As an example, we classify complete affine structures on the 2-torus (due to Kuiper). The Hopf manifolds introduced in §6.4 are fundamental examples of incomplete structures. That affine structures on compact manifolds are generally incomplete is one dramatic difference between affine geometry and traditional Riemannian geometry.

This requires, of course, relating geometric structures to *connections*, since all of the locally homogeneous geometric structures discussed in this book can be approached through this general concept. However, we do *not* discuss the general notion of *Cartan connections*, but rather refer to the excellent introduction to this subject by R. Sharpe [249].

Part Three: Affine and projective structures

Chapter 9 begins the classification of affine structures on surfaces. We prove Benzécri’s theorem [34] that a closed surface Σ admits an affine structure if and only if its Euler characteristic vanishes. We discuss the famous conjecture of Chern that the Euler characteristic of a closed affine manifold vanishes. Following Kostant-Sullivan [182] we prove this in the *complete* case. Chern’s conjecture has recently been proved in the volume-preserving case by Klingler [175].

Chapter 10 offers a detailed study of left-invariant affine structures on Lie groups. These provide many examples; in particular all the non-radiant affine structures on T^2 are *invariant* affine structures on the *Lie group* T^2 . For these structures the holonomy homomorphism and the developing map blend together in an intriguing way, and this perhaps this provides a conceptual basis for the unexpected relation between the one-dimensional property of geodesic completeness and the top-dimensional property of volume-preserving holonomy. Covariant differentiation of left-invariant vector fields lead to well-studied non-associative algebras *algèbres symétriques à gauche* (*left-symmetric algebras*, so defined as their associators are symmetric in the left two arguments. Commutator defines the structure of an underlying Lie algebra. Associative algebras correspond to *bi-invariant affine structures*, so the “group objects” in the category of affine manifolds correspond naturally to associative algebras.¹ As these structures were introduced by Ernest Vinberg [278] in his study of homogeneous convex cones in affine space, and further developed by Jean-Louis Koszul and his school, we call these algebras *Koszul-Vinberg algebras*. We take a decidedly geometric approach to these ubiquitous mathematical structures.

Most closed affine surfaces are invariant affine structures on the torus *group*.

Chapter 11 describes the question (apparently first raised by L. Markus [209]) of whether, for an closed orientable affine manifold, completeness is equivalent to *parallel volume*. The existence of a parallel volume form is equivalent to unimodularity of the linear holonomy group, that is, whether the holonomy is volume-preserving. This tantalizing question has led to much research, and subsumes various questions which we discuss. Carrière’s proof that compact flat Lorentzian manifolds are complete [56] is a special case of this conjecture. In

¹Apparently this is due to Vinberg, but I wonder if this was known earlier, perhaps to Cartan or Ehresmann.

particular we give the sharp classification of closed similarity manifolds by D. Fried [106] (a much different proof is independently due to Vaisman-Reischer [270]). The analog of this question for left-invariant affine structures on Lie groups is the conceptual and suggestive result that completeness is equivalent to parallelism of *right-invariant* vector fields, proved in §10.3.4.

Chapter 12 expounds the notions of “hyperbolicity” of Vey [275] and Kobayashi [179]. *Hyperbolic affine manifolds* are quotients of properly convex cones. Compact such manifolds are radiant suspensions of \mathbb{RP}^n -manifolds which are quotients of divisible domains. In particular we describe how a *completely incomplete* closed affine manifold must be affine hyperbolic in this sense. (That is, we tame the developing map of an affine structure with *no* two-ended complete geodesics.) This striking result is similar to the tameness where *all* geodesics are complete — complete manifolds are also quotients.

Chapter 13 summarizes some aspects of the now blossoming subject of \mathbb{RP}^2 -structures on surfaces, in terms of the explicit coordinates and deformations which extend some of the classic geometric constructions on the deformation space of hyperbolic structures on closed surfaces.

Chapter 14 describes the classic subject of \mathbb{CP}^1 -manifolds, which traditionally identify with *projective structures on Riemann surfaces*. Using the Schwarzian derivative, these structures are classified by the points of a holomorphic affine bundle over the Teichmüller space of Σ . This parametrization (presumably known to Poincaré), is remarkable in that it is completely *formal*, using standard facts from the theory of Riemann surfaces. One knows precisely the deformation space without any knowledge of the developing map (besides it being a local biholomorphism). This is notable because the developing maps can be pathological; indeed the first examples of pathological developing maps were \mathbb{CP}^1 -manifolds on hyperbolic surfaces. The theory of projective structures on Riemann surfaces is a paradigm for a successful classification of highly nontrivial geometric structures, and is a suggestive paradigm for the study of geometric structures in higher dimensions.

Chapter 15 surveys known results, and the many open questions, in dimension three. This complements Thurston’s book [266] and expository articles of Scott [246] and Bonahon [43], which deal with geometrization and the relations to 3-manifold topology. In particular we describe the classification (due to Serge Dupont) of projective structures on hyperbolic torus bundles

Acknowledgements

This book grew out of lecture notes *Projective Geometry on Manifolds*, from a course at the University of Maryland in 1988. Since then I have given minicourses at international conferences, graduate courses at the University of Maryland and Brown University in Fall 2017, which have been extremely useful in developing the material here. Courses I gave at the International Center for Theoretical Physics in Trieste, Lisbon (lectures in geometry), and a minicourse at University of Texas in Austin were also helpful in organizing this material.

I am grateful for notes written by Son Lam Ho and Greg Laun in later iterations of this course at Maryland.

Comments and suggestions by Suhyoung Choi, Daryl Cooper, Jeff Danciger, Darren Long, Ludovic Marquis, Fanny Kassel, Stephan Tillmann, and various anonymous referees have been extremely helpful. Recently I have benefitted from comments by Charles Daly, Misha Kapovich, Daniele Alessandrini and Rich Schwartz at various times in the development of this manuscript.

I am extremely grateful to Ed Dunne, Ina Mettes, and Eriko Hirone for many informative suggestions, comments and encouragement of this project.

My research in this subject began in my undergraduate thesis [114] written in Princeton in 1977. Dennis Sullivan and Bill Thurston suggested the study of affine and projective structures, which I continued to pursue in graduate school in Berkeley with Moe Hirsch. David Fried and John Smillie were spending summers there and we discussed affine structures extensively. Correspondence with Jacques Vey and Jacques Helmstetter in Grenoble, and extensive conversations with Dan Burns, Shiing-Shen Chern, Robert Gunning, Ravi Kulkarni, Shoshichi Kobayashi, Joe Wolf, and S.T. Yau were also extremely informative.

While a postdoc, I benefitted from many discussions with Scott Wolpert on symplectic geometry of Fricke space, from John Millson and Domingo Toledo on deformation theory of discrete groups. My collaboration with Millson was influenced by the generosity of Pierre Deligne, who, in an unpublished letter to Millson, taught us, among other things, the power of the scheme-theoretic viewpoint.

I also want to express my gratitude to François Labourie for many discussions at an early stage, as well as Yves Benoist for many interesting conversations and his support of this project, beginning with my 1988 notes. I am also grateful to Grisha Margulis for many discussions about proper affine actions, and conversations with John Loftin,

John Parker and Elisha Falbel about affine, projective and spherical CR geometry.

I have benefitted from many former graduate students working on various aspects of this theory, including experimental and visualization efforts. Ser Peow Tan, Todd Drumm, Virginie Charette, Sean Lawton, Andrew Sanders, and Jean-Philippe Burelle have been particularly influential in my understanding of this subject.

The series of conferences *Crystallographic Groups and their Generalizations* (1996, 1999, 2002, 2005, 2008, 2011), in Kortrijk, Belgium (and later in Oostende) were also very formative, and I am extremely grateful to Paul Igodt and Karel Dekimpe of the Katholieke University of Louvain in Kortrijk for organizing these highly stimulating events.

There are many people with whom conversations and correspondence have been useful in developing this theory. I have benefitted tremendously from a large community of research mathematicians at various career stages. I sincerely apologize for any omissions.

Work partially supported by National Science Foundation grants DMS070781, DMS1709701 a Semester Research Awards (Fall 2005 and 2010) from the General Research Board of the University of Maryland, and the Oswald Veblen Fund at the Institute for Advanced Study. I am grateful to the Simons Foundation for a Simons sabbatical grant for 2014-2015, and

The GEAR network, funded through the National Science Foundations's *Research Networks in the Mathematical Sciences* program directed by Steven Bradlow (main director), myself, Steven Kerckhoff, Richard Wentworth and Anna Wienhard, has facilitated much activity in this subject, which is reflected in this text.

I also wish to express my gratitude to the Mathematical Sciences Research Institute, the Institute for Advanced Study, Brown University and the Institute for Computation and Experimentation in Research in Mathematics, Institut Henri Poincaré, the Simons Foundation and the National Science Foundation their generous support and hospitality for this research.

Finally I would like to thank the enormous and deep support of my family: especially my wife Emily, offspring Evan, Lizzie and Michael, and grandchildren Amelia, Liana, Leonardo and Jonah, to whom this book is dedicated.

Prerequisites

This book is aimed roughly at first-year graduate students and advanced undergraduate students, although some knowledge of advanced material will be useful.

For general treatments of geometry, we refer to the two-volume text of Berger [39, 38] (see also Berry-Pansu-Berry-Saint Raymond [40]) and Coxeter [80].

We also assume basic familiarity with elementary topology, smooth manifolds, and the rudiments of Lie groups and Lie algebras. Much of this can be found in Lee’s book “Introduction to Smooth Manifolds” [201], including its appendices. For topology, we require basic familiarity with the notion of metric spaces, covering spaces and fundamental groups.

Fiber bundles, as discussed in the still excellent treatise of Steenrod [259], or the more modern treatment of principal bundles given in Sontz [257], will be used.

Some familiarity with the properties of proper maps and proper group actions will also be useful.

Some familiarity with the theory of connections in fiber bundles and vector bundles is useful, for example, Kobayashi-Nomizu [181], or Milnor [222], do Carmo [87] Lee [200], O’Neill [230].

We put the discussion of Fenchel-Nielsen coordinates on Fricke space in the context of Darboux’s theorem in symplectic geometry; we recommend §22 of Lee [201], §22 for a good general treatment consistent with our notation.

Notation, terminology and general background

In this section we collect various notational and terminological conventions, as well as some basic material which we use throughout.

Vectors and matrices. We work over a field \mathbf{k} , usually the field \mathbb{R} of real numbers, but sometimes the field \mathbb{C} of complex numbers. We shall denote vectors and matrices in bold font. Let \mathbf{V} be a vector space over \mathbf{k} of dimension n . A vector in \mathbf{V} corresponds to a column vector:

$$\mathbf{v} \longleftrightarrow \begin{bmatrix} v^1 \\ \vdots \\ v^n \end{bmatrix}$$

A *covector* is defined as a linear functional $\mathbf{V} \xrightarrow{\omega} \mathbf{k}$, corresponding to a row vector:

$$\omega \longleftrightarrow [\omega_1 \ \dots \ \omega_n]$$

and the duality pairing between V and V^* is:

$$\begin{aligned} V \times V^* &\longrightarrow \mathbf{k} \\ (\mathbf{v}, \omega) &\longmapsto v^i \omega_i \end{aligned}$$

(summation over paired indices). A linear transformation $\mathbf{k}^m \longrightarrow \mathbf{k}^n$ is defined by an $m \times n$ matrix

$$\mathbf{A} = [A^i_j]$$

mapping

$$\begin{aligned} \mathbf{k}^m &\xrightarrow{\mathbf{A}} \mathbf{k}^n \\ \mathbf{v} = \begin{bmatrix} v^1 \\ \vdots \\ v^m \end{bmatrix} &\longmapsto \begin{bmatrix} A^1_j v^j \\ \vdots \\ A^n_j v^j \end{bmatrix} \end{aligned}$$

where $j = 1, \dots, m$.

Affine vector fields on A correspond to affine maps $A \rightarrow A$:

$$\mathbf{A} := (A^i_j x^j + a^i) \partial_i \quad \longleftrightarrow \quad \hat{\mathbf{A}} := [A \mid a]$$

where

$$\mathbf{A} = \begin{bmatrix} A^1_1 & \dots & A^1_i & \dots & A^1_n \\ \vdots & & \vdots & & \vdots \\ A^i_1 & \dots & A^i_j & \dots & A^i_n \\ \vdots & & \vdots & & \vdots \\ A^n_1 & \dots & A^n_j & \dots & A^n_n \end{bmatrix}$$

is the linear part and

$$a = \begin{bmatrix} a^1 \\ \vdots \\ a^i \\ \vdots \\ a^n \end{bmatrix}$$

is the translational part. In this notation,

$$(1) \quad \hat{\mathbf{A}} = [A \mid a] = \left[\begin{array}{cccc|c} A^1_1 & \dots & A^1_n & & a^1 \\ \vdots & & \vdots & & \vdots \\ \dots & A^i_j & \dots & & a^i \\ \vdots & & \vdots & & \vdots \\ A^n_1 & \dots & A^n_n & & a^n \end{array} \right]$$

Projective equivalence. Denote the multiplicative group of nonzero scalars in \mathbf{k} by \mathbf{k}^\times , and let W be a vector space over \mathbf{k} . Then \mathbf{k}^\times acts by scalar multiplication on W . Say that nonzero vectors $\mathbf{w}, \mathbf{u} \in W$ are *projectively equivalent* if and only if $\exists \lambda \in \mathbf{k}^\times$ such that $\mathbf{w} = \lambda \mathbf{u}$. Projective equivalence classes $[\mathbf{v}]$ of nonzero vectors \mathbf{v} form the *projective space* $P(V)$ associated to V . Denote the projective equivalence class of

a vector $\mathbf{v} = \begin{bmatrix} v^1 \\ \vdots \\ v^n \end{bmatrix} \in V$ by

$$[\mathbf{v}] := \left[\begin{bmatrix} v^1 \\ \vdots \\ v^n \end{bmatrix} \right]$$

and the projective equivalence class of a covector $\omega = [\omega^1 \dots \omega^n] \in V^*$ by

$$[\omega] := \llbracket \omega^1 \quad \dots \quad \omega^n \rrbracket.$$

The set of projective equivalence classes of nonzero vectors in W is the *projective space* associated to W and denoted $P(W)$. (Projective equivalence classes of nonzero *covectors* comprise the projective space $P(W^*)$ *dual* to $P(W)$).

General Topology. For general background in topology we refer to Lee [201] and Willard [285].

If A is a topological space, and $B \subset A$ is a subspace, then we write $B \subset\subset A$ if B is compact (in the subspace topology).

We denote the space of mappings $A \rightarrow B$ by $\text{Map}(A, B)$, given the compact-open topology.

If f_n (for $n = 1, 2, \dots, \infty$) are mappings on a space X , write $f_n \rightrightarrows f_\infty$ if f_n converges *uniformly* to f_∞ on X .

Denote the group of diffeomorphisms of a smooth manifold X by $\text{Diff}(X)$, with the C^∞ topology (uniform convergence to all orders, on all $K \subset\subset X$). If f, g are smooth maps between smooth manifolds $X \rightarrow Y$, then we say that f and g are *isotopic* if and only if there is a smooth path

$$\phi_t \in \text{Diff}(X), \quad 0 \leq t \leq 1,$$

with $\phi_0 = \mathbb{I}_X$ such that $g = \phi_1 \circ f$. Denote this relation by $f \simeq g$.

Suppose (X, \mathbf{d}) is a metric space. If $x \in X, r > 0$, define the (*open*) *ball* with center x and radius r as:

$$B_r(x) := \{y \in X \mid \mathbf{d}(x, y) < r\}.$$

The open balls in a metric space are partially ordered by inclusion. More generally, if $A \subset X$, define

$$B_r(A) := \{y \in X \mid \exists a \in A \text{ such that } d(x, a) < r\}.$$

If (X, d) is a metric space, and $S, T \subset \subset X$, then define their *Hausdorff distance*

$$d(S, T) := \inf\{r \in \mathbb{R} \mid S \subset B_r(T) \text{ and } T \subset B_r(S)\}.$$

If X is compact, then the set of closed subsets of X with Hausdorff distance d is a metric space.

Denote group of isometries of a metric space (X, d) by $\text{Isom}(X, d)$, or just $\text{Isom}(X)$ if the context is clear.

Fundamental group and covering spaces. If $[a, b] \xrightarrow{\gamma} X$ is a continuous path, write

$$\gamma(a) \overset{\gamma}{\rightsquigarrow} \gamma(b)$$

to indicate that γ runs between its two endpoints $\gamma(a), \gamma(b)$. Fix an arbitrary basepoint $p_0 \in X$. A *loop based at p_0* is a path $p_0 \overset{\gamma}{\rightsquigarrow} p_0$. Two such *based loops* γ_1, γ_2 are *relatively homotopic* if they are homotopic by a homotopy which fixes p_0 . In that case we write $\gamma_1 \simeq \gamma_2$. In that The *fundamental group* $\pi_1(M; p_0)$ corresponding to p_0 consists of relative homotopy classes $[\gamma]$ of based loops γ .

The group operation is defined by *concatenation* of paths: If

$$[a_i, b_i] \xrightarrow{\gamma_i} X, \text{ for } i = 1, 2$$

are paths, with $\gamma_1(b_1) = \gamma_2(a_2)$, write $\gamma_1 \star \gamma_2$ for the continuous path

$$\gamma_1(a_1) \rightsquigarrow \gamma_2(b_2),$$

defined by:

$$[a_1, b_2] \xrightarrow{\gamma_1 \star \gamma_2} X$$

$$t \longmapsto \begin{cases} \gamma_1(t) & \text{if } a_1 \leq t \leq b_1 \\ \gamma_2(t) & \text{if } a_2 \leq t \leq b_2 \end{cases}$$

If γ_1, γ_2 are loops based at p_0 , so is $\gamma_1 \star \gamma_2$, and concatenation defines an binary operation on $\pi_1(X, p_0)$.

The *constant path* p_0 defines an identity element on $\pi_1(X, p_0)$ since $p_0 \star \gamma \simeq \gamma \star p_0 \simeq \gamma$. Define the *inverse* of a path $[a, b] \xrightarrow{\gamma} M$

$$[a, b] \xrightarrow{\gamma^{-1}} M$$

$$t \longmapsto \gamma(a + b - t).$$

If γ is a loop based at p_0 , then

$$\gamma \star \gamma^{-1} \simeq \gamma^{-1} \star \gamma \simeq p_0,$$

obtaining *inversion* in $\pi_1(M; p_0)$. If $[a_3, b_3] \xrightarrow{\gamma_3} X$ with $\gamma_2(b_2) = \gamma_3(a_2)$, then

$$(\gamma_1 \star \gamma_2) \star \gamma_3 \simeq \gamma_1 \star (\gamma_2 \star \gamma_3),$$

implying associativity. Thus $\pi_1(X, p_0)$ is indeed a group.

Under rather general conditions on X (such as being a topological manifold) defined the *universal covering space* (corresponding to p_0)

$$\widetilde{X^{(p_0)}} \xrightarrow{\Pi} X$$

as the collection of relative homotopy classes of paths γ starting at p_0 , with the other endpoint at $\Pi(\gamma)$. Give $\widetilde{X^{(p_0)}}$ the coarsest topology such that Π is continuous. Then Π is a local homeomorphism, and indeed a *Galois covering space* with covering group $\pi_1(X, p_0)$.

The (left) action on $\widetilde{X^{(p_0)}}$ by deck transformations from $\pi_1(X, p_0)$ is defined as follows. Choose a point $p \in X$, a path $p_0 \xrightarrow{\eta} p$ and a loop γ based at p_0 . The action of $[\gamma]$ on $[\eta]$ is defined by:

$$[\eta] \xrightarrow{[\gamma]} [\gamma \star \eta].$$

The action is free and proper, preserves $p = \Pi([\eta])$. The quotient map naturally identifies with Π and the quotient space $\widetilde{X^{(p_0)}}/\pi_1(X, p_0)$ naturally identifies with X .

Smooth manifolds. We shall work in the context of smooth manifolds, for which a good general reference is Lee [200]. This will enable the use of differential calculus locally, and notions of smooth mappings between manifolds. A *smooth manifold* is a Hausdorff space built from open subsets of \mathbb{R}^n , which we call *coordinate patches*. The *coordinate changes* are general smooth locally invertible maps. If M and N are given such structures, a continuous map $M \rightarrow N$ is *smooth* if in the local coordinate charts it is given by a smooth map.

This structure enables the *tangent bundle* $\mathbb{T}M$, whose points are the *infinitesimal displacements* of points in M . That is, to every smooth curve $(a, b) \xrightarrow{\gamma} M$, and parameter t with $a \leq t \leq b$, is a *velocity vector*

$$\gamma'(t) \in \mathbb{T}_{\gamma(t)}M$$

representing the infinitesimal effect of displacing $\gamma(t)$ along γ . Since the local coordinates change by general smooth locally invertible maps, there is no natural way of identifying these infinitesimal displacements at *different* points. Therefore we attach to each point $p \in M$, a “copy”

$\mathbb{T}_p M$ of the model space \mathbb{R}^n , which represents the vector space of *infinitesimal displacements of p* . It is important to note that although the *fibers* $\mathbb{T}_p M$ are disjoint, that the union

$$\mathbb{T}M := \bigcup_{p \in M} \mathbb{T}_p M$$

is topologized as a smooth manifold (indeed, a smooth *vector bundle*), and not as the disjoint union.

The velocity vector of a smooth curve is a *tangent vector at p* , which can be defined in two equivalent ways:

- Equivalence classes of smooth curves $\gamma(t)$ with $\gamma(0) = p$, where curves $\gamma_1 \sim \gamma_2$ if and only if

$$\left. \frac{d}{dt} \right|_{t=0} f \circ \gamma_1(t) = \left. \frac{d}{dt} \right|_{t=0} f \circ \gamma_2(t)$$

for all smooth functions $U \xrightarrow{f} \mathbb{R}$, where $U \subset \mathbf{A}$ is an open neighborhood of p .

- Linear operators $C^\infty(\mathbf{A}) \xrightarrow{D} \mathbb{R}$ satisfying

$$(2) \quad D(fg) = D(f)g(p) + f(p)D(g).$$

The tangent space $\mathbb{T}_p M$ is a vector space *linearizing* the smooth manifold M at the point $p \in M$.

The space of tangent vectors forms a smooth vector bundle $\mathbb{T}M \xrightarrow{\Pi} M$, with fiber $\Pi^{-1}(p) := \mathbb{T}_p M$. If $U \ni p$ is a coordinate patch, then $\Pi^{-1}(U)$ identifies with $U \times \mathbb{R}^n$, and this defines a smooth coordinate atlas on $\mathbb{T}M$.

Let M, N be smooth manifolds, and $p \in M$. A mapping

$$M \xrightarrow{f} N$$

is *differentiable at p* if every infinitesimal displacement $\mathbf{v} \in \mathbb{T}_p M$ maps to an infinitesimal displacement $D_p f(\mathbf{v}) \in \mathbb{T}_q N$, where $q = f(p)$. That is, if γ is a smooth curve with $\gamma(0) = p$ and $\gamma'(0) = \mathbf{v}$, then we require that $f \circ \gamma$ is a smooth curve through q at $t = 0$; then we call the new velocity $(f \circ \gamma)'(0) \in \mathbb{T}_q N$ the value of the *differential* or *derivative*

$$\begin{aligned} \mathbb{T}_p M &\xrightarrow{(Df)_p} \mathbb{T}_q N \\ \mathbf{v} &\longmapsto (f \circ \gamma)'(0) \end{aligned}$$

If P is a third smooth manifold, and $N \xrightarrow{g} P$ is a smooth map, the composition $M \xrightarrow{g \circ f} P$ is defined, and is a smooth map. The *Chain Rule* expresses the derivative of the composition as the composition of

the derivatives of f and g : and $M \xrightarrow{f} N \xrightarrow{g} P$ are smooth maps, then the differential of a composition

$$\begin{array}{ccccc} & & g \circ f & & \\ & \curvearrowright & & \curvearrowleft & \\ M & \xrightarrow{f} & N & \xrightarrow{g} & P \end{array}$$

induces a commutative diagram

$$\begin{array}{ccccc} & & (\mathbb{D}(g \circ f))_x & & \\ & \curvearrowright & & \curvearrowleft & \\ \mathbb{T}_x M & \xrightarrow{(\mathbb{D}f)_x} & \mathbb{T}_{f(x)} N & \xrightarrow{(\mathbb{D}g)_{f(x)}} & \mathbb{T}_{(g \circ f)(x)} P \end{array}$$

that is, $\mathbb{D}(g \circ f)_x = (\mathbb{D}g)_{f(x)} \circ (\mathbb{D}f)_x$.

If M, N are smooth manifolds, a *diffeomorphism* $M \rightarrow N$ is an invertible smooth mapping whose inverse is also smooth. In particular a diffeomorphism is a homeomorphism. If $M \xrightarrow{f} N$ is a smooth map and $p \in M$ such that the differential

$$\mathbb{T}_p M \xrightarrow{(\mathbb{D}f)_p} \mathbb{T}_{f(p)} N$$

is an isomorphism of vector spaces, the *Inverse Function Theorem* implies the existence of an open neighborhood $U \ni p$ such that the restriction $f|_U$ is a diffeomorphism $U \rightarrow f(U)$. In particular $f(U) \subset N$ is open and U can be chosen so that $(\mathbb{D}f)_q$ is an isomorphism for every $q \in U$. Such a map is called a *local diffeomorphism* (at p).

Under the C^∞ topology, diffeomorphisms $M \rightarrow M$ form a topological group, denoted by $\text{Diff}(M)$. Indeed $\text{Diff}(M)$ has more structure as a *Fréchet Lie group*. If N is a smooth manifold, then a map $N \rightarrow \text{Diff}(M)$ is *smooth* if the natural composition $N \times M \rightarrow M$ is smooth. A smooth homomorphism $\mathbb{R} \xrightarrow{\Phi} \text{Diff}(M)$ is called a *smooth flow* on M .

Vector fields. A vector field on M is a section of the tangent bundle $\mathbb{T}M \xrightarrow{\Pi} M$, that is a mapping $M \xrightarrow{\xi} \mathbb{T}_p M$ such that

$$\Pi \circ \xi = \mathbb{I}_M,$$

or equivalently, $\xi(p) \in \mathbb{T}_p M$. Denote the space of all vector fields on M by $\text{Vec}(M)$. Just as individual tangent vectors at $p \in M$ define derivations $C^\infty(M) \rightarrow \mathbb{R}$ over the evaluation map

$$\begin{array}{l} C^\infty(M) \longrightarrow \mathbb{R} \\ f \longmapsto f(p) \end{array}$$

(as in (2)), vector fields in $\text{Vec}(M)$ define derivations of the algebra $C^\infty(M)$.

Let $a < b \in \mathbb{R}$. A smooth curve $(a, b) \xrightarrow{\gamma} M$ is an *integral curve* for $\xi \in \text{Vec}(M)$ if and only if

$$\gamma'(t) = \xi(\gamma(t)) \in \mathbb{T}_{\gamma(t)}M$$

for all $a < t < b$. If Φ is a smooth flow as above, then for each $p \in M$,

$$\begin{aligned} (a, b) &\xrightarrow{\Phi_p} M \\ t &\longmapsto \Phi(t)(p) \end{aligned}$$

is a smooth curve in M with velocity vector field $(\Phi_p)'(t) \in \mathbb{T}_{\Phi_p(t)}M$. In particular

$$\xi(p) := (\Phi_p)'(0) \in \mathbb{T}_pM \text{ (since } \Phi_p(0) = p)$$

defines a smooth vector field $\xi \in \text{Vec}(M)$.

The *Fundamental Theorem on Flows* is a statement in the converse direction: every vector field $\xi \in \text{Vec}(M)$ is tangent to a *local flow*. That is, through every point there exists a unique maximal integral curve, defined for some open interval (a, b) containing 0. When M is a closed manifold), then the integral curves are defined on all of \mathbb{R} and corresponds to a flow Φ on M . Such a vector field is called *complete*. More generally (Lee [201], Theorem 9.16), if ξ is compactly supported, it is complete.

See Lee [201], §9, for full details; a precise statement of the Fundamental Theorem on Flows is given in Theorem 9.12. If f is a *local diffeomorphism*, and $\xi \in \text{Vec}(N)$, then define the *pullback* $f^*\xi \in \text{Vec}(M)$ by:

$$(3) \quad (f^*\xi)_p := ((Df)_p)^{-1}(\xi_{f(p)}).$$

In particular, in the terminology of Lee [201], the vector fields ξ and $f^*\xi$ are *f-related*.

Suppose that $M \xrightarrow{f} N$ is a smooth map and $\xi \in \text{Vec}(M)$ and $\eta \in \text{Vec}(N)$ are *f-related* vector fields, that is,

$$(Df)_p(\xi(p)) = \eta(f(p)), \quad \forall p \in M.$$

The *Naturality of Flows* (Lee [201], Theorem 9.13) implies that if $\Psi(t)$ is the local flow defined by $\xi \in \text{Vec}(M)$ and $\Psi(t)$ the local flow on N defined by $\eta \in \text{Vec}(N)$, then

$$f(\Phi_t(p)) = \Psi_t(f(p))$$

whenever these objects are defined.

The vector fields on M form a Lie algebra $\text{Vec}(M)$ under Lie bracket.

Part 1

Affine and projective geometry

CHAPTER 1

Affine geometry

This section introduces the geometry of affine spaces. After a rigorous definition of affine spaces and affine maps, we discuss how linear algebraic constructions define geometric structures on affine spaces. Affine geometry is then transplanted to manifolds. The section concludes with a discussion of affine subspaces, vector fields, volume and the notion of center of gravity.

1.1. Euclidean space

We begin with a short summary of Euclidean geometry in terms of its underlying space and its group of isometries.

Euclidean geometry can be described in many different ways. Here is one simple approach. Denote by \mathbf{E}^n the set of points in the vector space \mathbb{R}^n (that is, ordered n -tuples of real numbers) with the distance function

$$\begin{aligned} \mathbf{E}^n \times \mathbf{E}^n &\xrightarrow{d} \mathbb{R} \\ (p, q) &\longmapsto \|p - q\|. \end{aligned}$$

EXERCISE 1.1.1. Let $(\mathbf{E}^n, d) \xrightarrow{g} (\mathbf{E}^n, d)$ be an isometry. Then

$$g(p) = \mathbf{A}p + \mathbf{b}$$

for an orthogonal matrix $\mathbf{A} \in \mathbf{O}(n)$ and a vector $\mathbf{b} \in \mathbb{R}^n$.

That is, the isometry g of Euclidean n -space \mathbf{E}^n is a composition of the linear isometry defined by \mathbf{A} and the translation

$$p \xrightarrow{\tau_{\mathbf{b}}} p + \mathbf{b}$$

by \mathbf{b} .

Two objects X, Y are *parallel* if they are related by the action of a translation, in which case we write $X \parallel Y$.

EXERCISE 1.1.2. Show that translations form a normal subgroup $\text{Trans}(\mathbf{E}^n)$ isomorphic to \mathbb{R}^n and

$$\text{Isom}(\mathbf{E}^n) = \text{Trans}(\mathbf{E}^n) \rtimes \mathbf{O}(n).$$

Deduce that $\text{Isom}(\mathbf{E}^n)$ preserves the relation of parallelism.

Another feature of Euclidean geometry is the notion of *angle*:

EXERCISE 1.1.3. *Every isometry of \mathbf{E}^n preserves angles. (Hint: use the fact that angles can be defined in terms of inner products on \mathbb{R}^n .)*

That is, every isometry is angle-preserving, or *conformal*. The equivalence relation of *similarity* is generated by the group $\text{Sim}(\mathbf{E}^n)$ of conformal transformations of \mathbf{E}^n .

An element of $\text{Sim}(\mathbf{E}^n)$ which is not an isometry is the *homothety* given by scalar multiplication $p \mapsto \lambda p$, where $\lambda \in \mathbb{R}$ and $\lambda \neq \pm 1$. (See §1.5.2 for the general definition of homotheties.) Denote the group of scalar multiplications by $\lambda > 0$ by \mathbb{R}^+ .

EXERCISE 1.1.4. *The group $\text{Sim}(\mathbf{E}^n)$ is generated by $\text{Isom}(\mathbf{E}^n)$ and \mathbb{R}^+ . Indeed,*

$$\text{Sim}(\mathbf{E}^n) = \text{Sim}_0(\mathbf{E}^n) \ltimes \mathbb{R}^+$$

where

$$\text{Sim}_0(\mathbf{E}^n) := \mathbb{R}^+ \times \text{O}(n)$$

is the group of linear similarities of \mathbf{E}^n . Explicitly a transformation g of \mathbf{E}^n lies in $\text{Sim}(\mathbf{E}^n)$ if it has the form

$$p \mapsto \lambda \mathbf{A}(p) + \mathbf{b}$$

where $\mathbf{A} \in \text{O}(n)$ and $\lambda \in \mathbb{R}^+$.

Yet another feature of Euclidean geometry is *volume*:

EXERCISE 1.1.5. *Show that an orientation-preserving isometry of Euclidean space is volume-preserving. Show that an orientation-preserving similarity transformation preserves volume if and only if it is an isometry.*

Compare §1.4.2 for further discussion of volume in affine geometry.

1.2. Affine space

What geometric properties of \mathbf{E}^n do not involve the metric notions of distance, angle and volume? For example, the notion of *straight line* is invariant under translations and more general linear maps which are not Euclidean isometries. Although it admits a metric definition as “the shortest path joining two points,” it enjoys a more fundamental characterization as a curve of zero acceleration, that is, a *geodesic*. However to define the acceleration of a smooth curve, one needs to compare the velocity vectors at *different* points along the curve. This is achieved by the *parallel transport* of the velocity along the curve, and hence involves the notion of *parallelism*. (This is the notion of an *affine*

connection, which is way to “connect” the infinitesimal displacements at different locations.)

Here is our first definition of an *affine transformation*:

DEFINITION 1.2.1. *An affine transformation of \mathbb{R}^n is a mapping of the form*

$$\begin{aligned} \mathbb{R}^n &\xrightarrow{g} \mathbb{R}^n \\ p &\longmapsto \mathbf{A}p + \mathbf{b} \end{aligned}$$

where $\mathbf{A} \in \mathrm{GL}(n, \mathbb{R})$ is an $n \times n$ invertible matrix and $\mathbf{b} \in \mathbb{R}^n$ is a vector. \mathbf{A} is called the linear part of g , and denoted $\mathbf{L}(g)$ and \mathbf{b} is called the translational part of g , and denoted $\mathbf{u}(g)$.

Thus an affine transformation g is:

- a translation if and only if $\mathbf{L}(g) = \mathbb{I}$;
- a Euclidean isometry if and only if $\mathbf{L}(g) \in \mathrm{O}(n)$;
- a Euclidean similarity (conformal transformation) if and only if $\mathbf{L}(g) \in \mathrm{Sim}_0(\mathbb{R}^n) = \mathbb{R}^+ \times \mathrm{O}(n)$;
- a volume-preserving affine transformation if and only if $\mathbf{L}(g) \in \mathrm{SL}(n, \mathbb{R})$.

1.2.1. The geometry of parallelism. Here is a more formal definition of an affine space. Although less intuitive, it embodies the idea that affine geometry is the geometry of *parallelism*.

Recall that subsets $X, Y \subset \mathbb{E}^n$ are *parallel* (written $X \parallel Y$) if and only if $\tau_{\mathbf{v}}(X) = Y$ for some vector $\mathbf{v} \in \mathbb{R}^n$. (Here $\tau_{\mathbf{v}} \in \mathrm{Trans}(\mathbb{E}^n)$ denotes the translation $p \longmapsto p + \mathbf{v}$.) Affine geometry is the geometry arising from the simply transitive action of the vector space of translations (isomorphic to \mathbb{R}^n).

Recall that an action of a group G on a space X is *simply transitive* if and only if for some (and then necessarily every) $x \in X$, the evaluation map

$$\begin{aligned} G &\longrightarrow X \\ g &\longmapsto g \cdot x \end{aligned}$$

is bijective: that is, for all $x, y \in X$, a unique $g \in G$ takes x to y . Equivalently, the action is both:

- *Transitive*: There is only one orbit, and
- *Free*: No nontrivial element fixes a point.

For further general discussion about group actions, see §A.3.

DEFINITION 1.2.2. *Let G be a group. A G -torsor is a space X with a simply transitive G -action.*

Thus a G -torsor is like the group G , except that the special role of its identity element is “forgotten.” Thus all the points are regarded as equivalent. What is remembered is the algebraic structure of the transformations in G which transport uniquely between the points.

Now we give the formal definition of an affine space:

DEFINITION 1.2.3. *An affine space is a V -torsor A , where V is a vector space. We call V the vector space underlying A , and denote it by $\text{Trans}(A)$, the elements of which are the translations of A .*

This abstract approach provides the usual coordinates for an affine space. Namely, choose a basepoint $p_0 \in A$ which will correspond to the *origin*. That is, it will be labeled by the *zero vector* $\mathbf{0} \in V$. Any other point $p \in A$ is related to p_0 by a unique translation $\tau \in \text{Trans}(A)$ satisfying $p = \tau(p_0)$. (This translation exists because the action is transitive, and is unique because the action is free.) Identifying the transformations $\text{Trans}(A)$ with vectors $\mathbf{v} \in V$ in the usual coordinates, the vector \mathbf{v} corresponding to τ is just $\mathbf{v} = p - p_0$, and τ is the mapping

$$\begin{aligned} A &\xrightarrow{\tau} A \\ p &\longmapsto p + \mathbf{v}. \end{aligned}$$

1.2.2. Affine transformations. Here is the second definition of affine transformations.

Affine maps are maps between affine spaces which are *compatible* with these simply transitive actions of vector spaces. Suppose A, A' are affine spaces with underlying vector spaces

$$V \longleftrightarrow \text{Trans}(A), \quad V' \longleftrightarrow \text{Trans}(A').$$

Then a map

$$A \xrightarrow{f} A'$$

is *affine* if for each $\tau \in \text{Trans}(A)$, there exists a translation $\tau' \in \text{Trans}(A')$ such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & A' \\ \tau \downarrow & & \downarrow \tau' \\ A & \xrightarrow{f} & A' \end{array}$$

commutes. Necessarily τ' is unique and evidently the correspondence

$$\tau \longmapsto \tau'$$

defines a homomorphism of groups $V \longrightarrow V'$, that is, a linear map between vector spaces. This linear map is the *linear part* of f , denoted

$L(f)$. Denoting the space of all affine maps $A \rightarrow A'$ by $\text{aff}(A, A')$ and the space of all linear maps $V \rightarrow V'$ by $\text{Hom}(V, V')$, linear part defines a map

$$\text{aff}(A, A') \xrightarrow{L} \text{Hom}(V, V')$$

The set of affine *endomorphisms* of an affine space A will be denoted by $\text{aff}(A)$ and the group of affine *automorphisms* of A will be denoted $\text{Aff}(A)$.

The notion of *translational part* involves choosing basepoints $p_0 \in A$ and $p'_0 \in A'$, respectively. Then the *translational part* of $f \in \text{aff}(A, A')$ is simply

$$\mathbf{u}(f) := f(p_0) - p'_0,$$

that is, the vector in V' corresponding to the translation $\tau_f \in \text{Trans}(A)$ taking $p'_0 \in A'$ to $f(p_0) \in A'$. Then $(\tau_f)^{-1} \circ f$ maps p_0 to p'_0 and corresponds to the linear part $L(f)$ as follows. Let $p \in A$ be an arbitrary point which corresponds to the vector

$$\mathbf{x} = p - p_0 \in V,$$

that is, the translation $\tau \in \text{Trans}(A)$ corresponding to \mathbf{x} maps p_0 to p , as in §1.2. Then

$$((\tau_f)^{-1} \circ f) \circ \tau(p_0) = \tau' \circ ((\tau_f)^{-1} \circ f)p_0 = \tau'(p'_0) = p'_0 + L(f)\mathbf{x}$$

where $\tau' \in \text{Trans}(A')$ corresponds to $L(f)\mathbf{x}$. Thus f is the composition of the translation $\tau_f \longleftrightarrow \mathbf{u}(f)$ with the linear part $L(f)$.

The space $\text{aff}(A, A')$ has the natural structure of an affine space. Namely the vector space

$$\text{Hom}(V, V') \oplus V'$$

acts simply transitively on $\text{aff}(A, A')$. Furthermore the Cartesian product $\text{Aff}(A) \times \text{Aff}(A')$ acts by composition on $\text{aff}(A, A')$, preserving the affine structure. (Compare Vey [275] for a discussion of the affine structure on the space of affine mappings.)

$\text{Aff}(A)$ is a Lie group and its Lie algebra identifies with $\text{aff}(A)$, which we later identify with *affine vector fields* on A . Furthermore $\text{Aff}(A)$ is isomorphic to the semidirect product $\text{Aut}(V) \ltimes V$, where V is the normal subgroup consisting of translations and

$$\text{Aut}(V) = \text{GL}(V)$$

is the group of linear automorphisms of the vector space $V \longleftrightarrow \text{Trans}(A)$.

Affine geometry is the study of affine spaces and affine maps between them. If $U \subset A$ is an open subset, then a map $U \xrightarrow{f} A'$ is *locally affine* if for each connected component U_i of U , there exists an affine map $f_i \in \text{aff}(A, A')$ such that the restrictions of f and f_i to U_i

are identical. Note that two affine maps which agree on a nonempty open set are identical.

1.3. The connection on affine space

Now we discuss the structure of an affine space \mathbf{A} as a smooth manifold. To analyze the differentiable structure on \mathbf{A} , we consider smooth paths in \mathbf{A} and their velocity vector fields, which live in the tangent bundle \mathbf{TA} . From this we “connect” the tangent spaces to define covariant differentiation enabling us to define acceleration as the covariant derivative of the velocity. Geodesics are curves of zero acceleration.

1.3.1. The tangent bundle of an affine space. Let $\gamma(t)$ denote a *smooth curve* in \mathbf{A} ; that is, in coordinates

$$\gamma(t) = (x^1(t), \dots, x^n(t))$$

where $x^j(t)$ are smooth functions of the time parameter, which ranges in an interval $[t_0, t_1] \subset \mathbb{R}$. The vector $\gamma(t) - \gamma(t_0)$ corresponds to the unique translation taking $\gamma(t_0)$ to $\gamma(t)$, and lies in the vector space \mathbf{V} underlying \mathbf{A} . It represents the displacement of the curve γ as it goes from $t = t_0$ to t . Define its *velocity vector* $\gamma'(t) \in \mathbf{V}$ as the derivative of this path in the vector space \mathbf{V} of translations. It represents the *infinitesimal* displacement of $\gamma(t)$ as t varies.

The set of tangent vectors is a vector space, denoted $\mathbf{T}_p M$, and naturally identifies with \mathbf{V} as follows. If $\mathbf{v} \in \mathbf{V}$ is a vector, then the path $\gamma_{(p, \mathbf{v})}(t)$ defined by:

$$(4) \quad t \mapsto p + t\mathbf{v} = \tau_{t\mathbf{v}}(p)$$

is a smooth path with $\gamma(0) = p$ and velocity vector $\gamma'(0) = \mathbf{v}$. Conversely, definition above of an infinitesimal displacement, shows that every smooth path through $p = \gamma(0)$ with velocity $\gamma'(0) = \mathbf{v}$ is tangent to the curve (4) as above.

The tangent spaces $\mathbf{T}_p M$ *linearize* M as follows. A mapping

$$M \xrightarrow{f} M'$$

is *differentiable at p* if every infinitesimal displacement $\mathbf{v} \in \mathbf{T}_p M$ maps to an infinitesimal displacement $\mathbf{D}_p f(\mathbf{v}) \in \mathbf{T}_q M'$, where $q = f(p)$. That is, if γ is a smooth curve with $\gamma(0) = p$ and $\gamma'(0) = \mathbf{v}$, then we require that $f \circ \gamma$ is a smooth curve through q at $t = 0$; then we call the new

velocity $(f \circ \gamma)'(0)$ the value of the *derivative*

$$\begin{aligned} T_p M &\xrightarrow{D_p f} T_q M' \\ \mathbf{v} &\longmapsto (f \circ \gamma)'(0) \end{aligned}$$

1.3.2. Parallel transport. On an affine space A , all the tangent spaces identify with each other. Namely, if $x, y \in A$, let $\tau \in \text{Trans}(A)$ be the unique translation taking x to y . (τ corresponds to the vector $y - x$.) The differential $(D\tau)_x$ maps $T_x A$ isomorphically to $T_y A$ and we denote this by:

$$T_x A \xrightarrow{\mathbb{P}_{x,y}} T_y A$$

We call this map *parallel transport* from x to y .

EXERCISE 1.3.1. *Another construction involves the linear structure of $\mathbf{V} \longleftrightarrow \text{Trans}(A)$. Namely, the action of \mathbf{V} by translations identifies the vector space \mathbf{V} with $T_x A$. Denoting this isomorphism by $\mathbf{V} \xrightarrow{\alpha_x} T_x A$, show that $\mathbb{P}_{x,y} = \alpha_y \circ (\alpha_x)^{-1}$.*

A vector field $\xi \in \text{Vec}(A)$ is *parallel* if it is invariant under parallel transport. That is, $\mathbb{P}_{x,y}(\xi_x) = \xi_y$ for any $x, y \in A$. This just means that ξ is a “constant vector field,” defined by a constant map $A \xrightarrow{\mathbf{v}} \mathbf{V}$: as a differential operator

$$\begin{aligned} C^\infty(A) &\xrightarrow{\xi} \mathbb{R} \\ f &\longmapsto v^i(x) \frac{\partial f}{\partial x^i} \end{aligned}$$

where $\mathbf{v}(x)$ is constant. Thus \mathbf{V} identifies with the space of parallel vector fields on A , and is based by the coordinate vector fields

$$\frac{\partial}{\partial x^i} \in \text{Vec}(A),$$

which we abbreviate simply by ∂_i .

EXERCISE 1.3.2. *Show that $\xi \in \text{Vec}(A)$ is parallel if and only if it generates a one-parameter group of translations.*

Similarly, the dual vector space \mathbf{V}^* identifies with parallel 1-forms as follows. A 1-form (covector field) on A corresponds to a constant map $A \rightarrow \mathbf{V}^*$. The basis of parallel covector fields dual to the coordinate basis $\{\partial_1, \dots, \partial_n\}$ of parallel vector fields is denoted $\{dx^1, \dots, dx^n\}$ (as usual).

EXERCISE 1.3.3. *Show that a parallel 1-form is exact, and hence closed.*

1.3.3. Acceleration and geodesics. The velocity vector field $\gamma'(t)$ of a smooth curve $\gamma(t)$ is an example of a *vector field along the curve* $\gamma(t)$: For each t , the tangent vector $\gamma'(t) \in T_{\gamma(t)}\mathbf{A}$. Differentiating the velocity vector field raises a significant difficulty: since the values of the vector field live in different vector spaces, we need a way to compare, or to *connect* them. The natural way is use the simply transitive action of the group \mathbf{V} of translations of \mathbf{A} . That is, suppose that $\gamma(t)$ is a smooth path, and $\xi(t)$ is a vector field along $\gamma(t)$. Let τ_s^t denote the translation taking $\gamma(t+s)$ to $\gamma(t)$, that is, in coordinates:

$$\begin{aligned} \mathbf{A} &\xrightarrow{\tau_s^t} \mathbf{A} \\ p &\longmapsto p + (\gamma(t+s) - \gamma(t)) \end{aligned}$$

Its differential

$$T_{\gamma(t+s)}\mathbf{A} \xrightarrow{D\tau_s^t} T_{\gamma(t)}\mathbf{A}$$

then maps $\xi(t+s)$ into $T_{\gamma(t)}\mathbf{A}$ and the *covariant derivative* $\frac{D}{dt}\xi(t)$ is the derivative of this smooth path in the *fixed* vector space $T_{\gamma(t)}\mathbf{A}$:

$$\begin{aligned} \frac{D}{dt}\xi(t) &:= \left. \frac{d}{ds} \right|_{s=0} (D\tau_s^t)(\xi(t+s)) \\ &= \lim_{s \rightarrow 0} \frac{(D\tau_s^t)(\xi(t+s)) - \xi(t)}{s} \end{aligned}$$

In this way, define the *acceleration* as the covariant derivative of the velocity:

$$\gamma''(t) := \frac{D}{dt}\gamma'(t)$$

A curve with zero acceleration is called a *geodesic*.

EXERCISE 1.3.4. *Given a point p and a tangent vector $\mathbf{v} \in T_p\mathbf{A}$, show that the unique geodesic $\gamma(t)$ with*

$$(\gamma(0), \gamma'(0)) = (p, \mathbf{v})$$

is given by (4).

In other words, geodesics in \mathbf{A} are parametrized curves which are Euclidean straight lines traveling at *constant* speed. However, in affine geometry the *speed* itself is not defined, but “motion along a straight line at constant speed” is affinely invariant (zero acceleration).

This leads to the following important definition:

DEFINITION 1.3.5. Let $p \in \mathbf{A}$ and $\mathbf{v} \in \mathbf{T}_p(\mathbf{A}) \cong \mathbf{V}$. Then the exponential mapping is defined by:

$$\begin{aligned} \mathbf{T}_p\mathbf{A} &\xrightarrow{\text{Exp}_p} \mathbf{A} \\ \mathbf{v} &\longmapsto p + \mathbf{v}. \end{aligned}$$

Thus the unique geodesic with initial position and velocity (p, \mathbf{v}) equals

$$t \longmapsto \text{Exp}_p(t\mathbf{v}) = p + t\mathbf{v}.$$

1.4. Parallel structures

Many important refinements of affine geometry involve structures which are *parallel*. Parallelism generalizes the notion of “constant” when the targets vary from point to point.

For example, the most familiar geometry is *Euclidean geometry*, extremely rich with metric notions such as distance, angle, area and volume. We have seen that *affine geometry* underlies it with the more primitive notion of *parallelism*. Euclidean geometry arises from affine geometry by introducing a Riemannian structure on \mathbf{A} , which is *parallel*.

Parallel vector fields and 1-forms were introduced back in §1.3.2, where parallel vector fields correspond to vectors in \mathbf{V} and parallel 1-forms (parallel covector fields) correspond to covectors in \mathbf{V}^* . Now we consider parallel tensor fields of higher order.

1.4.1. Parallel Riemannian structures. Let \mathbf{B} be an inner product on \mathbf{V} and $\mathbf{O}(\mathbf{V}; \mathbf{B}) \subset \mathbf{GL}(\mathbf{A})$ the corresponding orthogonal group. Then \mathbf{B} defines a flat Riemannian metric on \mathbf{A} and the inverse image

$$\mathbf{L}^{-1}(\mathbf{O}(\mathbf{V}; \mathbf{B})) \cong \mathbf{O}(\mathbf{V}; \mathbf{B}) \cdot \mathbf{Trans}(\mathbf{A})$$

is the full group of isometries, that is, the *Euclidean group*. If \mathbf{B} is a non-degenerate indefinite form, then there is a corresponding flat pseudo-Riemannian metric on \mathbf{A} and the inverse image $\mathbf{L}^{-1}(\mathbf{O}(\mathbf{V}; \mathbf{B}))$ is the full group of isometries of this pseudo-Riemannian metric.

EXERCISE 1.4.1. Show that an affine automorphism g of Euclidean n -space \mathbb{R}^n is conformal (that is, preserves angles) if and only if its linear part is the composition of an orthogonal transformation and scalar multiplication.

Such a transformation will be called a *similarity transformation* and the group of similarity transformations will be denoted $\mathbf{Sim}(\mathbf{E}^n)$. The scalar multiple is called the *scale factor* $\lambda(g) \in \mathbb{R}^+$ and defines

a homomorphism $\text{Sim}(\mathbf{E}^n) \xrightarrow{\lambda} \mathbb{R}^+$. In general, if $g \in \text{Sim}(\mathbf{E}^n)$, then $\exists A \in \text{O}(n), \mathbf{b} \in \mathbb{R}^n$ such that

$$g(x) = \lambda(g)Ax + \mathbf{b}.$$

1.4.2. Parallel tensor fields. Namely, any tangent vector $\mathbf{v}_p \in \mathbb{T}_p\mathbf{A}$ extends uniquely to a vector field on \mathbf{A} invariant under the group of translations. As we saw in §1.4.1, Euclidean structures are defined by extending an inner product from a single tangent space to all of \mathbf{E} .

Dual to parallel vector fields are *parallel 1-forms*. Every tangent covector $\omega_p \in \mathbb{T}_p^*\mathbf{A}$ extends uniquely to a translation-invariant 1-form.

EXERCISE 1.4.2. *Prove that a parallel 1-form is closed. Express a parallel 1-form in local coordinates.*

If $n = \dim(\mathbf{A})$, then an exterior n -form ω must be $f(x) dx^1 \wedge \cdots \wedge dx^n$ in local coordinates, where $f \in C^\infty(\mathbf{A})$ is a smooth function. Then ω is parallel if and only if $f(x)$ is constant.

EXERCISE 1.4.3. *Prove that an affine transformation $g \in \text{Aff}(\mathbf{A})$ preserves a parallel volume form if and only if $\det \mathbf{L}(g) = 1$.*

1.4.3. Complex affine geometry. We have been working entirely over \mathbb{R} , but it is clear one may study affine geometry over any field. If $\mathbf{k} \supset \mathbb{R}$ is a field extension, then every \mathbf{k} -vector space is a vector space over \mathbb{R} and thus every \mathbf{k} -affine space is an \mathbb{R} -affine space. In this way we obtain more refined geometric structures on affine spaces by considering affine maps whose linear parts are linear over \mathbf{k} .

EXERCISE 1.4.4. *Show that 1-dimensional complex affine geometry is the same as (orientation-preserving) 2-dimensional similarity geometry.*

This structure is another case of a parallel structure on an affine space, as follows. Recall a complex vector space has an underlying structure as a real vector space \mathbf{V} . The difference is a notion of scalar multiplication by $\sqrt{-1}$, which is given by a linear map

$$\mathbf{V} \xrightarrow{J} \mathbf{V}$$

such that $J \circ J = -\mathbb{I}$. Such an automorphism is called a *complex structure* on \mathbf{V} , and “turns \mathbf{V} into” a *complex vector space*.

If M is a manifold, an endomorphism field (that is, a $(1, 1)$ -tensor field) J where, for each $p \in M$, the value J_p is a complex structure on the tangent space \mathbb{T}_pM is called an *almost complex structure*. Necessarily $\dim(M)$ is even.

Recall that a *complex manifold* is a manifold with an atlas of coordinate charts where coordinate changes are biholomorphic. (Such an atlas is called a *holomorphic atlas*. Every complex manifold admits an almost complex structure, but not every almost complex structure arises from a holomorphic atlas, except in dimension two.

EXERCISE 1.4.5. *Prove that a complex affine space is the same as a affine space with a parallel almost complex structure.*

1.5. Affine vector fields

A vector field X on A is said to be *affine* if it generates a one-parameter group of affine transformations. Affine vector fields include parallel vector fields and radiant vector fields. Parallel vector fields generate one-parameter groups of translations, and *radiant* vector fields generate one-parameter groups of homotheties. Covariant differentiation provides general criteria characterizing affine vector fields.

1.5.1. Translations and parallel vector fields.

EXERCISE 1.5.1. *A vector field X on A is parallel if, for every $p, q \in A$, the values $X_p \in T_pA$ and $X_q \in T_qA$ are parallel.*

Since translation τ by $\mathbf{v} = q - p$ is the unique translation taking p to q , this simply means that the differential $D\tau$ maps X_p to X_q .

EXERCISE 1.5.2. *Let $X \in \text{Vec}(A)$ be a vector field on an affine space A . The following conditions are equivalent:*

- X is parallel
- The coefficients of X (in affine coordinates) are constant.
- $\nabla_Y X = 0$ for all $Y \in \text{Vec}(A)$.
- The covariant differential $\nabla X = 0$.
- The linear part $L(X) = 0$.

The vector space V identifies with the space of parallel vector fields on A . The parallel vector fields form an abelian Lie algebra of vector fields on A .

1.5.2. Homotheties and radiant vector fields. Another important class of affine vector fields are the *radiant vector fields*, or *infinitesimal homotheties*:

DEFINITION 1.5.3. *An affine transformation $\phi \in \text{Aff}(A)$ is a homothety if it is conjugate by a translation to scalar multiplication $\mathbf{v} \mapsto \lambda \mathbf{v}$, for some scalar $\lambda \in \mathbb{R}^\times$. An affine vector field is radiant if it generates a one-parameter group of homotheties.*

Observe that a homothety fixes a unique point $p \in \mathbf{A}$, which we often take to be the origin. The only zero of the corresponding radiant vector field is p .

Radiant vector fields are also called *Euler vector fields*, due to their role in Euler's theorem on homogeneous functions: Recall that a function $\mathbb{R}^n \xrightarrow{f} \mathbb{R}$ is *homogeneous of degree m* if and only if

$$f(\lambda x) = \lambda^m f(x)$$

for all $\lambda \in \mathbb{R}^+$.

THEOREM 1.5.4 (Euler). *Suppose that \mathbf{R} is the radiant vector field vanishing at the origin $\mathbf{0}$. Then f is homogeneous of degree m if and only if the directional derivative $\mathbf{R}f = mf$.*

EXERCISE 1.5.5. *Prove Euler's theorem above.*

Radiant vector fields play an important role in our study. Many important examples of affine manifolds admit radiant vector fields, and radiant vector fields provide a link between affine structures in dimension n and projective structures in dimension $n - 1$.

EXERCISE 1.5.6. *Let $\mathbf{R} \in \text{Vec}(\mathbf{A})$ be a vector field. Then the following conditions are equivalent:*

- \mathbf{R} is radiant;
- $\nabla \mathbf{R} = \mathbb{I}_{\mathbf{A}}$ (where $\mathbb{I}_{\mathbf{A}} \in \mathcal{T}^1(\mathbf{A}; \mathbb{T}\mathbf{A})$ is the identity map $\mathbb{T}\mathbf{A} \rightarrow \mathbb{T}\mathbf{A}$, regarded as an endomorphism field on \mathbf{A});
- there exists $b^i \in \mathbb{R}$ for $i = 1, \dots, n$ such that

$$\mathbf{R} = \sum_{i=1}^n (x^i - b^i) \frac{\partial}{\partial x^i}.$$

Note that $b = (b^1, \dots, b^n)$ is the unique zero of \mathbf{R} and that \mathbf{R} generates the one-parameter group of homotheties fixing b . (Thus a radiant vector field is a special kind of affine vector field.) Furthermore \mathbf{R} generates the center of the isotropy group of $\text{Aff}(\mathbf{A})$ at b , which is conjugate (by translation by b) to $\text{GL}(\mathbf{A})$. Show that the radiant vector fields on \mathbf{A} form an affine space isomorphic to \mathbf{A} .

1.5.3. Affineness criteria. Affine vector fields can be characterized in terms of the covariant differential operation

$$\mathcal{T}^p(M; \mathbb{T}M) \xrightarrow{\nabla} \mathcal{T}^{p+1}(M; \mathbb{T}M)$$

where $\mathcal{T}^p(M; \mathbb{T}M)$ denotes the space of $\mathbb{T}M$ -valued covariant p -tensor fields on M , that is, the tensor fields of type $(1, p)$. Thus $\mathcal{T}^0(M; \mathbb{T}M) = \text{Vec}(M)$, the space of vector fields on M . The space $\mathcal{T}^1(M; \mathbb{T}M)$ is

comprises \mathbb{T}^*M -valued vector fields on M , which identify with $\mathbb{T}M$ -valued 1-forms, or equivalently *endomorphism fields* on M .

EXERCISE 1.5.7. *X is affine if and only if it satisfies any of the following equivalent conditions:*

- For all $Y, Z \in \text{Vec}(\mathbf{A})$,

$$\nabla_Y \nabla_Z X = \nabla_{(\nabla_Y Z)} X.$$

- $\nabla \nabla X = 0$.
- The coefficients of X are affine functions, that is,

$$X = \sum_{i,j=1}^n (a^i_j x^j + b^i) \frac{\partial}{\partial x^i}$$

for constants $a^i_j, b^i \in \mathbb{R}$.

Write

$$\mathbf{L}(X) = \sum_{i,j=1}^n a^i_j x^j \frac{\partial}{\partial x^i}$$

for the linear part (which corresponds to the matrix $(a^i_j) \in \mathfrak{gl}(\mathbb{R}^n)$) and

$$X(0) = \sum_{i=1}^n b^i \frac{\partial}{\partial x^i}$$

for the translational part (the translational part of an affine vector field is a parallel vector field).

EXERCISE 1.5.8. *Under this correspondence, covariant derivative corresponds to composition of affine maps (matrix multiplication):*

$$\nabla_B A \quad \longleftrightarrow \quad \hat{\mathbf{A}} \hat{\mathbf{B}}$$

The Lie bracket of two affine vector fields is given by:

- $\mathbf{L}([X, Y]) = [\mathbf{L}(X), \mathbf{L}(Y)] = \mathbf{L}(X)\mathbf{L}(Y) - \mathbf{L}(Y)\mathbf{L}(X)$
(matrix multiplication)
- $[X, Y](0) = \mathbf{L}(X)Y(0) - \mathbf{L}(Y)X(0)$.

In this way the space $\text{aff}(\mathbf{A}) = \text{aff}(\mathbf{A}, \mathbf{A})$ of affine endomorphisms $\mathbf{A} \circlearrowleft$ is a Lie algebra.

Let M be an affine manifold. A vector field $\xi \in \text{Vec}(M)$ is *affine* if in local coordinates ξ appears as a vector field in $\text{aff}(\mathbf{A})$. We denote the space of affine vector fields on an affine manifold M by $\text{aff}(M)$.

EXERCISE 1.5.9. *Let M be an affine manifold.*

- (1) *Show that $\text{aff}(M)$ is a subalgebra of the Lie algebra $\text{Vec}(M)$.*

- (2) Show that the identity component of the affine automorphism group $\mathbf{Aut}(M)$ has Lie algebra $\mathfrak{aff}(M)$.
- (3) If ∇ is the flat affine connection corresponding to the affine structure on M , show that a vector field $\xi \in \mathbf{Vec}(M)$ is affine if and only if

$$\nabla_{\xi}v = [\xi, v]$$

$$\forall v \in \mathbf{Vec}(M).$$

1.6. Affine subspaces

Suppose that $A_1 \xrightarrow{\iota} A$ is an injective affine map; then we say that $\iota(A_1)$ (or with slight abuse, ι itself) is an *affine subspace*. If A_1 is an affine subspace then for each $x \in A_1$ there exists a linear subspace $V_1 \subset \mathbf{Trans}(A)$ such that A_1 is the orbit of x under V_1 (that is, “an affine subspace in a vector space is just a coset (or translate) of a linear subspace $A_1 = x + V_1$.”) An affine subspace of dimension 0 is thus a point and an affine subspace of dimension 1 is a line.

EXERCISE 1.6.1. Show that if l, l' are (affine) lines and

$$(x, y) \in l \times l, \quad x \neq y$$

$$(x', y') \in l' \times l', \quad x' \neq y'$$

are pairs of distinct points. Then there is a unique affine map $l \xrightarrow{f} l'$ such that

$$f(x) = x',$$

$$f(y) = y'.$$

If $x, y, z \in l$ (with $x \neq y$), then define $[x, y, z]$ to be the image of z under the unique affine map $l \xrightarrow{f} \mathbb{R}$ with $f(x) = 0$ and $f(y) = 1$. Show that if $l = \mathbb{R}$, then $[x, y, z]$ is given by the formula

$$[x, y, z] = \frac{z - x}{y - x}.$$

This is called an *affine parameter* along the line.

1.7. Volume in affine geometry

Although an affine automorphism of an affine space A need not preserve a natural measure on A , Euclidean volume nonetheless does behave rather well with respect to affine maps. The Euclidean volume form ω can almost be characterized affinely by its parallelism: it is invariant under all translations. Moreover two $\mathbf{Trans}(A)$ -invariant volume forms differ by a scalar multiple but there is no natural way to

normalize. Such a volume form will be called a *parallel volume form*. If $g \in \text{Aff}(\mathbf{A})$, then the distortion of volume is given by

$$g^*\omega = \det \mathbf{L}(g) \cdot \omega.$$

Thus although there is no canonically normalized volume or measure there is a natural affinely invariant line of measures on an affine space. The subgroup $\text{SAff}(\mathbf{A})$ of $\text{Aff}(\mathbf{A})$ consisting of volume-preserving affine transformations is the inverse image $\mathbf{L}^{-1}(\text{SL}(\mathbf{V}))$, sometimes called the *special affine group* of \mathbf{A} . Here $\text{SL}(\mathbf{V})$ denotes, as usual, the *special linear group*

$$\text{Ker}(\text{GL}(\mathbf{V}) \xrightarrow{\det} \mathbb{R}^\times) = \{g \in \text{GL}(\mathbf{V}) \mid \det(g) = 1\}.$$

1.7.1. Centers of gravity. Given a finite subset $F \subset \mathbf{A}$ of an affine space, its *center of gravity* or *centroid* $\bar{F} \in \mathbf{A}$ is point associated with F in an affinely invariant way: that is, given an affine map $\mathbf{A} \xrightarrow{\phi} \mathbf{A}'$ we have

$$\overline{\phi(F)} = \phi(\bar{F}).$$

This operation can be generalized as follows.

THEOREM 1.7.1. *Let μ be a probability measure on an affine space \mathbf{A} . Then there exists a unique point $\bar{x} \in \mathbf{A}$ (the centroid of μ) such that for all affine maps $\mathbf{A} \xrightarrow{f} \mathbb{R}$,*

$$(5) \quad f(x) = \int_{\mathbf{A}} f \, d\mu$$

PROOF. Let (x^1, \dots, x^n) be an affine coordinate system on \mathbf{A} . Let $\bar{x} \in \mathbf{A}$ be the points with coordinates $(\bar{x}^1, \dots, \bar{x}^n)$ given by

$$\bar{x}^i = \int_{\mathbf{A}} x^i \, d\mu.$$

This uniquely determines $\bar{x} \in \mathbf{A}$; we must show that (5) is satisfied for all affine functions. Suppose $\mathbf{A} \xrightarrow{f} \mathbb{R}$ is an affine function. Then there exist a_1, \dots, a_n, b such that

$$f = a_1x^1 + \dots + a_nx^n + b$$

and thus

$$\begin{aligned} f(\bar{x}) &= a_1 \int_{\mathbf{A}} x^1 \, d\mu + \dots + a_n \int_{\mathbf{A}} x^n \, d\mu \\ &\quad + b \int_{\mathbf{A}} d\mu = \int_{\mathbf{A}} f \, d\mu \end{aligned}$$

as claimed. □

Now let $C \subset A$ be a *convex body*, that is, a convex open subset having compact closure. Then C determines a probability measure μ_C on A by

$$\mu_C(X) = \frac{\int_{X \cap C} \omega}{\int_C \omega}$$

where ω is any parallel volume form on A .

PROPOSITION 1.7.2. *Let $C \subset A$ be a convex body. Then the centroid \bar{C} of C lies in C .*

PROOF. C is the intersection of halfspaces, that is, C consists of all $x \in A$ such that $f(x) > 0$ for all affine maps

$$A \xrightarrow{f} \mathbb{R}$$

such that $f|_C > 0$. If f is such an affine map, then clearly $f(\bar{C}) > 0$. Therefore $\bar{C} \in C$. \square

1.7.2. Divergence. If $\xi \in \text{Vec}(M)$, then the infinitesimal distortion of volume is the *divergence* of ξ , defined as the function $\text{div}(\xi)$ such that

$$\mathcal{L}_\xi(\omega) = \text{div}(\xi)\omega$$

where ω is (any) parallel volume form and \mathcal{L}_ξ denotes Lie differentiation with respect to ξ . If, in coordinates $\xi = \xi^i \partial_i$, then

$$\text{div}(\xi) = \partial_i \xi^i$$

(the usual formula).

EXERCISE 1.7.3. *The Lie algebra of the special affine group $\text{SAff}(A)$ consists of affine vector fields of divergence zero.*

1.8. Linearizing affine geometry

Associated to every affine space A is an embedding \mathcal{A}' of A as an *affine hyperplane* in a vector space W as follows.

EXERCISE 1.8.1. *Let A be an affine space over a field k with underlying vector space $V := \text{Trans}(A)$. Let $W := V \oplus k$ and let $W \xrightarrow{\psi} k$ denote linear projection onto the second summand.*

- For each $s \in k$, the group V acts simply transitively on the affine hyperplane $\psi^{-1}(s)$.
 - A identifies with $\psi^{-1}(1)$.
 - V identifies with $\text{Ker}(\psi) = \psi^{-1}(0)$.

- Define a bijective correspondence between n -dimensional affine spaces \mathbf{A} and pairs (\mathbf{W}, ψ) where \mathbf{W} is an $n + 1$ -dimensional vector space and $\psi \in \mathbf{W}^*$ is a nonzero covector, where \mathbf{A} corresponds to $\psi^{-1}(1)$.
- Identify the affine group $\text{Aff}(\mathbf{A})$ with the subgroup of $\text{GL}(\mathbf{W})$ preserving this hyperplane, as well as the stabilizer of ψ .
- If $[\mathbf{A} \mid \mathbf{a}]$ represents an affine transformation with linear part \mathbf{A} and translational part \mathbf{a} , show that the corresponding linear transformation of \mathbf{W} is represented by the block matrix

$$\begin{bmatrix} \mathbf{A} & \mathbf{a} \\ \mathbf{0} & 1 \end{bmatrix}$$

where $\mathbf{0}$ is the row vector representing the zero map $\mathbf{V} \rightarrow \mathbb{R}$.

In coordinates, $\mathcal{A}'(\mathbf{v}) = \begin{bmatrix} v^1 \\ \vdots \\ v^n \\ 1 \end{bmatrix} \in \mathbf{W}$ and $\psi = [0 \dots 0 \ 1] \in \mathbf{W}^*$. The

affine transformation has linear part $\mathbf{A} = \begin{bmatrix} A^1_1 & \dots & A^1_n \\ \vdots & & \vdots \\ A^n_1 & \dots & A^n_n \end{bmatrix} \in \text{GL}(\mathbf{V})$ and

translational part $\mathbf{a} = \begin{bmatrix} a^1 \\ \vdots \\ a^n \end{bmatrix} \in \mathbf{V}$:

$$\begin{aligned} \mathbf{v} \xrightarrow{[\mathbf{A} \mid \mathbf{a}]} & \begin{bmatrix} A^1_1 & \dots & A^1_n \\ \vdots & A^i_j & \vdots \\ A^n_1 & \dots & A^n_n \end{bmatrix} \begin{bmatrix} v^1 \\ \vdots \\ v^j \\ \vdots \\ v^n \end{bmatrix} + \begin{bmatrix} a^1 \\ \vdots \\ a^i \\ \vdots \\ a^n \end{bmatrix} = \begin{bmatrix} A^1_j v^j + a^1 \\ \vdots \\ A^i_j v^j + a^i \\ \vdots \\ A^n_j v^j + a^n \end{bmatrix} \\ \mathcal{A}'(\mathbf{v}) \mapsto & \left[\begin{array}{ccc|c} A^1_1 & \dots & A^1_n & a^1 \\ \vdots & & \vdots & \vdots \\ A^n_1 & \dots & A^n_n & a^n \\ \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 \end{array} \right] \begin{bmatrix} v^1 \\ \vdots \\ v^n \\ 1 \end{bmatrix} = \begin{bmatrix} A^1_i v^i + a^1 \\ \vdots \\ A^n_i v^i + a^n \\ \dots \\ 1 \end{bmatrix} \end{aligned}$$

CHAPTER 2

Projective geometry

Projective geometry arose historically out of the efforts of Renaissance artists to understand perspective. Imagine a painter looking at a 2-dimensional canvas with one eye closed. His open eye plays the role of the the origin in the 3-dimensional vector space \mathbb{W} and the canvas plays the role of an affine hyperplane $A \subset \mathbb{W}$ as in §1.8. As the canvas tilts, the geometry seen by the painter changes. Parallel lines no longer appear parallel (like the railroad tracks described above) and distance and angle are distorted. But lines stay lines and the basic relations of collinearity and concurrence are unchanged. The change in perspective given by “tilting” the canvas is determined by a linear transformation of \mathbb{W} , since a point on A is determined completely by the 1-dimensional linear subspace of \mathbb{V} containing it. (One must solve systems of linear equations to write down the effect of such transformation.) Projective geometry is the study of points, lines and the incidence relations between them.

Projective space *closes off* affine space — that is, it *compactifies* affine space by adding *ideal points* at infinity. To develop an intuitive feel for projective geometry, consider how points in A^n may go to infinity:

The easiest way to go to infinity in A^n is by following geodesics, since they have zero acceleration. Furthermore two geodesics approach the same ideal point if they are parallel. A good model are railroad tracks running parallel to each other — they meet ideally at the horizon. We thus force parallel lines to intersect by attaching ideal points where the extended parallel lines are to intersect.

A good general reference for projective geometry (especially in dimension two) is Coxeter [81]), as well as Berger [37, 38] and Coxeter [80]. More classical treatments are Busemann-Kelly [54], Semple-Kneebone [248] and Veblen-Young [271, 272].

2.1. Ideal points

Parallelism of lines in A is an equivalence relation. Define an *ideal point* of A as an equivalence class. The *ideal set* of an affine space A is

the space $\mathbb{P}_\infty(\mathbf{A})$ of ideal points, with the quotient topology. If $l, l' \subset \mathbf{A}$ are parallel lines, then the point in \mathbb{P}_∞ corresponding to their parallelism class is defined as their intersection. So two lines are parallel if and only if they intersect at infinity.

Projective space is defined as the union $\mathbb{P} := \mathbf{A} \cup \mathbb{P}_\infty(\mathbf{A})$, with a suitable topology. The natural structure on \mathbb{P} is perhaps most easily seen in terms of an alternate, maybe more familiar, description. Embed \mathbf{A} as an affine hyperplane in a vector space $\mathbf{W} \cong \mathbf{V} \oplus \mathbf{k}$ as in §1.8, where $\mathbf{V} = \text{Trans}(\mathbf{A})$ is the vector space underlying \mathbf{A} . For example, if

$$p = \begin{bmatrix} p^1 \\ \vdots \\ p^n \end{bmatrix} \in \mathbf{A}^n,$$

then its embedding in $\mathbf{W} = \mathbf{k}^{n+1}$ is the nonzero vector

$$\mathcal{A}'(p) = \begin{bmatrix} p^1 \\ \vdots \\ p^n \\ 1 \end{bmatrix} \in \mathbf{W}.$$

Furthermore the line $\mathbf{k}\mathcal{A}'(p)$ it spans meets the affine hyperplane $\mathbf{A} \leftrightarrow \mathbf{k}^n \times \{1\}$ in a single point. Think of \mathbf{A} as the “canvas” or *viewing hyperplane*, and the line $\mathbf{k}\mathcal{A}'(p)$ as the line of sight as the point p is viewed from the origin $\mathbf{0} \in \mathbf{W}$ (the eye of the painter). The nonzero elements of this line $\mathbf{k}^\times \mathcal{A}'(p)$ form the *projective equivalence class* $[\mathcal{A}'(p)]$.

Now suppose that p travels to infinity along an affine geodesic $\ell \subset \mathbf{A}$:

$$p(t) := p + t\mathbf{v},$$

where $t \in \mathbf{k}$ and $\mathbf{v} \in \mathbf{V}$ is a nonzero vector. Denote the corresponding path of vectors in \mathbf{W} by

$$\mathbf{w}(t) := \mathcal{A}'(p(t)) \in \mathbf{W}.$$

Although $\lim_{t \rightarrow \infty} \mathbf{w}(t)$ does not exist, the corresponding lines $\mathbf{k} \cdot \mathbf{w}(t)$ converge to the line $\mathbf{k} \cdot \mathbf{v}$ corresponding to $\mathbf{v} \in \mathbf{W}$. This limiting line defines the *ideal point* of the affine line ℓ :

$$\lim_{t \rightarrow \infty} [\mathbf{w}(t)] = [\mathbf{v}]$$

This motivates the following fundamental definition:

DEFINITION 2.1.1. *Let \mathbf{W} denote a vector space over \mathbf{k} . The projective space associated to \mathbf{W} is the space $\mathbb{P}(\mathbf{W})$ of projective equivalence classes $[\mathbf{w}]$ of nonzero vectors $\mathbf{w} \in \mathbf{W}$, with the quotient topology.*

Thus a point in $P(W)$ (a “projective point”) corresponds to a *line* (that is, a one-dimensional linear subspace) in W . If $\mathbf{w} = \begin{bmatrix} w^1 \\ \vdots \\ w^{n+1} \end{bmatrix} \in W$, the corresponding projective point is

$$p := [\mathbf{w}] = \left[\begin{array}{c} w^1 \\ \vdots \\ w^{n+1} \end{array} \right] \in P(W)$$

and w^1, \dots, w^{n+1} are the *homogeneous coordinates* of p .

Since linear transformations of W preserve lines, $GL(W) = \text{Aut}(W)$ acts on $P(W)$; the induced transformations are the *projective transformations* or *collineations* of $P(W)$.

EXERCISE 2.1.2. *The action of $GL(W)$ on $P(W)$ is not effective. Its kernel consists of the group k^\times of nonzero scalings, which forms the center of $GL(W)$. The projective group or collineation group is the quotient*

$$PGL(W) := GL(W)/k^\times$$

which does act effectively.

2.2. Projective subspaces

Returning to the projective geometry of the line ℓ , note that the affine line $\mathbf{w}(t)$ in W lies in the linear 2-plane $\text{span}(p, \mathbf{v}) \subset W$. The one-dimensional linear subspaces contained in this linear 2-plane is a *projective line*.

DEFINITION 2.2.1. *Let $P = P(W)$ be a projective space, and let d be a nonnegative integer. A d -dimensional projective subspace S of P is the collection of all projective equivalence classes $[\mathbf{v}]$ of nonzero vectors \mathbf{v} lying in a fixed $d + 1$ -dimensional linear subspace $S \subset W$. We write $S = P(S)$ and call S the projectivization of S .*

Thus a *projective line* is the projectivization of a linear 2-plane in W and a *projective hyperplane* is the projectivization a linear hyperplane in W .

A linear embedding $S_1 \hookrightarrow S_2 \subset W$ of linear subspaces induces an embedding of projective subspaces $P(S_1) \hookrightarrow P(S_2)$ and we say that the subspaces $P(S_1)$ and $P(S_2)$ are *incident*. Clearly projective transformations preserve the relation of incidence. The converse, that an incidence-preserving transformation of projective space, is a deep theorem, sometimes called the *fundamental theorem of projective geometry*.

EXERCISE 2.2.2. *Show that the set of ideal points is a projective hyperplane.*

Suppose that $S_1, S_2 \subset \mathbf{P}$ are disjoint projective subspaces. Then, writing $S_i = \mathbf{P}(S_i)$ for respective linear subspaces $S_i \subset \mathbf{W}$, the projectivization $\mathbf{P}(S_1 + S_2)$ is a projective subspace $\text{span}(S_1, S_2) \subset \mathbf{P}$ and

$$\dim(\text{span}(S_1, S_2)) = \dim(S_1) + \dim(S_2) + 1.$$

If S_1 and S_2 are points, then $\text{span}(S_1, S_2)$ is a line, and we use the more familiar notation $\overleftrightarrow{S_1 S_2}$. If S_i are projective subspaces and $S_1 \cap S_2 \neq \emptyset$, then $S_1 \cap S_2$ is a projective subspace and

$$\dim(\text{span}(S_1, S_2)) + \dim(S_1 \cap S_2) = \dim(S_1) + \dim(S_2).$$

Evidently $\text{span}(S_1, S_2)$ is the smallest projective subspace containing S_1 and S_2 .

2.2.1. Affine patches. Ideal points are only special when projective space is the completion of affine space; by changing the viewing hyperplane, one gets different notions of “ideal.” Indeed, every projective point has neighborhoods which are affine subspaces.

Let \mathbf{P} be d -dimensional projective space and $H \subset \mathbf{P}$ be a projective hyperplane. Then the complement $\mathbf{P} \setminus H$ is an *affine patch* and has the structure as a d -dimensional affine space with underlying vector space \mathbf{V} via an *affine chart*

$$\mathbf{V} \xrightarrow[\approx]{\mathcal{A}} \mathbf{P} \setminus H,$$

defined as follows. Write $\mathbf{P} = \mathbf{P}(\mathbf{W})$. Choose a covector $\psi \in \mathbf{W}^*$ such that $H = \mathbf{P}(\mathbf{V})$ where $\mathbf{V} := (\text{Ker}(\psi))$. Choose a vector $\mathbf{w}_0 \in \mathbf{W}$ with $\psi(\mathbf{w}_0) = 1$ to define an origin in the affine patch. Then

$$\begin{aligned} \mathbf{V} &\xrightarrow{\mathcal{A}(\psi, \mathbf{w}_0)} \mathbf{P} \setminus H \\ \mathbf{v} &\longmapsto [\mathbf{w}_0 + \mathbf{v}] \end{aligned}$$

defines an affine chart on $\mathbf{P} \setminus H$. Compare §1.8.

Every projective point $p \in \mathbf{P}$ lies in an affine patch. Writing

$$p = [\mathbf{X}] = \left[\begin{array}{c} X^1 \\ \vdots \\ X^{d+1} \end{array} \right] \in \mathbf{P}^d,$$

some homogeneous coordinate X^i of the nonzero vector \mathbf{X} is nonzero. Then $X^i \neq 0$ defines an affine patch with chart

$$\mathbf{k}^d \xrightarrow{\mathcal{A}^i} \mathbf{P}^d$$

$$\begin{bmatrix} v^1 \\ \vdots \\ v^d \end{bmatrix} \mapsto \begin{bmatrix} v^1 \\ \vdots \\ v^{i-1} \\ 1 \\ v^i \\ \vdots \\ v^d \end{bmatrix}$$

and $p \in \mathbf{A}^{(i)} := \mathcal{A}^{(i)}(\mathbf{V})$. These $d + 1$ coordinate affine patches define a covering by contractible open sets.

EXERCISE 2.2.3. Suppose that $1 \leq i \neq j \leq d + 1$.

- (1) Express the intersection $\mathbf{A}^{(i)} \cap \mathbf{A}^{(j)}$ in terms of the charts $\mathcal{A}^{(i)}, \mathcal{A}^{(j)}$.
- (2) Compute the change of coordinates

$$(\mathcal{A}^{(i)})^{-1}(\mathbf{A}^{(i)} \cap \mathbf{A}^{(j)}) \xrightarrow{(\mathcal{A}^{(j)})^{-1} \circ \mathcal{A}^{(i)}} (\mathcal{A}^{(j)})^{-1}(\mathbf{A}^{(i)} \cap \mathbf{A}^{(j)}).$$

- (3) Let $\mathbf{k} = \mathbb{R}$ and let $a, b \in \mathbb{R}$ with $a < b$. Suppose that

$$(a, b) \xrightarrow{\gamma} \mathbf{V}$$

is a curve such that $\mathcal{A}^{(i)}(\gamma(t)) \in \mathbf{A}^{(i)}$ and $\mathcal{A}^{(j)}(\gamma(t)) \in \mathbf{A}^{(j)}$ for $a < t < b$. Suppose that $\mathcal{A}^{(i)} \circ \gamma$ is a geodesic in $\mathbf{A}^{(i)}$. Show there exists a reparametrization, that is, a diffeomorphism

$$(a, b) \xrightarrow[\approx]{\tau} \tau((a, b)) \subset \mathbb{R}$$

such that the composition $\mathcal{A}^{(j)} \circ \gamma \circ \tau$, that is, the map

$$t \mapsto \mathcal{A}^{(j)}(\gamma(\tau(t))),$$

is a geodesic in $\mathbf{A}^{(j)}$.

In general, the topology of projective space is complicated. Since it arises from a *quotient* and not a *subset* construction, it is more sophisticated than a subset. Indeed, projective space generally does not arise as a *hypersurface* in Euclidean space. Although \mathbf{P}^d can be covered by $d + 1$ contractible open sets, it cannot be covered by fewer contractible open sets. For either $\mathbf{k} = \mathbb{R}$ or \mathbb{C} , projective space $\mathbf{P}^d(\mathbf{k})$ is a compact smooth manifold. We summarize some basic facts about the topology.

EXERCISE 2.2.4. Suppose $k = \mathbb{R}$. Exhibit $\mathbb{P}^d(\mathbb{R})$ as a quotient of the unit sphere $S^d \subset \mathbb{R}^{d+1}$ by the antipodal map. Show that $\mathbb{P}^1(\mathbb{R}) \approx S^1$ and for $d > 1$ the fundamental group $\mathbb{P}^d(\mathbb{R})$ has order two. Show that $\mathbb{P}^d(\mathbb{R})$ is orientable if and only if d is odd.

EXERCISE 2.2.5. Suppose that $k = \mathbb{C}$. Exhibit $\mathbb{P}^d(\mathbb{C})$ as a quotient of the unit sphere $S^{2d-1} \subset \mathbb{C}^{d+1}$ by the group \mathbb{T} of unit complex numbers. Show that $\mathbb{P}^1(\mathbb{C}) \approx S^2$, and $\mathbb{P}^d(\mathbb{C})$ is simply connected and orientable for all $d \geq 1$.

EXERCISE 2.2.6. Find a natural S^1 -fibration $\mathbb{P}^{2d+1}(\mathbb{R}) \rightarrow \mathbb{P}^d(\mathbb{C})$.

2.3. Projective mappings

Linear mappings $V \xrightarrow{\phi} W$ between vector spaces define mappings between the corresponding projective spaces. However, if ϕ is not injective, the corresponding projective map is not defined on all of $\mathbb{P}(V)$. We begin discussing with projective maps defined by injective linear maps, particularly emphasizing *automorphisms*, classically known as *collineations*. Collineations arise from linear automorphisms of the vector space W .

2.3.1. Embeddings and Collineations. A projective subspace $S \subset \mathbb{P}(W)$ determines a projective map, determined by the linear inclusion $S \hookrightarrow W$, where S is the linear subspace of W projectivizing to S .

EXERCISE 2.3.1. Show that an injective projective map $\mathbb{P}(S) \rightarrow \mathbb{P}(W)$ is determined by an injective linear map $S \rightarrow W$ is determined by an injective linear map, unique up to composition with a homothety of S on the left and a homothety of W on the right.

A linear automorphism of a vector space W induces an invertible transformation $\mathbb{P}(W) \xrightarrow{\phi} \mathbb{P}(W)$. We call such a transformation a *collineation* or a *homography*. Evidently a collineation preserves projective subspaces, and the relations between them. An *involution* is a collineation of order two, that is, $\phi = \phi^{-1}$.

EXERCISE 2.3.2. Let $\mathbb{R}P^n$ be a real projective space of dimension n , and let ϕ be an involution of $\mathbb{R}P^n$.

- Suppose n is even. Then $\text{Fix}(\phi)$ is the union of two disjoint projective subspaces of dimensions d_1, d_2 where $d_1 + d_2 = n - 1$.
- Suppose $n = 2m + 1$ is odd. Then either:
 - $\text{Fix}(\phi) \neq \emptyset$ and equals the union of two disjoint projective subspaces of dimensions d_1, d_2 where $d_1 + d_2 = n - 1$, or

– $\text{Fix}(\phi) = \emptyset$, and ϕ leaves invariant an S^1 -fibration

$$\mathbb{R}\mathbb{P}^n \longrightarrow \mathbb{C}\mathbb{P}^m.$$

For example, an involution ϕ of $\mathbb{R}\mathbb{P}^2$ has an isolated fixed point p and a disjoint fixed line l . In an affine patch containing p , ϕ looks like a symmetry in p , preserving the local orientation. In contrast, ϕ looks like a reflection in l in an affine patch containing l , reversing the local orientation. Since $\mathbb{R}\mathbb{P}^2$ is nonorientable no *global* orientation exists to either preserve or reverse.

Here is an explicit example. The involution defined in homogeneous coordinates by:

$$\begin{aligned} \mathbb{R}\mathbb{P}^2 &\xrightarrow{\iota} \mathbb{R}\mathbb{P}^2 \\ \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} &\longmapsto \begin{bmatrix} -X \\ -Y \\ Z \end{bmatrix} \end{aligned}$$

fixes the point

$$p = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

and the projective line defined by $Z = 0$, that is, $l = [0 \ 0 \ 1]$. In the affine chart \mathcal{A}^3 , the isolated fixed point has coordinates $(0, 0)$ and ι appears as the *symmetry*

$$\begin{bmatrix} v^1 \\ v^2 \end{bmatrix} \longmapsto \begin{bmatrix} -v^1 \\ -v^2 \end{bmatrix}.$$

and in the affine chart \mathcal{A}^1 , the fixed line has coordinates $(*, 0)$ and ι appears as the reflection

$$\begin{bmatrix} v^1 \\ v^2 \end{bmatrix} \longmapsto \begin{bmatrix} -v^1 \\ v^2 \end{bmatrix}$$

fixing the vertical axis. That a single reflection can appear simultaneously as a symmetry in a point and reflection in a line indicates the topological complexity of \mathbb{P}^2 : A reflection in a line *reverses* a local orientation about a point on the line, and a symmetry in a point *preserves* a local orientation about the point.

2.3.2. Singular projective mappings. When the linear map ϕ is not injective, then $\mathbb{P}(\text{Ker}(\phi))$ is a projective subspace, upon which the projectivization $[\phi]$ of ϕ is not defined. For that reason, we call it the *undefined set* of $[\phi]$, and denote it $\mathbb{U}([\phi])$.

EXERCISE 2.3.3. Show that the projective automorphisms of \mathbb{P} form a group $\text{Aut}(\mathbb{P})$ which has the following description. If

$$\mathbb{P} \xrightarrow{f} \mathbb{P}$$

is a projective automorphism, some linear isomorphism

$$\mathbb{V} \xrightarrow{\tilde{f}} \mathbb{V}$$

induces f . Indeed,

$$1 \longrightarrow \mathbb{R}^\times \longrightarrow \text{GL}(\mathbb{V}) \longrightarrow \text{Aut}(\mathbb{P}) \longrightarrow 1$$

is a short exact sequence, where $\mathbb{R}^\times \longrightarrow \text{GL}(\mathbb{V})$ is the inclusion of the group of multiplications by nonzero scalars. This quotient, the projective general linear group

$$\text{PGL}(\mathbb{V}) := \text{GL}(\mathbb{V}) / \mathbb{R}^\times \cong \text{Aut}(\mathbb{P}^n),$$

is also denoted $\text{PGL}(n+1, \mathbb{R})$. If n is even, then

$$\text{PGL}(n+1, \mathbb{R}) \cong \text{SL}(n+1; \mathbb{R}).$$

If n is odd, then $\text{PGL}(n+1, \mathbb{R})$ has two connected components, and its identity component is doubly covered by $\text{SL}(n+1; \mathbb{R})$ and is isomorphic to $\text{SL}(n+1; \mathbb{R}) / \{\pm \mathbb{I}\}$.

If \mathbb{V}, \mathbb{V}' are vector spaces with associated projective spaces \mathbb{P}, \mathbb{P}' then a linear map $\mathbb{V} \xrightarrow{\tilde{f}} \mathbb{V}'$ maps lines through 0 to lines through 0. But \tilde{f} only induces a map $\mathbb{P} \xrightarrow{f} \mathbb{P}'$ if it is injective, since $f(x)$ can only be defined if $\tilde{f}(\tilde{x}) \neq 0$ (where \tilde{x} is a point of $\Pi^{-1}(x) \subset \mathbb{V} - \{0\}$). Suppose that \tilde{f} is a (not necessarily injective) linear map and let

$$\mathbb{U}(f) = \Pi(\text{Ker}(\tilde{f})).$$

The resulting *projective endomorphism* of \mathbb{P} is defined on the complement $\mathbb{P} - \mathbb{U}(f)$. If $\mathbb{U}(f) \neq \emptyset$, the corresponding projective endomorphism is by definition a *singular projective transformation* of \mathbb{P} . If f is singular, its image is a proper projective subspace, called the *range* of f and denoted $\mathcal{R}(f)$.

A projective map $\mathbb{P}_1 \xrightarrow{\iota} \mathbb{P}$ corresponds to a linear map $\mathbb{V}_1 \xrightarrow{\tilde{\iota}} \mathbb{V}$ between the corresponding vector spaces (well-defined up to scalar multiplication). Since ι is defined on all of \mathbb{P}_1 , $\tilde{\iota}$ is an injective linear map and hence corresponds to an embedding. Such an embedding (or its image) will be called a *projective subspace*. Projective subspaces of dimension k correspond to linear subspaces of dimension $k+1$. (By convention the empty set is a projective space of dimension -1.) Note that the “bad set” $\mathbb{U}(f)$ of a singular projective transformation is a

projective subspace. Two projective subspaces of dimensions k, l where $k+l \geq n$ intersect in a projective subspace of dimension at least $k+l-n$. The rank of a projective endomorphism is defined to be the dimension of its image.

EXERCISE 2.3.4. Let \mathbf{P} be a projective space of dimension n . Show that the (possibly singular) projective transformations of \mathbf{P} form themselves a projective space of dimension $(n+1)^2 - 1$. We denote this projective space by $\text{End}(\mathbf{P})$. Show that if $f \in \text{End}(\mathbf{P})$, then

$$\dim N(f) + \text{rank}(f) = n - 1.$$

Show that $f \in \text{End}(\mathbf{P})$ is nonsingular (in other words, a collineation) if and only if $\text{rank}(f) = n$, that is, $\mathcal{U}(f) = \emptyset$. Equivalently, $\mathcal{R}(f) = \mathbf{P}$.

An important kind of projective endomorphism is a *projection*, also called a *perspectivity*. Let $A^k, B^l \subset \mathbf{P}^n$ be disjoint projective subspaces whose dimensions satisfy $k+l = n-1$. Define the projection $\Pi = \Pi_{A^k, B^l}$ onto A^k from B^l

$$\mathbf{P}^n - B^l \xrightarrow{\Pi} A^k$$

as follows. For every $x \in \mathbf{P}^n - A^k$ the minimal projective subspace

$$\overleftrightarrow{x B^l} := \text{span}(\{x\} \cup B^l)$$

containing $\{x\} \cup B^l$ is unique and has dimension $l+1$. It intersects A^k transversely in a 0-dimensional projective subspace, that is, a unique point $\Pi_{A^k, B^l}(x)$, that is:

$$\{\Pi_{A^k, B^l}(x)\} = \overleftrightarrow{x B^l} \cap A.$$

Perspectivities are projective mappings obtained as restrictions of projections:

EXERCISE 2.3.5. Let $A' \subset \mathbf{P}$ be a projective subspace of dimension k disjoint from B .

- Restriction $\Pi_{A, B}|_{A'}$ is a projective isomorphism $A' \rightarrow A$.
- Express an arbitrary projective isomorphism between projective subspaces as a composition of perspectivities.




FIGURE 2.1. A perspectivity between two lines l, l' in the plane. The undefined set of $\{O\}$ of the projection is the *center* of the perspectivity.

2.3.3. Locally projective maps. If P, P' are projective spaces and $U \subset P$ is an open set then a map $U \xrightarrow{f} P'$ is *locally projective* if for each component $U_i \subset U$ there exists a linear map

$$V(P) \xrightarrow{\tilde{f}_i} V(P')$$

such that the restrictions of $f \circ \Pi$ and $\Pi \circ \tilde{f}_i$ to $\Pi^{-1}U_i$ agree. Locally projective maps (and hence also locally affine maps) satisfy the *Unique Extension Property*: if $U \subset U' \subset P$ are open subsets of a projective space with U nonempty and U' connected, then any two locally projective maps $f_1, f_2 : U' \rightarrow P'$ which agree on U must be identical. (Compare §5.1.1.)

The passage between the geometry of P and the algebra of V is a “dictionary” between linear algebra and projective geometry. Linear maps and linear subspaces correspond geometrically to projective maps and projective subspaces: inclusions, intersections and linear spans correspond to incidence relations in projective geometry. Thus projective geometry lets us visually understand linear algebra and linear algebra enables to prove theorems in geometry by calculation.

EXERCISE 2.3.6. *Let $U \subset P$ be a connected open subset of a projective space of dimension greater than 1. Let $U \xrightarrow{f} P'$ be a local diffeomorphism. Then f is locally projective if and only if for each line $l \subset P$, the image $f(l \cap U)$ is a line.*

2.4. Affine patches

Let $H \subset \mathbb{P}$ be a projective hyperplane (projective subspace of codimension one). Then the complement $\mathbb{P} - H$ has a natural affine geometry, that is, is an affine space in a natural way. Indeed the group of projective automorphisms $\mathbb{P} \rightarrow \mathbb{P}$ leaving fixed each $x \in H$ and whose differential $T_x\mathbb{P} \rightarrow T_x\mathbb{P}$ is a volume-preserving linear automorphism is a vector group acting simply transitively on $A = \mathbb{P} \setminus H$. Moreover the subgroup of $\text{Aut}(\mathbb{P})$ leaving H invariant is $\text{Aff}(A)$. In this way affine geometry *embeds* in projective geometry.

Here is how it looks in terms of matrices. Let $A = \mathbb{R}^n$; then the affine subspace of

$$V = \text{Trans}(A) \oplus \mathbb{R} = \mathbb{R}^{n+1}$$

corresponding to A is $\mathbb{R}^n \times \{1\} \subset \mathbb{R}^{n+1}$, the point of A with *affine* or *inhomogeneous* coordinates (x^1, \dots, x^n) has homogeneous coordinates $[x^1, \dots, x^n, 1]$. Let $f \in \text{Aff}(E)$ be the affine transformation with linear part $A \in \text{GL}(n; \mathbb{R})$ and translational part $\mathbf{b} \in \mathbb{R}^n$, that is, $f(x) = Ax + \mathbf{b}$, is then represented by the $(n + 1)$ -square matrix

$$\begin{bmatrix} A & \mathbf{b} \\ 0 & 1 \end{bmatrix}$$

where \mathbf{b} is a column vector and 0 denotes the $1 \times n$ zero row vector.

2.4.1. Projective vector fields. In the affine space A^n , let

$$\mathfrak{B}(A^n) \longrightarrow A^n$$

denote the *bundle of bases*, more commonly known as the *affine frame bundle* over A^n : its fiber \mathfrak{B}_p over a point $p \in A^n$ consists of the set of bases for the tangent space $T_p A^n$. Using the simply transitive action of $\mathfrak{k}^n = \text{Trans}(A^n)$ on A^n , the total space $\mathfrak{B}(A^n)$ is a torsor for the affine automorphism group $\text{Aff}(A^n)$: an affine automorphism is determined uniquely by its action on a basepoint $p_0 \in A^n$ and a basis $\beta_0 \in \mathfrak{B}_{p_0}$ of $T_{p_0} A^n$. and every $(p, \beta) \in \mathfrak{B}$ is the image of (p_0, β_0) .

Let $g \in \text{Aut}(\mathbb{P}^n)$ be a projective automorphism. Fixing a basepoint p_0 and a basis β_0 of $T_{p_0} A^n$, let $h \in \text{Aff}(A^n)$ be the unique affine automorphism taking (p_0, β_0) to (p, β) . Then $h^{-1} \circ g$ is a projective automorphism fixing p_0 and acts identically on $T_{p_0} \mathbb{P}^n = T_{p_0} A^n$. In the affine chart A^{n+1} where p_0 is the origin, such a projective transformation is

defined in homogeneous coordinates by:

$$\begin{bmatrix} X^1 \\ \vdots \\ X^n \\ X^{n+1} \end{bmatrix} \mapsto \begin{bmatrix} X^1 \\ \vdots \\ X^n \\ \sum_{i=1}^n \xi_i X^i + X^{n+1} \end{bmatrix}$$

for a row vector ξ^\dagger for $\xi \in \mathbf{k}^n$, that is, by the block matrix

$$\begin{bmatrix} \mathbb{I}_n & \mathbf{0} \\ \xi & 1 \end{bmatrix}.$$

In affine coordinates such a transformation is given by

$$(x^1, \dots, x^n) \xrightarrow{g_\xi} \left(\frac{x^1}{1 + \sum_{i=1}^n \xi_i x^i}, \dots, \frac{x^n}{1 + \sum_{i=1}^n \xi_i x^i} \right).$$

EXERCISE 2.4.1. *Show that this group is isomorphic to a n -dimensional vector group, and g_ξ lies in the one-parameter group*

$$t \mapsto g_{t\xi}.$$

The corresponding vector field equals the product

$$-\sum_{i=1}^n \xi_i x^i \mathbf{R}$$

where \mathbf{R} is the radiant vector field defined in §1.5.2. Denote the Lie algebra of such vector fields by \mathfrak{g}_1 , the Lie algebra of parallel vector fields on \mathbf{A}^n by \mathfrak{g}_{-1} , and the Lie algebra of linear vector fields on \mathbf{A}^n by \mathfrak{g}_0 , show that, for $\lambda = 0, \pm 1$, the subalgebra \mathfrak{g}_λ of the Lie algebra \mathfrak{g} of projective vector fields equals the λ -eigenspace of $\text{Ad}\mathbf{R}$. Furthermore $[\mathfrak{g}_\lambda, \mathfrak{g}_\mu] \subset \mathfrak{g}_{\lambda+\mu}$ where $\mathfrak{g}_\nu := 0$ if $\nu \neq 0, \pm 1$. In particular as a vector space \mathfrak{g} decomposes as a direct sum

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \cong \mathbf{k}^n \oplus \mathfrak{gl}(n) \oplus \mathbf{k}^n.$$

Describe the corresponding group-theoretic decomposition of the projective automorphism group

$$\text{Aut}(\mathbf{P}^n) := \text{PGL}(n+1).$$

2.5. Classical projective geometry

This section surveys standard results in projective geometry. The *fundamental theorem of projective geometry* characterizes projective mappings. The *cross ratio* of four points is the fundamental invariant in one-dimensional projective geometry. The classical notion of




FIGURE 2.2. The four points on the horizontal line form a harmonic quadruple. Such a quadruple is characterized by having cross-ratio -1 .

harmonic sets is introduced, and is applied to the study of projective reflections and their products.

2.5.1. One-dimensional reflections. Let l be a projective line $x, z \in l$ be distinct points. Then there exists a unique reflection (a *harmonic homology* in classical terminology)

$$l \xrightarrow{\rho_{x,z}} l$$

whose fixed-point set equals $\{x, z\}$. We say that a pair of points y, w are *harmonic* with respect to x, z if $\rho_{x,z}$ interchanges them. In that case one can show that x, z are harmonic with respect to y, w . Furthermore this relation is equivalent to the existence of lines p, q through x and lines r, s through z such that

$$(6) \quad y = \overleftrightarrow{(p \cap r)(q \cap s)} \cap l$$

$$(7) \quad z = \overleftrightarrow{(p \cap s)(q \cap r)} \cap l$$

This leads to a projective-geometry construction of reflection, as follows. Let $x, y, z \in l$ be fixed; we seek the harmonic conjugate of y with respect to x, z , that is, the image $R_{x,z}(y)$. Erect arbitrary lines (in general position) p, q through x and a line r through z . Through y draw the line

$$l' := \overleftrightarrow{y (r \cap q)}$$

through $r \cap q$; join its intersection with p with z to form a line:

$$s = \overleftrightarrow{z (p \cap l')}.$$

Then $R_{x,z}(y)$ will be the intersection of s with l .




FIGURE 2.3. Non-Euclidean tessellations by equilateral triangles

EXERCISE 2.5.1. Consider the projective line $\mathbb{P}^1 = \mathbb{R} \cup \{\infty\}$. Show that for every rational number $x \in \mathbb{Q}$ there exists a sequence

$$x_0, x_1, x_2, x_3, \dots, x_n \in \mathbb{P}^1$$

such that:

- $x = \lim_{i \rightarrow \infty} x_i$;
- $\{x_0, x_1, x_2\} = \{0, 1, \infty\}$;
- For each $i \geq 3$, there is a harmonic quadruple (x_i, y_i, z_i, w_i) with

$$y_i, z_i, w_i \in \{x_0, x_1, \dots, x_{i-1}\}.$$

If x is written in reduced form p/q then what is the smallest n for which x can be reached in this way?

EXERCISE 2.5.2 (Synthetic arithmetic). Using the above synthetic geometry construction of harmonic quadruples, show how to add, subtract, multiply, and divide real numbers by a straightedge-and-pencil construction. In other words, draw a line l on a piece of paper and choose three points to have coordinates $0, 1, \infty$ on it. Choose arbitrary points corresponding to real numbers x, y . Using only a straightedge (not a ruler!) construct the points corresponding to

$$x + y, x - y, xy, \text{ and } x/y \text{ if } y \neq 0.$$

2.5.2. Fundamental theorem of projective geometry. One version of what is sometimes called the *fundamental theorem of projective geometry* is that the projective transformations (defined by linear transformations of the associated vector space) are precisely the transformations of projective space which preserve the ternary relation of collinearity (hence *collineations*). Collinearity is a special instance of the set of *incidence relations* between projective subspaces. For example, two distinct projective points p, q are incident to a unique projective line $\overleftrightarrow{p, q}$ and (p, q, r) is a collinear triple if and only if $r \in \overleftrightarrow{p, q}$. We do not develop this theory in detail, but refer to the texts of Berger [39, 38] and Coxeter [80] for discussion, and in particular the relation with automorphisms of the ground field k .

If $l \subset \mathbb{P}$ and $l' \subset \mathbb{P}'$ are projective lines, part of the *Fundamental Theorem of Projective Geometry* asserts that for given triples $x, y, z \in l$ and $x', y', z' \in l'$ of distinct points there exists a unique projective map

$$\begin{aligned} l &\xrightarrow{f} l' \\ x &\longmapsto x' \\ y &\longmapsto y' \\ z &\longmapsto z' \end{aligned}$$

If $w \in l$ then the *cross-ratio* $[w, x, y, z]$ is defined to be the image of w under the unique collineation $l \xrightarrow{f} \mathbb{P}^1$ with

$$\begin{aligned} x &\xrightarrow{f} 1 \\ y &\xrightarrow{f} 0 \\ z &\xrightarrow{f} \infty \end{aligned}$$

If $l = \mathbb{P}^1$, then this linear fractional transformation is:

$$f(w) := \frac{w - y}{w - z} \bigg/ \frac{x - y}{x - z}$$

so

$$(8) \quad [w, x, y, z] := \frac{w - y}{w - z} \bigg/ \frac{x - y}{x - z},$$

thus defining¹ the *cross-ratio*. Cross-ratio extends to quadruples of four points, of which at least three are distinct.

EXERCISE 2.5.3. *Let σ be a permutation on four symbols. Show that there exists a linear fractional transformation Φ_σ such that*

$$[x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}, x_{\sigma(4)}] = \Phi_\sigma([x_1, x_2, x_3, x_4]).$$

Determine which permutations leave the cross-ratio invariant.

The cross-ratio is a function of 4 variables, and therefore transforms under the action of the symmetric group \mathfrak{S}_4 on 4 symbols. The group \mathfrak{S}_4 is a split extension

$$\mathbb{Z}/2 \oplus \mathbb{Z}/2 \trianglelefteq \mathfrak{S}_4 \twoheadrightarrow \mathfrak{S}_3$$

where the normal subgroup $\mathbb{Z}/2 \oplus \mathbb{Z}/2$ consists of products of disjoint transpositions and a section $\mathfrak{S}_3 \hookrightarrow \mathfrak{S}_4$ corresponds to the inclusion $\{2, 3, 4\} \hookrightarrow \{1, 2, 3, 4\}$.

EXERCISE 2.5.4. *Show that the cross-ratio is invariant under the normal subgroup $\mathbb{Z}/2 \oplus \mathbb{Z}/2$, and transforms under $\mathfrak{S}_3 = \text{Aut}\{2, 3, 4\}$*

¹The literature has several variations of the cross-ratio; the version here is used by Veblen-Young [271], Kneebone-Semple [248], Coxeter [82], Ahlfors [3]. Other versions can be found in Goldman [124], Hubbard [154], Ovsienko-Tabachnikov [231].

by the rules, where $z = [z_1, z_2, z_3, z_4]$:

$$\begin{aligned} [z_1, z_2, z_4, z_3] &= 1/z \\ [z_1, z_3, z_2, z_4] &= 1 - z \\ [z_1, z_3, z_4, z_2] &= 1 - 1/z \\ [z_1, z_4, z_2, z_3] &= 1/(1 - z) \\ [z_1, z_4, z_3, z_2] &= z/(z - 1) \end{aligned}$$

This corresponds to an embedding $\mathfrak{S}_3 \hookrightarrow \mathrm{GL}(2, \mathbb{Z})$ taking

$$\begin{aligned} (34) &\longmapsto \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ (23) &\longmapsto \begin{bmatrix} -1 & 1 \\ 0 & 1 \end{bmatrix} \\ (234) &\longmapsto \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \\ (243) &\longmapsto \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix} \\ (24) &\longmapsto \begin{bmatrix} -1 & 0 \\ -1 & +1 \end{bmatrix} \end{aligned}$$

which induces an isomorphism $\mathfrak{S}_3 \cong \mathrm{GL}(2, \mathbb{Z}/2)$.

A pair $\{w, x\}$ is harmonic with respect to the pair $\{y, z\}$ (in which case we say that (x, y, w, z) is a *harmonic quadruple*) if and only if the cross-ratio $[x, y, w, z] = -1$.

EXERCISE 2.5.5. When $z = \infty$, the expression for the cross-ratio simplifies:

$$[x, y, w, \infty] := \frac{x - w}{y - w}$$

and defines a fundamental affine invariant.

- (x, y, z, ∞) is a harmonic quadruple if and only if y is the midpoint of \overline{xz} .
- Suppose $(x, y, w), (x', y', w')$ are ordered triples of distinct points of \mathbb{A}^1 . Show that $[x, y, w, \infty] = [x', y', w', \infty]$ if and only if $\exists g \in \mathrm{Aff}(\mathbb{A}^1)$ such that

$$\begin{aligned} x' &= g(x) \\ y' &= g(y) \\ w' &= g(w). \end{aligned}$$

The process of extending a triple of points on \mathbf{P}^1 to a harmonic quadruple is equivalent to applying a projective reflection in one pair to the remaining element. This process is called *harmonic subdivision*. Iterated harmonic subdivisions produce a countable dense subsets of \mathbf{P}^1 corresponding to the rational numbers $\mathbb{Q} \subset \mathbb{R}$. Such a subset is called a *harmonic net* (Coxeter [81], §3.5) or a *net of rationality* in Veblen-Young [271], p.84). Compare also Busemann-Kelly [54], §I.6. These ideas provide an approach to the fundamental theorem:

EXERCISE 2.5.6. Let $\mathbf{P}^1 \xrightarrow{f} \mathbf{P}^1$ be a homeomorphism. Show that the following conditions are equivalent:

- f is projective;
- f preserves harmonic quadruples
- f preserves cross-ratios, that is, for all quadruples (x, y, w, z) , the cross-ratios satisfy

$$[f(x), f(y), f(w), f(z)] = [x, y, w, z].$$

Determine the weakest hypothesis on f to obtain these conditions.

2.5.3. Distance via cross-ratios. For later use in §12.1, here are some explicit formulas for cross-ratios on 1-dimensional spaces \mathbb{R}^+ and \mathbf{I} . Their infinitesimal forms yield the Poincaré metrics on \mathbb{R}^+ and \mathbf{I} .

EXERCISE 2.5.7. (Parameters on the positive ray and unit intervals)

$$[0, e^a, e^b, \infty] = [-1, \tanh(a/2), \tanh(b/2), 1] = e^{b-a}$$

2.5.4. Products of reflections. If ϕ, ψ are collineations, each of which fix a point $O \in \mathbf{P}$, their composition $\phi\psi = \phi \circ \psi$ fixes O . In particular its derivative

$$D(\phi\psi)_O = (D\phi)_O \circ (D\psi)_O$$

acts linearly on the tangent space $T_O\mathbf{P}$. We consider the case when ϕ, ψ are reflections in \mathbf{P}^2 . As in Exercise 2.3.2, a reflection ϕ is completely determined by its set $\text{Fix}(\phi)$ which consists of a point p_ϕ and a line l_ϕ such that $p_\phi \notin l_\phi$. Define

$$O := l_\phi \cap l_\psi$$

and \mathbf{P}_O the projective line whose points are the lines incident to O .

EXERCISE 2.5.8. Let ρ denote the cross-ratio of the four lines

$$l, \overleftrightarrow{Op}, \overleftrightarrow{Op'}, l'$$

as elements of \mathbf{P}_O .

- The linear automorphism

$$\mathbb{T}_O\mathbb{P} \xrightarrow{D(\phi\psi)_O} \mathbb{T}_O\mathbb{P}$$

of the tangent space $\mathbb{T}_O\mathbb{P}$ leaves invariant a positive definite inner product \mathfrak{g}_O on $\mathbb{T}_O\mathbb{P}$.

- Furthermore $D(\phi\psi)_O$ represents a rotation of angle θ in the tangent space $\mathbb{T}_O(\mathbb{P})$ with respect to \mathfrak{g}_O if and only if

$$\rho = \frac{1}{2}(1 + \cos \theta)$$

for $0 < \theta < \pi$ and is a rotation of angle π (that is, an involution) if and only if $p \in l'$ and $p' \in l$.

2.6. Asymptotics of projective transformations

We shall be interested in the singular projective transformations since they occur as limits of nonsingular projective transformations. The collineation group $\mathbf{Aut}(\mathbb{P})$ of $\mathbb{P} = \mathbb{P}^n$ is a large noncompact group which is naturally embedded in the projective space $\mathbf{End}(\mathbb{P})$ as an open dense subset as in Exercise 2.3.4. Thus understanding precisely what it means for a sequence of collineations to converge to a (possibly singular) projective transformation is crucial.

A *singular projective transformation* of \mathbb{P} is a projective map f defined on the complement of a projective subspace $\mathbb{U}(f) \subset \mathbb{P}$, called the *undefined subspace* of f and taking values in a projective subspace $\mathcal{R}(f) \subset \mathbb{P}$, called the *range* of f . Furthermore

$$\dim \mathbb{P} = \dim \mathbb{U}(f) + \dim \mathcal{R}(f) + 1.$$

PROPOSITION 2.6.1. *Let $g_m \in \mathbf{Aut}(\mathbb{P})$ be a sequence of collineations of \mathbb{P} and let $g_\infty \in \mathbf{End}(\mathbb{P})$. Then the sequence g_m converges to g_∞ in $\mathbf{End}(\mathbb{P})$ if and only if the restrictions $g_m|_K$ converge uniformly to $g_\infty|_K$ for all compact sets $K \subset \subset \mathbb{P} - \mathbb{U}(g_\infty)$.*

Convergence in $\mathbf{End}(\mathbb{P})$ may be described as follows. Let $\mathbb{P} = \mathbb{P}(\mathbb{V})$ where $\mathbb{V} \cong \mathbb{R}^{n+1}$ is a vector space. Then $\mathbf{End}(\mathbb{P})$ is the projective space associated to the vector space $\mathbf{End}(\mathbb{V})$ of $(n+1)$ -square matrices. If $a = (a_j^i) \in \mathbf{End}(\mathbb{V})$ is such a matrix, let

$$\|a\| = \sqrt{\mathrm{Tr}(aa^\dagger)} = \sqrt{\sum_{i,j=1}^{n+1} |a_j^i|^2}$$

denote its Euclidean norm; projective endomorphisms then correspond to matrices a with $\|a\| = 1$, uniquely determined up to the antipodal

map $a \mapsto -a$. The following lemma will be useful in the proof of Proposition 2.6.1.

LEMMA 2.6.2. *Let \mathbf{V}, \mathbf{V}' be vector spaces and let $\mathbf{V} \xrightarrow{\tilde{f}_n} \mathbf{V}'$ be a sequence of linear maps converging to $\mathbf{V} \xrightarrow{\tilde{f}_\infty} \mathbf{V}'$. Let $\tilde{K} \subset \mathbf{V}$ be a compact subset of $\mathbf{V} - \text{Ker}(\tilde{f}_\infty)$. Define:*

$$\begin{aligned} \mathbf{V} &\xrightarrow{f_i} \mathbf{V}' \\ x &\mapsto \frac{\tilde{f}_i(x)}{\|\tilde{f}_i(x)\|}. \end{aligned}$$

Then f_n converges uniformly to f_∞ on \tilde{K} as $n \rightarrow \infty$.

PROOF. Choose $C > 0$ such that $C \leq \|\tilde{f}_\infty(x)\| \leq C^{-1}$ for $x \in \tilde{K}$. For $\epsilon > 0$, $\exists N$ such that if $n > N$, then $\forall x \in \tilde{K}$,

$$(9) \quad \begin{aligned} \|\tilde{f}_\infty(x) - \tilde{f}_n(x)\| &< \frac{C\epsilon}{2}, \\ \left| 1 - \frac{\tilde{f}_n(x)}{\|\tilde{f}_\infty(x)\|} \right| &< \frac{\epsilon}{2}. \end{aligned}$$

Let $x \in \tilde{K}$. Then

$$\begin{aligned} \|f_n(x) - f_\infty(x)\| &= \left\| \frac{\tilde{f}_n(x)}{\|\tilde{f}_n(x)\|} - \frac{\tilde{f}_\infty(x)}{\|\tilde{f}_\infty(x)\|} \right\| \\ &= \frac{1}{\|\tilde{f}_\infty(x)\|} \left\| \frac{\|\tilde{f}_\infty(x)\|}{\|\tilde{f}_n(x)\|} \tilde{f}_n(x) - \tilde{f}_\infty(x) \right\| \\ &\leq \frac{1}{\|\tilde{f}_\infty(x)\|} \left(\left\| \frac{\|\tilde{f}_\infty(x)\|}{\|\tilde{f}_n(x)\|} \tilde{f}_n(x) - \tilde{f}_n(x) \right\| \right. \\ &\quad \left. + \|\tilde{f}_n(x) - \tilde{f}_\infty(x)\| \right) \\ &= \left| 1 - \frac{\|\tilde{f}_n(x)\|}{\|\tilde{f}_\infty(x)\|} \right| + \frac{1}{\|\tilde{f}_\infty(x)\|} \|\tilde{f}_n(x) - \tilde{f}_\infty(x)\| \\ &< \frac{\epsilon}{2} + C^{-1} \left(\frac{C\epsilon}{2} \right) = \epsilon \quad (\text{by (9)}), \end{aligned}$$

completing the proof of Lemma 2.6.2. \square

PROOF OF PROPOSITION 2.6.1. Suppose g_m is a sequence of locally projective maps defined on a connected domain $\Omega \subset \mathbf{P}$ converging

uniformly on all compact subsets of Ω to a map

$$\Omega \xrightarrow{g_\infty} \mathbf{P}'.$$

Lift g_∞ to a linear transformation \tilde{g}_∞ of norm 1, and lift g_m to linear transformations \tilde{g}_m , also linear transformations of norm 1, converging to \tilde{g}_∞ . Then

$$g_m \longrightarrow g_\infty$$

in $\text{End}(\mathbf{P})$. Conversely if $g_m \longrightarrow g_\infty$ in $\text{End}(\mathbf{P})$ and

$$K \subset \mathbf{P} - \mathbb{U}(g_\infty),$$

choose lifts as above and $\tilde{K} \subset\subset V$ such that $\Pi(\tilde{K}) = K$. By Lemma 2.6.2,

$$\tilde{g}_m / \|\tilde{g}_m\| \rightrightarrows \tilde{g}_\infty / \|\tilde{g}_\infty\|$$

on \tilde{K} . Hence $g_m|_K \rightrightarrows g_\infty|_K$, completing the proof of Proposition 2.6.1. \square

2.6.1. Some examples. Let us consider a few examples of this convergence. Consider the case first when $n = 1$. Let $\lambda_m \in \mathbb{R}$ be a sequence converging to $+\infty$ and consider the projective transformations given by the diagonal matrices

$$g_m = \begin{bmatrix} \lambda_m & 0 \\ 0 & (\lambda_m)^{-1} \end{bmatrix}$$

Then $g_m \longrightarrow g_\infty$ where g_∞ is the singular projective transformation corresponding to the matrix

$$g_\infty = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

— this singular projective transformation is undefined at $\mathbb{U}(g_\infty) = \{[0, 1]\}$; every point other than $[0, 1]$ is sent to $[1, 0]$. It is easy to see that a singular projective transformation of \mathbf{P}^1 is determined by the ordered pair of points $(\mathbb{U}(f), \mathcal{R}(f))$. Note that in the next example, the two points $\mathbb{U}(\phi_\infty), \mathcal{R}(\phi_\infty)$ coincide.

EXERCISE 2.6.3. Consider the sequence of projective transformations of \mathbf{P}^1

$$\phi_n(x) := \frac{x}{1 - nx}, \quad \text{as } n \longrightarrow +\infty$$

- Show that the pointwise limit equals the constant function 0:

$$\lim_{n \rightarrow \infty} \phi_n(x) = 0, \quad \forall x \in \mathbf{P}^1.$$

- Show that ϕ_n does not converge uniformly on any subset $S \subset \mathbf{P}^1$ which contains an infinite subsequence $s_j > 0$ with $s_j \searrow 0$ as $j \longrightarrow \infty$.

- Show that ϕ_n does converge to the singular projective transformation ϕ_∞ defined on the complement of $\mathbb{U}(\phi_\infty) = \{0\}$ and has constant value 0.
- Use this idea, together with the techniques involved in the proof of Lemma 2.6.2, to prove a statement converse to Proposition 2.6.1: If $g_n \in \text{Aut}(\mathbb{P})$ is a sequence of projective transformations converging to a singular projective transformation $g_\infty \in \text{End}(\mathbb{P})$, then, for any open subset $S \subset \mathbb{P}$ which meets $\mathbb{U}(g_\infty)$, the restrictions $g_n|_S$ do not converge uniformly.

EXERCISE 2.6.4. (1) The projective group $\text{PGL}(2, \mathbb{R}) = \text{Aut}(\mathbb{RP}^1)$ is an open dense subset of $\text{End}(\mathbb{RP}^1) \approx \mathbb{RP}^3$. Its complement naturally identifies with the Cartesian product $\mathbb{RP}^1 \times \mathbb{RP}^1$ under the correspondence

$$\begin{aligned} \text{End}(\mathbb{RP}^1) \setminus \text{Aut}(\mathbb{RP}^1) &\longleftrightarrow \mathbb{RP}^1 \times \mathbb{RP}^1 \\ [f] &\longleftrightarrow (\mathbb{U}(f), \mathcal{R}(f)) \end{aligned}$$

- (2) Prove the analogous statements for $\text{PGL}(2, \mathbb{C})$ and \mathbb{CP}^1 , that is, when \mathbb{R} is replaced by \mathbb{C} .
- (3) Show that if $\Gamma < \text{PGL}(2, \mathbb{C})$ is a discrete subgroup with limit set $\Lambda \subset \mathbb{CP}^1$, then $\bar{\Gamma} \setminus \Gamma$ identifies with $\Lambda \times \Lambda \subset \mathbb{CP}^1 \times \mathbb{CP}^1$.

2.6.2. Higher dimensional projective maps. More interesting phenomena arise when $n = 2$. Let $g_m \in \text{Aut}(\mathbb{P}^2)$ be a sequence of diagonal matrices

$$\begin{bmatrix} \lambda_m & 0 & 0 \\ 0 & \mu_m & 0 \\ 0 & 0 & \nu_m \end{bmatrix}$$

where $0 < \lambda_m < \mu_m < \nu_m$ and $\lambda_m \mu_m \nu_m = 1$. Corresponding to the three eigenvectors (the coordinate axes in \mathbb{R}^3) are three fixed points

$$p_1 = [1, 0, 0], \quad p_2 = [0, 1, 0], \quad p_3 = [0, 0, 1].$$

They span three invariant lines

$$l_1 = \overleftarrow{p_2 p_3}, \quad l_2 = \overleftarrow{p_3 p_1}, \quad l_3 = \overleftarrow{p_3 p_1}.$$

Since $0 < \lambda_m < \mu_m < \nu_m$, the collineation has an repelling fixed point at p_1 , a saddle point at p_2 and an attracting fixed point at p_3 . Points on l_2 near p_1 are repelled from p_1 faster than points on l_3 and points on l_2 near p_3 are attracted to p_3 more strongly than points on l_1 . Suppose that g_m does not converge to a nonsingular matrix; it follows that $\nu_m \rightarrow +\infty$ and $\lambda_m \rightarrow 0$ as $m \rightarrow \infty$. Suppose that $\mu_m/\nu_m \rightarrow \rho$;

then g_m converges to the singular projective transformation g_∞ determined by the matrix

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & \rho & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

which, if $\rho > 0$, has undefined set $\mathbb{U}(g_\infty) = p_1$ and range l_1 ; otherwise $\mathbb{U}(g_\infty) = l_2$ and $\text{Image}(g_\infty) = p_2$.

2.6.3. Limits of similarity transformations. Convergence to singular projective transformations is perhaps easiest for translations of affine space, or, more generally, Euclidean isometries.

EXERCISE 2.6.5. *Suppose $g_m \in \text{Isom}(\mathbb{E}^n)$ be a divergent sequence of Euclidean isometries. Show that $\exists p \in \mathbb{P}_\infty^{n-1}$ a subsequence g_{m_k} , that $g_{m_k}|_K \rightrightarrows p$ for every compact $K \subset \subset \mathbb{E}^n$.*

Indeed the boundary of the translation group \mathbb{V} of \mathbb{A}^n is the projective space \mathbb{P}_∞^{n-1} . More generally the boundary of $\text{Isom}(\mathbb{E}^n)$ identifies with \mathbb{P}_∞^{n-1} .

EXERCISE 2.6.6. *Suppose $g_m \in \text{Sim}(\mathbb{E}^n)$ be a divergent sequence of similarities of Euclidean space. Then \exists a subsequence g_{m_k} , and a point*

$$p \in \mathbb{E}^n \coprod \mathbb{P}_\infty^{n-1}$$

such that one of the three possibilities occur:

- $p \in \mathbb{P}_\infty^{n-1}$ and $g_{m_k}|_K \rightrightarrows p$, $\forall K \subset \subset \mathbb{E}^n$;
- $p \in \mathbb{P}_\infty^{n-1}$ and $\exists q \in \mathbb{E}^n$ such that

$$g_{m_k}|_K \rightrightarrows p, \forall K \subset \subset \mathbb{E}^n \setminus \{q\};$$

- $p \in \mathbb{E}_\infty^n$ and

$$g_{m_k}|_K \rightrightarrows p, \forall K \subset \subset \mathbb{E}^n.$$

The scale factor homomorphism $\text{Sim}(\mathbb{E}^n) \xrightarrow{\lambda} \mathbb{R}^+$ defined in §1.4.1 of Chapter 1 controls the asymptotics of linear similarities. The two latter cases occur when $\lim_{k \rightarrow \infty} \lambda(g_{m_k}) = \infty$ and $\lim_{k \rightarrow \infty} \lambda(g_{m_k}) = 0$, respectively.

These results will be used in Fried's classification of closed similarity manifolds (§11.4 of Chapter 11).

2.6.4. Normality domains. Convergence to singular projective transformations closely relates to the notion of *normality*, introduced by Kulkarni-Pinkall [194], and extending the classical notion of *normal families* in complex analysis. Let G be a group acting on a space X strongly effectively. A point $x \in X$ is a *point of normality* with respect to G if and only if x admits an open neighborhood W such the the set of restrictions

$$G|_W := \{g|_W \mid g \in G\}$$

is a compact subset of $\text{Map}(W, X)$ with respect to the compact-open topology on $\text{Map}(W, X)$. (This means that $G|_W$ is a *normal family* in the sense of Montel.) Denote the set of points of normality by $\text{Nor}(G, X)$. Clearly $\text{Nor}(G, X)$ is a G -invariant open subset of X , called the *normality domain*.

PROPOSITION 2.6.7. *Suppose that $\Gamma < \text{Aut}(\mathbb{P})$ is a discrete group of collineations of a projective space \mathbb{P} . Let $\bar{\Gamma} \subset \text{End}(\mathbb{P})$ denote its closure in the set of singular projective transformations. Then the normality domain $\text{Nor}(\Gamma, X)$ consists of the complement*

$$\mathcal{U}_\Gamma := X \setminus \bigcup_{\bar{\gamma} \in \bar{\Gamma}} \mathbb{U}(\bar{\gamma})$$

in \mathbb{P} of the union $\bigcup_{\bar{\gamma} \in \bar{\Gamma}} \mathbb{U}(\bar{\gamma})$.

PROOF. Observe first that $\bigcup_{\bar{\gamma} \in \bar{\Gamma}} \mathbb{U}(\bar{\gamma}) \subset\subset X$ since each projective subspace $\mathbb{U}(\bar{\gamma}) \subset\subset X$. Since the parameter space $\bar{\Gamma}$ is compact. Thus it's closed, and its complement

$$\mathcal{U}_\Gamma := X \setminus \bigcup_{\bar{\gamma} \in \bar{\Gamma}} \mathbb{U}(\bar{\gamma})$$

is open. We claim that $\mathcal{U}_\Gamma = \text{Nor}(\Gamma, X)$.

We first show any point $x \in \mathcal{U}_\Gamma$ is a point of normality. To this end, we show that. set of restrictions $\Gamma|_{\mathcal{U}_\Gamma}$ is precompact in $\text{Map}(\mathcal{U}_\Gamma, X)$. This follows immediately from Proposition 2.6.1 as follows. Consider an infinite sequence $\gamma_n \in \Gamma$. Proposition 2.6.1 ensures a subsequence γ_n and a singular projective transformation $\bar{\gamma}_\infty \in \text{End}(\mathbb{P})$, such that

$$\gamma_n|_K \rightrightarrows \bar{\gamma}_\infty|_K, \quad \forall K \subset\subset \mathcal{U}_\Gamma$$

as desired. (Since $K \cap \mathbb{U}(\bar{\gamma}_\infty) = \emptyset$, the restriction $\bar{\gamma}_\infty|_K$ is defined.)

Conversely suppose that $x \in \mathbb{U}(\bar{\gamma})$ for some $\bar{\gamma} \in \bar{\Gamma}$. Choose a sequence $g_n \in \Gamma$ converging to $\bar{\gamma}$, and a precompact open neighborhood $S \ni x$. By Exercise 2.6.3, the restrictions $g_n|_S$ do not converge uniformly, and the restrictions to the closure $\bar{S} \subset X$ do not converge uniformly. Thus $x \notin \text{Nor}(\Gamma, X)$ as claimed. \square

CHAPTER 3

Duality and non-Euclidean geometry

The axiomatic development of projective geometry enjoys a basic symmetry: In \mathbf{P}^2 , a pair of distinct points lie on a unique line and a pair of distinct lines meet in a unique point. Consequently any statement about the geometry of \mathbf{P}^2 can be *dualized* by replacing “point” by “line,” “line” by “point,” “collinear” with “concurrent,” etc; as long as it is done in a *completely consistent fashion*.

One of the oldest nontrivial theorems of projective geometry is Pappus’ theorem (300 A.D.), asserting that if $l, l' \subset \mathbf{P}^2$ are distinct lines and $A, B, C \in l$ and $A', B', C' \in l'$ are triples of distinct points, then the three points

$$\overleftrightarrow{AB'} \cap \overleftrightarrow{A'B}, \quad \overleftrightarrow{BC'} \cap \overleftrightarrow{B'C}, \quad \overleftrightarrow{CA'} \cap \overleftrightarrow{C'A}$$

are collinear. Dual to Pappus’ theorem is the following: if $p, p' \in \mathbf{P}^2$ are distinct points and a, b, c are distinct lines all passing through p and a', b', c' are distinct lines all passing through p' , then the three lines

$$\overleftrightarrow{(a \cap b) \ (a' \cap b)}, \quad \overleftrightarrow{(b \cap c) \ (b' \cap c)}, \quad \overleftrightarrow{(c \cap a) \ (c' \cap a)}$$

are concurrent. (According to [80], Hilbert observed that Pappus’ theorem is equivalent to the commutative law of multiplication.)

This chapter introduces the projective models for elliptic and hyperbolic geometry through the classical notion of *polarities*. We begin by working over both \mathbb{R} and \mathbb{C} , but specialize to the case $\mathbf{k} = \mathbb{R}$ when we discuss important cases of polarities:

- *Elliptic* polarities, corresponding to definite symmetric bilinear forms, and leading to elliptic (spherical) geometry;
- *Hyperbolic* polarities, corresponding to Lorentzian bilinear forms, and leading to hyperbolic geometry;
- *Null* polarities, corresponding to nondegenerate skew-symmetric bilinear forms, and leading to contact projective geometry (in odd dimensions). (Compare Semple and Kneebone [248].)

See Ratcliffe [239] for a good elementary treatment of the Beltrami-Klein model for hyperbolic geometry, as well as Thurston [266], §2.3.

The various derivations of hyperbolic geometry are aided by symmetry. The Beltrami-Klein metric generalizes to the Hilbert metric on convex domains. For an extensive modern discussion of Hilbert geometry, see Papadopoulos-Troyanov [233]. These, in turn, are special cases of intrinsic metrics, which we introduce here, and discuss later in the context of affine and projective structures on manifolds.

3.1. Dual projective spaces

In terms of our projective geometry/linear algebra dictionary, projective duality translates into duality between vector spaces as follows. Let $P = P(W)$ be a projective space associated to the vector space W . A nonzero linear functional

$$W \xrightarrow{\psi} k$$

defines a projective hyperplane H_ψ in P ; two such functionals define the same hyperplane if and only if they are *projectively equivalent*, that is, they differ by scalar multiplication by a nonzero scale factor. Equivalently, they determine the same line in the vector space W^* dual to W .

The *projective space P^* dual to P* consists of lines in the dual vector space W^* , which correspond to hyperplanes in P . The line joining two points in P^* corresponds to the intersection of the corresponding hyperplanes in P , and a hyperplane in P^* corresponds to a point in P .

EXERCISE 3.1.1. *Show that an n -dimensional projective space enjoys a natural correspondence*

$$\{k - \text{dimensional subspaces of } P\} \longleftrightarrow \{l - \text{dimensional subspaces of } P^*\}$$

where $k + l = n - 1$.

Since vector spaces of the same dimension are isomorphic, a projective space P is projectively isomorphic to the dual of P^* , but *in many different ways*.

Let $P \xrightarrow{f} P'$ be a projective map. Then for each hyperplane $H' \subset P'$ the preimage $f^{-1}(H')$ is a hyperplane in P . This defines a projective map $(P')^* \xrightarrow{f^*} P^*$.

Here is the detailed construction. Suppose that f is defined by a linear mapping of vector spaces $W \xrightarrow{F} W'$, where P, P' projectivize W, W' respectively. Since f is defined on all of P , the linear map F

is injective. The projective hyperplane H corresponds to the linear hyperplane

$$S := \text{Ker}(\psi) \subset W,$$

for some nonzero covector $\psi \in W^*$. Since $F^{-1}(S) = \text{Ker}(\psi \circ F)$, the preimage $f^{-1}(H)$ is the projective hyperplane defined by the covector $\psi \circ F \in (W')^*$.

EXERCISE 3.1.2. *If f is the projectivization of a linear map $W \xrightarrow{F} W'$, show that f^* is the projectivization of the dual map $(W')^* \xrightarrow{F^*} W^*$ given by matrix transpose. That is, represent F by a matrix M by choosing bases of W and W' respectively. Show that the matrix representing F^* in the respective dual bases of W^* and $(W')^*$ is the transpose M^\dagger of the matrix M .*

3.2. Correlations and polarities

DEFINITION 3.2.1. *Let P be a projective space. A correlation of P is a projective map $P \rightarrow P^*$.*

That is, a correlation associates to every projective point p a projective hyperplane $C(p)$ in such a way to *preserve incidences*: if $p, q, r \in P$ are distinct projective points, then p, q, r are collinear if and only if the projective subspace $C(p) \cap C(q) \cap C(r)$ has codimension two (not three).

EXERCISE 3.2.2. *Let W be a vector space such that $P = P(W)$. Correlations of P identify with projective equivalence classes of nondegenerate bilinear forms*

$$W \times W \rightarrow \mathbf{k},$$

or, equivalently, linear isomorphisms $W \rightarrow W^$.*

3.2.1. Example: Elliptic geometry. Here is an example of a correlation, corresponding to the standard Euclidean inner product on $W = \mathbb{R}^3$. This correlation associates a point $p = [\mathbf{v}] \in \mathbb{R}P^2$ the projective line p^* corresponding to the orthogonal complement \mathbf{v}^\perp . The corresponding linear isomorphism $W \rightarrow W^*$ is the usual *transpose* operation, interchanging column vectors and row vectors.

EXERCISE 3.2.3. *p and p^* are never incident.*

An important property of this correlation is that it is *self-inverse* in the following sense:

EXERCISE 3.2.4. *Let $p \mapsto p^*$ be the correlation defined as above. If $l, m \in P^*$ are distinct projective lines, with*

$$l = p^*, \quad m = q^*,$$

for respective projective points p, q , then

$$(l \cap m)^* = \overleftrightarrow{pq}.$$

DEFINITION 3.2.5. A self-inverse correlation is called a polarity.

3.2.2. Elliptic polarities and elliptic geometry. Our interest in projective correlations stems from their use to define models of non-Euclidean geometry. The above correlation defines a distance d on \mathbb{RP}^2 , making (\mathbb{RP}^2, d) into a metric space, called the *elliptic plane*. It is a basic example of *non-Euclidean geometry*.

For any two points $p, q \in \mathbb{RP}^2$, define the distance $d(p, q)$ as follows. If $p = q$, define their distance to be zero. Otherwise, p, q span a unique projective line $l := \overleftrightarrow{pq}$, and we extend (p, q) to a quadruple on l and compute its cross-ratio. Since p and p^* (respectively q and q^*) are not incident, $p^* \cap l$ and $q^* \cap l$ are points on l , denoted p', q' respectively. Define:

$$d(p, q) := \tan^{-1} \sqrt{[p, q, p', q']}.$$

EXERCISE 3.2.6. (\mathbb{RP}^2, d) is a metric space satisfying:

- $\text{SO}(3)$ acts isometrically and transitively on this metric space.
- (Busemann-Kelly [54], Exercise 38.1, p. 237) This metric space arises from a Riemannian structure, defined in an affine patch with coordinates $(x, y) \in \mathbb{R}^2$, by the metric tensor

$$\frac{dx^2 + dy^2 + (x dy - y dx)^2}{x^2 + y^2 + 1}$$

- Relate this metric space to the Euclidean unit sphere in \mathbb{R}^3 .
- The Gaussian curvature equals $+1$.
- The geodesics are exactly the projective lines.

This construction extends to higher dimensions, and the metric geometry is *Elliptic Geometry*. The corresponding Riemannian metric is called the *Fubini-Study metric*. In 1866 Beltrami showed that the only Riemannian metrics on domains in \mathbb{P}^n where the geodesics are straight line segments are (up to a collineation and change of scale factor) Euclidean metrics and these two metrics. Hilbert's fourth problem was to determine all metric space structures on domains in \mathbb{P}^n whose geodesics are straight line segments. There are many unusual such metrics, see Busemann-Kelly [54], and Pogorelov [236]. For more general discussion compare Coxeter [82] and Goldman [124], §1.3.

3.2.3. Polarities. The correlation defining elliptic geometry is an example of a *polarity*, which is self-inverse in the sense of Exercise 3.2.4. Polarities give rise to more general and fascinating geometries, the most important being *hyperbolic non-Euclidean geometry*.

We discuss polarities in general. For expository simplicity, we henceforth restrict to the case $k = \mathbb{R}$, although the complex case is quite interesting and basic.

Using the dictionary between projective geometry and linear algebra, one sees that if W is the vector space corresponding to $P = P(W)$, then $P^* = P(W^*)$ and a correlation θ is realized as a linear isomorphism $W \xrightarrow{\tilde{\theta}} W^*$, which is uniquely determined up to homotheties. Linear maps $W \xrightarrow{\tilde{\theta}} W^*$, correspond to bilinear forms

$$W \times W \xrightarrow{B_{\tilde{\theta}}} \mathbb{R}$$

under the correspondence

$$\tilde{\theta}(v)(w) = B_{\tilde{\theta}}(v, w)$$

and $\tilde{\theta}$ is an isomorphism if and only if $B_{\tilde{\theta}}$ is nondegenerate. Thus correlations can be interpreted analytically as projective equivalence classes of nondegenerate bilinear forms.

EXERCISE 3.2.7. *A correlation θ is a polarity (that is, θ is self-inverse) if and only if a corresponding bilinear form B_{θ} is either symmetric or skew-symmetric.*

Let θ be a polarity on P . A point $p \in P$ is *conjugate* if it is incident to its polar hyperplane, that is, if $p \in \theta(p)$. By our dictionary we see that the conjugate points of a polarity correspond to *null vectors* of the associated quadratic form, that is, to nonzero vectors $v \in W$ such that $B_{\theta}(v, v) = 0$. A polarity is said to be *elliptic* if it admits no conjugate points, which are definite. The polarity of § 3.2.1 which associates to a point $p = [v] \in P^2$ the line $P(v^\perp) \subset P^2$ is an elliptic polarity (compare Exercise 3.2.3).

At the other extreme, a polarity is *null* if and only if every point is conjugate.

EXERCISE 3.2.8. *Null polarities of a projective space P correspond to nondegenerate skew-symmetric bilinear forms on the vector space W , where $P = P(W)$. A projective space P admits a null polarity if and only if $\dim P$ is odd.*

Exercise 6.1.2 develops the theory of geometric structures (called *contact projective structures*) related to null polarities.

3.2.4. Quadrics. As in §3.2.2 an elliptic polarity describes the structure of elliptic geometry on $\mathbb{R}P^n$. In particular it induces a Riemannian structure on $\mathbb{R}P^n$, namely the *Fubini-Study metric*. This construction readily generalizes to polarities which are not null, and correspond to nondegenerate symmetric bilinear forms, which are *indefinite*. The isomorphism class of such a bilinear form is determined by its *signature* (p, q) where p, q are positive integers with $p + q = n + 1$. A standard example is the $B_{p,q}$ defined by:

$$(x, y) \xrightarrow{B_{p,q}} x^1 y^1 + \cdots + x^p y^p - x^{p+1} y^{p+1} - \cdots - x^{n+1} y^{n+1}$$

corresponding to the quadratic form

$$x \longmapsto (x^1)^2 + \cdots + (x^p)^2 - (x^{p+1})^2 - \cdots - (x^{p+q})^2.$$

EXERCISE 3.2.9. Let $B = B_\theta$ be a nondegenerate symmetric bilinear form on the vector space W of signature (p, q) where $p + q = n + 1 = \dim(W)$. Consider the quadric hypersurfaces:

$$Q^- := \{\mathbf{w} \in W \mid B(\mathbf{w}, \mathbf{w}) = -1\}$$

$$Q^0 := \{\mathbf{w} \in W \mid B(\mathbf{w}, \mathbf{w}) = 0\}$$

$$Q^+ := \{\mathbf{w} \in W \mid B(\mathbf{w}, \mathbf{w}) = 1\}.$$

Their respective projectivizations in $P = P(W)$ decompose P as a disjoint union:

$$P = P(Q^-) \amalg P(Q^0) \amalg P(Q^+).$$

- If $p = 0$, then B is negative definite, and $P(Q^-) = P(Q^0) = \emptyset$ and $P(Q^+) = P$.
- If $0 < p, q < n + 1$, then B is indefinite and $P(Q^-), P(Q^0), P(Q^+)$ are each connected and nonempty. Indeed,

$$P(Q^0) \approx S^{p-1} \times S^{q-1}$$

if $p, q > 1$ and

$$P(Q^0) \approx S^{q-1}$$

if $p = 1$.

- If $p = n + 1$, then B is positive definite, $P(Q^+) = P(Q^0) = \emptyset$ and $P(Q^-) = P$.

PROPOSITION 3.2.10. $P(Q^+)$ (respectively $P(Q^-)$) admits a $PO(p, q)$ -invariant pseudo-Riemannian structure of signature $(p-1, q)$ (respectively $(p, q+1)$).

PROOF. The symmetric bilinear form \mathbf{Q} on W induces an invariant pseudo-Riemannian structure on \mathbf{Q}^+ . Let $\mathbf{v} \in \mathbf{Q}^+$. Then the tangent space $\mathbf{T}_{\mathbf{v}}\mathbf{Q}$ identifies with the orthogonal complement $\mathbf{v}^\perp \subset W$. The restriction of the pseudo-Riemannian structure to \mathbf{v}^\perp has signature $(p-1, q)$ and is evidently $\mathrm{PO}(p, q)$ -invariant. The case of \mathbf{Q}^- is completely analogous. \square

The set of conjugate points of a polarity θ is the quadric $\mathbf{P}(\mathbf{Q}_\theta^0)$, comprising points $[\mathbf{w}] \in \mathbf{P}$ with $\mathbf{B}_\theta(\mathbf{w}, \mathbf{w}) = 0$.

The quadric \mathbf{Q} determines the polarity θ as follows.

For brevity we consider only the case $q = 1$, in which case the complement $\mathbf{P} \setminus \mathbf{Q}$ has two components, a convex component

$$\Omega = \{[x^0 : x^1 : \cdots : x^n] \mid (x^0)^2 - (x^1)^2 - \cdots - (x^n)^2 < 0\}$$

and a nonconvex component

$$\Omega^\dagger = \{[x^0 : x^1 : \cdots : x^n] \mid (x^0)^2 - (x^1)^2 - \cdots - (x^n)^2 > 0\}$$

diffeomorphic to the total space of the tautological line bundle over \mathbf{P}^{n-1} (for $n = 2$ this is a Möbius band). If $x \in Q$, let $\theta(x)$ denote the hyperplane tangent to Q at x . If $x \in \Omega^\dagger$ the points of Q lying on tangent lines to Q containing x all lie on a hyperplane which is $\theta(x)$. If $H \in \mathbf{P}^*$ is a hyperplane which intersects Q , then either H is tangent to Q (in which case $\theta(H)$ is the point of tangency) or there exists a cone tangent to Q meeting Q in $Q \cap H$ — the vertex of this cone will be $\theta(H)$. If $x \in \Omega$, then there will be no tangents to Q containing x , but by representing x as an intersection $H_1 \cap \cdots \cap H_n$, we obtain $\theta(x)$ as the hyperplane containing $\theta(H_1), \dots, \theta(H_n)$.

EXERCISE 3.2.11. Show that $\mathbf{P} \xrightarrow{\theta} \mathbf{P}^*$ is projective.

Observe that a polarity on \mathbf{P} of signature (p, q) determines, for each non-conjugate point $x \in \mathbf{P}$ a unique reflection R_x which preserves the polarity. The group of collineations preserving such a polarity is the *projective orthogonal group* $\mathrm{PO}(p, q)$, that is, the image of the orthogonal group $\mathrm{O}(p, q) \subset \mathrm{GL}(n+1, \mathbb{R})$ under the projectivization homomorphism

$$\mathrm{GL}(n+1, \mathbb{R}) \longrightarrow \mathrm{PGL}(n+1, \mathbb{R})$$

having kernel the scalar matrices $\mathbb{R}^\times \subset \mathrm{GL}(n+1, \mathbb{R})$. Let

$$\Omega = \{\Pi(v) \in \mathbf{P} \mid \mathbf{B}(v, v) < 0\};$$

then by projection from the origin Ω can be identified with the hyperquadric

$$\{v \in \mathbb{R}^{p,q} \mid \mathbf{B}(v, v) = -1\}$$

whose induced pseudo-Riemannian metric has signature $(q, p - 1)$ and constant nonzero curvature. In particular if $(p, q) = (1, n)$ then Ω is a model for hyperbolic n -space \mathbf{H}^n in the sense that the group of isometries of \mathbf{H}^n are represented precisely as the group of collineations of \mathbf{P}^n preserving Ω^n . In this model, geodesics are the intersections of projective lines in \mathbf{P} with Ω ; more generally intersections of projective subspaces with Ω define totally geodesic subspaces.

Consider the case that $\mathbf{P} = \mathbf{P}^2$. Points “outside” Ω correspond to geodesics in \mathbf{H}^2 . If $p_1, p_2 \in \Omega^\dagger$, then

$$\overleftarrow{p_1 p_2} \cap \Omega \neq \emptyset$$

if and only if the geodesics $\theta(p_1), \theta(p_2)$ are ultra-parallel in \mathbf{H}^2 ; in this case $\theta(\overleftarrow{p_1 p_2})$ is the geodesic orthogonal to both $\theta(p_1), \theta(p_2)$. (Geodesics $\theta(p)$ and l are orthogonal if and only if $p \in l$.) Furthermore $\overleftarrow{p_1 p_2}$ is tangent to Q if and only if $\theta(p_1)$ and $\theta(p_2)$ are parallel. For more information on this model for hyperbolic geometry, see [80]. or [265], §2. This model for non-Euclidean geometry seems to have first been discovered by Cayley in 1858.

3.3. Projective model of hyperbolic geometry

The case when $q = 1$ is fundamentally important. Then $\mathbf{P}(\mathbf{Q}^-)$ is equivalent to the *unit ball* $\mathbb{B} \subset \mathbb{R}^n$ defined by

$$\|x\|^2 = x \cdot x = \sum_{i=1}^n (x^i)^2 < 1.$$

and the induced Riemannian metric

$$\begin{aligned} ds_{\mathbb{B}}^2 &= \frac{-4}{\sqrt{1 - \|x\|^2}} d^2 \sqrt{1 - \|x\|^2} \\ &= \frac{-4}{1 - \|x\|^2} \left\{ (1 - \|x\|^2) dx \cdot dx + (x \cdot dx)^2 \right\} \\ (10) \quad &= \frac{4}{(1 - \|x\|^2)^2} \sum_{i=1}^n (x^i dx^i)^2 + (1 - \|x\|^2)^2 (dx^i)^2 \end{aligned}$$

defines a complete $\mathbf{PO}(n, 1)$ -invariant Riemannian structure of constant curvature -1 on \mathbb{B} . The resulting Riemannian manifold is (*real*) *hyperbolic space* \mathbf{H}^n .

In 1894 Hilbert discovered a beautiful general construction of the distance function on the underlying metric space involving projective geometry. Suppose $x, y \in \mathbb{B} \subset \mathbf{P}$ are distinct points. They span a unique projective line $\overleftarrow{xy} \subset \mathbf{P}$. Then \overleftarrow{xy} meets $\partial\mathbb{B}$ in two points




FIGURE 3.1. The *Klein-Beltrami* projective model of hyperbolic space. The Hilbert metric on the convex domain bounded by a quadric in projective space is defined in terms of cross-ratio. This metric is a Riemannian metric of constant negative curvature.

x_0, y_0 as in Figure 3.1. Then the cross-ratio $[x, y, x_0, y_0]$ is defined and the *Hilbert distance*

$$d(x, y) := \log[x, y, x_0, y_0]$$

makes (\mathbb{B}, d) into a metric space.

EXERCISE 3.3.1. *Show that this metric space underlies the Riemannian structure defined above. Show that its group of isometries is $\mathrm{PO}(n, 1)$ and acts transitively not just on \mathbb{H}^n but on its unit tangent bundle.*

This metric is analogous to the construction of the Fubini-Study metric given in Exercise 3.2.6. Its generalizes to arbitrary properly convex domains, as will be discussed in the next chapter.

3.3.1. The hyperbolic plane. Due to its fundamental role, we discuss the projective model of the hyperbolic plane \mathbb{H}^2 in detail:

EXERCISE 3.3.2. *Let \mathbb{H}^2 denote the upper halfplane $\mathbb{R} \times \mathbb{R}^+$ with the Poincaré metric*

$$g = y^{-2}(dx^2 + dy^2).$$

Our model for the Lorentzian vector space $\mathbb{R}^{2,1}$ is the Lie algebra $\mathfrak{sl}(2, \mathbb{R})$ of traceless 2×2 real matrices with the quadratic form

$$\frac{1}{2} \mathrm{Tr} \begin{bmatrix} a & b \\ c & -a \end{bmatrix}^2 = a^2 + bc = a^2 + \left(\frac{b+c}{2}\right)^2 - \left(\frac{b-c}{2}\right)^2.$$

The mapping

$$(11) \quad \mathbb{H}^2 \xrightarrow{\mathcal{J}} \mathfrak{sl}(2, \mathbb{R}) \cap \mathrm{SL}(2, \mathbb{R}) \subset \mathbb{R}^{2,1}$$

$$(x, y) \longmapsto y^{-1} \begin{bmatrix} x & -(x^2 + y^2) \\ 1 & -x \end{bmatrix}$$

isometrically embeds \mathbb{H}^2 as the component of the hypersphere in $\mathbb{R}^{2,1}$

$$\left\{ \begin{bmatrix} a & b \\ c & -a \end{bmatrix} \mid a^2 + bc = -1, \quad c > 0 \right\}.$$

Composition $\mathsf{P} \circ \mathcal{J}$ with projectivization isometrically embeds \mathbb{H}^2 in projective space, as above. $\mathsf{P} \circ \mathcal{J}$ is equivariant, mapping the isometry group $\mathrm{PGL}(2, \mathbb{R})$ of \mathbb{H}^2 isomorphically onto the projective automorphism group $\mathrm{PO}(2, 1) \cong \mathrm{SO}(2, 1)$. For any point $z \in \mathbb{H}^2$, the matrix $\mathcal{J}(z)$ corresponds to the symmetry about z , that is, the orientation-preserving involutive isometry of \mathbb{H}^2 fixing z .

Appendix B.4 derives the Levi-Civita connection of this metric.

3.3.2. The upper halfspace model of hyperbolic 3-space.

Hyperbolic 3-space is fundamentally important as well. Its group of orientation-preserving isometries is isomorphic to $\mathrm{PGL}(2, \mathbb{C}) \cong \mathrm{PSL}(2, \mathbb{C})$ under a local isomorphism

$$\mathrm{PSL}(2, \mathbb{C}) \longrightarrow \mathrm{O}(3, 1).$$

(See Appendix E for a construction of such a local isomorphism.) It admits a useful model as the *upper halfspace* in the division algebra of *quaternions*.

3.3.2.1. *Quaternions*. Recall that the (Hamilton) quaternions are defined as a 4-dimensional real vector space \mathbb{H} with basis denoted $\{\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$. Its multiplicative structure is defined by the multiplication table:

	$\mathbf{1}$	\mathbf{i}	\mathbf{j}	\mathbf{k}
$\mathbf{1}$	$\mathbf{1}$	\mathbf{i}	\mathbf{j}	\mathbf{k}
\mathbf{i}	\mathbf{i}	$-\mathbf{1}$	\mathbf{k}	$-\mathbf{j}$
\mathbf{j}	\mathbf{j}	$-\mathbf{k}$	$-\mathbf{1}$	\mathbf{i}
\mathbf{k}	\mathbf{k}	\mathbf{j}	$-\mathbf{i}$	$-\mathbf{1}$

$\mathbf{1}$ is a two-sided identity element, and \mathbb{H} is an associative (but *not* commutative!) division algebra over \mathbb{R} . We write

$$q := r\mathbf{1} + x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = r\mathbf{1} + \mathbf{v}$$

where $r, x, y, z \in \mathbb{R}$ are scalars and

$$\mathbf{v} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k} \in \mathbb{R}^3$$

is a vector.

Quaternionic conjugation

$$\begin{aligned}\mathbb{H} &\longrightarrow \mathbb{H} \\ q = r\mathbf{1} + \mathbf{v} &\longmapsto \bar{q} := r\mathbf{1} - \mathbf{v}\end{aligned}$$

is an *anti-automorphism*, (that is, $\overline{q_1 q_2} = \overline{q_2} \overline{q_1}$). Its fixed set is thus a subalgebra, the image of the embedding

$$\begin{aligned}\mathbb{R} &\hookrightarrow \mathbb{H} \\ r\mathbf{1} &\longmapsto r\mathbf{1} + 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}.\end{aligned}$$

(In fact, $\mathbb{R} = \mathbb{R}\mathbf{1}$ is the *center* of \mathbb{H} .) In particular the *real part* is the projection

$$\begin{aligned}\mathbb{H} &\xrightarrow{\Re} \mathbb{R} \\ q &\longmapsto r = \frac{1}{2}(q + \bar{q})\end{aligned}$$

A quaternion is *pure* if its real part is zero, and pure quaternions identify with the vector space \mathbb{R}^3 . Multiplication of pure quaternions correspond to the operations of dot and cross product of vectors in \mathbb{R}^3 :

$$\mathbf{v}_1 \mathbf{v}_2 = -(\mathbf{v}_1 \cdot \mathbf{v}_2) \mathbf{1} + \mathbf{v}_1 \times \mathbf{v}_2.$$

Furthermore

$$q\bar{q} = \|q\|^2 = r^2 + \|\mathbf{v}\|^2 \geq 0$$

and $q\bar{q} > 0$ if $q \neq 0$. Thus \mathbb{H} is a *division algebra*, with

$$q^{-1} := \|q\|^{-2} \bar{q}.$$

3.3.2.2. *Hyperbolic 3-space*. The embedding

$$\begin{aligned}\mathbb{C} &\hookrightarrow \mathbb{H} \\ r\mathbf{1} + x\mathbf{i} &\longmapsto r\mathbf{1} + x\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}\end{aligned}$$

makes \mathbb{H} into a (left) vector space over \mathbb{C} . Define the *upper halfspace* \mathbb{H}^3 as the subset of \mathbb{H} consisting of $z + h\mathbf{j}$, where $z \in \mathbb{C}$ and $h > 0$.

EXERCISE 3.3.3. *Let*

$$g := \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}(2, \mathbb{C}),$$

that is, $a, b, c, d \in \mathbb{C}$ and $ad - bc = 1$.

- Show that

$$g(w) := (aw + b)(cw + d)^{-1}$$

defines a (left) action of $\mathrm{SL}(2, \mathbb{C})$ on \mathbb{H} . Interpret this action as a projective action over \mathbb{H} .

- Show that \mathbf{H}^3 is invariant under this action.
- Find a Riemannian metric on \mathbf{H}^3 upon which this group acts as its group of orientation-preserving isometries.
- Prove that the subspace $\mathbf{H}^2 \subset \mathbf{H}^3$ defined by $\Im(z) = 0$ (that is, $\mathbb{R} \times \mathbb{R}^+$) is an isometrically embedded hyperbolic plane. Determine its group of isometries.
- Show that

$$z + h\mathbf{j} \mapsto \bar{z} + h\mathbf{j}$$

is an isometry of \mathbf{H}^3 fixing \mathbf{H}^2 (reflection in \mathbf{H}^2) and given by quaternion conjugation by

$$q \mapsto -\mathbf{i}\bar{q}\mathbf{i}.$$

- Show that the symmetry in the point $\mathbf{j} \in \mathbf{H}^3$ is given by the quaternionic formula

$$q \mapsto \mathbf{i}\bar{q}^{-1}\mathbf{i}.$$




FIGURE 3.2. Projective model of a $(3, 3, 4)$ -triangle tessellation of H^2




FIGURE 3.3. Projective deformation of hyperbolic $(3, 3, 4)$ -triangle tessellation

CHAPTER 4

Convexity

The Klein-Beltrami projective model of hyperbolic geometry extends to geometries defined on properly convex domains in \mathbb{RP}^n . This chapter concerns this notion of convexity. After surveying some basic examples, we describe the *Hilbert metric*, a projective-geometry construction which includes the Klein-Beltrami hyperbolic metric. This metric is only Riemannian only in this special case, and is generally *Finsler*, that is, arises from a norm on the tangent spaces. From there we describe another construction, due to Vinberg [278], which leads to natural Riemannian structure. We use this structure to give new proofs of several results of Benzécri [35], on spaces of convex bodies in projective space. These results lead to regularity properties for domains arising from convex \mathbb{RP}^n -manifolds.

A convex domain $\Omega \subset \mathbb{P}$ is said to be *quasi-homogeneous* if $\text{Aut}(\Omega)$ acts syndetically — that is, the quotient $\Omega/\text{Aut}(\Omega)$ is compact, but not necessarily Hausdorff. (Benzécri calls such domains *balayable*, translated as “sweepable”. If the action is *proper*, (that is, the quotient is Hausdorff), then the domain is said to be *divisible*. We are particularly interested in these compactness criteria. For reasons of space, we do not discuss the currently very active field of *finite volume* properly convex manifolds, but refer to relatively recent work by Marquis [212, 210, 211], Cooper-Long-Tillmann [78, 79] Ballas-Danciger-Lee-Marquis [18], Ballas-Cooper-Leitner [17], and Choi [74].

Later in §12.2 we revisit these ideas in the more general setting of the *intrinsic metrics* introduced by Carathéodory and Kobayashi, in the analogous context of complex manifolds. Our treatment of the projective theory closely follows Kobayashi [177, 178, 179] and Vey [275].

4.1. Convex domains and cones

Let W be a real vector space. Recall that a subset $\Omega \subset W$ is *convex* if, whenever $x, y \in \Omega$, the line segment $\overline{xy} \subset \Omega$. Equivalently, W is closed under *convex combinations*: if $\mathbf{w}_1, \dots, \mathbf{w}_m \in W$ and $t_1, \dots, t_m \in \mathbb{R}$ satisfy $t_i \geq 0$ and

$$t_1 + \dots + t_m = 1,$$

then

$$t_1 \mathbf{w}_1 + \cdots + t_m \mathbf{w}_m \in \Omega.$$

A domain $\Omega \subset \mathbf{W}$ is a *cone* if and only if it is invariant under the group \mathbb{R}^+ of positive homotheties, that is, scalar multiplication by positive real numbers. A cone is *sharp* (in French, *saillant*) if it contains no complete affine line.

Since $\Omega' \subset \mathbf{W} \setminus \{\mathbf{0}\}$ is convex, Ω must be disjoint from at least one hyperplane H in \mathbf{P} . (In particular \mathbf{P} is itself *not* convex.) Equivalently $\Omega \subset \mathbf{P}$ is convex if a hyperplane $H \subset \mathbf{P}$ exists such that Ω is a convex set in the complementary affine space $\mathbf{P} \setminus H$.

A domain $\Omega \subset \mathbf{P}$ is *properly convex* if and only if a *sharp* properly convex cone $\Omega' \subset \mathbf{W}$ exists such that $\Omega = \Pi(\Omega')$. Equivalently, Ω is properly convex if and only if a hyperplane $H \subset \mathbf{P}$ exists such that $\overline{\Omega}$ is a convex subset of the affine space $\mathbf{P} \setminus H$. If $\Omega \subset \mathbf{P}$ is properly convex, then its intersection $\Omega \cap \mathbf{P}'$ with any projective subspace $\mathbf{P}' \subset \mathbf{P}$ is either empty or properly convex in \mathbf{P}' . In particular every projective line intersecting Ω meets $\partial\Omega$ in exactly two points.

For example, \mathbf{W} itself and the upper half-space

$$\mathbb{R}^n \times \mathbb{R}^+ = \{(x^0, \dots, x^n) \in \mathbf{W} \mid x^0 > 0\}$$

are both convex cones but neither is sharp. The *positive orthant*

$$(\mathbb{R}^+)^{n+1} := \{(x^0, \dots, x^n) \in \mathbf{W} \mid x^i > 0 \text{ for } i = 0, 1, \dots, n\}$$

and the positive light-cone

$$C_{n+1} = \{(x^0, \dots, x^n) \in \mathbf{W} \mid x^0 > 0 \text{ and } -(x^0)^2 + (x^1)^2 + \dots + (x^n)^2 < 0\}$$

are both properly convex cones.

EXERCISE 4.1.1. Define the parabolic convex domain as

$$(12) \quad \mathcal{P} := \{(x, y) \in \mathbb{R}^2 \mid y > x^2\}.$$

- Show that \mathcal{P} is a properly convex affine domain but not affinely equivalent to a cone.
- Describe the group of affine automorphisms of \mathcal{P} .
- Describe the group of projective automorphisms of \mathcal{P} .

EXERCISE 4.1.2. Show that the set $\mathfrak{P}_n(\mathbb{R})$ of all positive definite symmetric $n \times n$ real matrices is a properly convex cone in the $n(n+1)/2$ -dimensional vector space \mathbf{W} of $n \times n$ symmetric matrices. Are there any affine transformations of \mathbf{W} preserving $\mathfrak{P}_n(\mathbb{R})$? What is its group of affine automorphisms?

Convex affine domains have the structure of principal \mathbb{R} -bundles over sharp convex cones:

PROPOSITION 4.1.3. *Let $\Omega \subset V$ be an open convex cone in a vector space. Then there exists a unique linear subspace $W \subset V$ such that:*

- Ω is invariant under translation by vectors in W (that is, Ω is W -invariant;)
- There exists a properly convex cone $\Omega_0 \subset V/W$ such that $\Omega = \pi_W^{-1}(\Omega_0)$ where $\pi_W : V \rightarrow V/W$ denotes linear projection with kernel W .

PROOF. Let

$$W = \{w \in V \mid x + tw \in \Omega, \forall x \in \Omega, t \in \mathbb{R}\}.$$

Then W is a linear subspace of V and Ω is W -invariant. Let

$$\Omega_0 = \pi_W(\Omega) \subset V/W;$$

then $\Omega = \pi_W^{-1}(\Omega_0)$. We must show that Ω_0 is properly convex. To this end we can immediately reduce to the case $W = 0$. Suppose that Ω contains a complete affine line $\{y + tw \mid t \in \mathbb{R}\}$ where $y \in \Omega$ and $w \in V$. Then for each $s, t \in \mathbb{R}$

$$x_{s,t} = \frac{s}{s+1}x + \frac{1}{s+1}\left(y + stw\right) \in \Omega$$

whence

$$\lim_{s \rightarrow \infty} x_{s,t} = x + tw \in \bar{\Omega}.$$

Thus $x + tw \in \bar{\Omega}$ for all $t \in \mathbb{R}$. Since $x \in \Omega$ and Ω is open and convex, $x + tw \in \Omega$ for all $t \in \mathbb{R}$ and $w \in W$ as claimed. \square

4.1.1. Halfspaces and supporting hyperplanes. Let $\Omega \subset A$ be a proper convex domain in an affine space A . In general Ω will be an intersection of open halfspaces.

Here is a sketch of the proof, using the Hahn-Banach theorem.

The Hahn-Banach theorem (as stated in Theorem 11.4.1 of Berger [37]) asserts that every affine subspace (for example a point) disjoint from Ω extends to an affine hyperplane disjoint from Ω . Since Ω is proper, $A \setminus \Omega$ is nonempty. Choose a point $p \in A \setminus \Omega$. The Hahn-Banach theorem guarantees an affine hyperplane $H \subset V$ containing p disjoint from Ω . The two components of its complement $V \setminus H$ are halfspaces. Since Ω is connected, one of them contains Ω .

Let \mathcal{W} denote the set of halfspaces $W \supset \Omega$. Suppose that

$$p \in \left(\bigcap_{W \in \mathcal{W}} W \right) \setminus \Omega.$$

Applying the Hahn-Banach theorem again guarantees a halfspace containing Ω but not containing p , a contradiction. Thus Ω is the intersection of open halfspaces.

The boundary of a minimal open halfspace containing Ω is a hyperplane, called a *supporting hyperplane* for Ω . By Proposition 11.5.2 of Berger [37], at every point of $\partial\Omega$ is a supporting hyperplane for Ω .

4.1.2. Convexity in projective space. Convexity is somewhat more subtle in projective space \mathbb{P} . First observe first that convexity is invariant under translations, and thus invariant under affine transformations. Say that a domain $\Omega \subset \mathbb{P}$ is *convex* if and only if Ω lies in some affine patch $A \subset \mathbb{P}$, and is a convex subset of A . The above remarks imply that this notion is independent of the choice of A . Equivalently, if $\mathbb{P} = \mathbb{P}(W)$ and

$$W \setminus \{0\} \xrightarrow{\Pi} \mathbb{P}$$

denotes projectivization, then Ω is convex if $\Omega = \Pi(\Omega')$ for some convex cone $\Omega' \subset W$.

For studying convex subsets of projective space, passing to the double covering space, the *sphere of directions* is useful, especially for calculations.

4.2. The Hilbert metric

In 1894 Hilbert introduced a projectively invariant metric $d = d_\Omega$ on any properly convex domain $\Omega \subset \mathbb{P}$ as follows. This was introduced in §3.3 as an explicit form of the metric on \mathbb{H}^n in the Klein-Beltrami model. After reviewing its definition and basic properties, we discuss the other basic example of an open simplex, in which case the metric does not arise from a Riemannian structure. A simple example is Vey's semisimplicity Theorem 4.3.1, which we prove in a special case (used later in §12 to classify completely incomplete affine structures, following Kobayashi [179] and Vey [277, 276]).

4.2.1. Definition and basic properties. Let $x, y \in \Omega$ be a pair of distinct points; then the line \overleftrightarrow{xy} meets $\partial\Omega$ in two points which we denote by x_∞, y_∞ (the point closest to x will be x_∞ , etc). The *Hilbert distance*

$$d = d_\Omega^{\text{Hilb}}$$

between x and y in Ω will be defined as the logarithm of the cross-ratio of this quadruple:

$$d(x, y) = \log[x_\infty, x, y, y_\infty]$$

(where the cross-ratio is defined in (8)). Clearly $d(x, y) \geq 0$, so that $d(x, y) = d(y, x)$. Since Ω contains no complete affine line, $x_\infty \neq y_\infty$, and $d(x, y) > 0$ if $x \neq y$.

Similarly

$$\Omega \times \Omega \xrightarrow{d} \mathbb{R}$$

is *proper*, or *finitely compact*: that is, for each $x \in \Omega$ and $r > 0$, the closed “ r -ball”

$$B_r(x) = \{y \in \Omega \mid d(x, y) \leq r\}$$

is compact. Once the triangle inequality is established, the completeness of the metric space (Ω, d) follows. The triangle inequality results from the convexity of Ω , although we deduce it by showing that the Hilbert metric agrees with the general intrinsic metric introduced by Kobayashi [179]. Thus we enforce the triangle inequality as part of the construction of the metric. In this metric the geodesics are represented by straight lines.

By Exercise 3.3.1, the Hilbert metric on a quadric domain agrees with the Beltrami-Klein Riemannian structure. The other fundamental example is the open simplex, which is the projectivization of an *orthant* in \mathbb{R}^n (for example the *positive orthant* $(\mathbb{R}^+)^n \subset \mathbb{R}^n$). When $n = 3$, the projective domain is just a triangle.

4.2.2. The Hilbert metric on a triangle. Let $\Delta \subset \mathbb{P}^2$ denote a domain bounded by a triangle. Then the balls in the Hilbert metric are hexagonal regions. (In general if Ω is a convex k -gon in \mathbb{P}^2 then the unit balls in the Hilbert metric will be interiors of $2k$ -gons.)

EXERCISE 4.2.1. (*Unit balls in the Hilbert metric*)

- Prove that $\text{Aut}(\Delta)$ is conjugate to the group of diagonal matrices with positive eigenvalues.
- Deduce that $\text{Aut}(\Delta)$ acts transitively on Δ .
- Conclude that all the unit balls are isometric.

Here is a construction which illustrates the Hilbert geometry of Δ . (Compare Figure 2.3.) Start with a triangle Δ and choose line segments l_1, l_2, l_3 from an arbitrary point $p_1 \in \Delta$ to the vertices v_1, v_2, v_3 of Δ . Choose another point p_2 on l_1 , say, and form lines l_4, l_5 joining it to the remaining vertices. Let

$$\rho = \log \left| \left[v_1, p_1, p_2, l_1 \cap \overleftarrow{v_2 v_3} \right] \right|$$

where $[\cdot]$ denotes the cross-ratio of four points on l_1 . The lines l_4, l_5 intersect l_2, l_3 in two new points which we call p_3, p_4 . Join these two points to the vertices by new lines l_i which intersect the old l_i in new

points p_i . In this way one generates infinitely many lines and points inside Δ , forming a configuration of smaller triangles T_j inside Δ . For each p_i , the union of the T_j with vertex p_i is a convex hexagon which is a Hilbert ball in Δ of radius ρ . Note that this configuration is combinatorially equivalent to the tessellation of the plane by congruent equilateral triangles. Indeed, this tessellation of Δ arises from an action of a (3,3,3)-triangle group by collineations and converges (in an appropriate sense) to the Euclidean equilateral-triangle tessellation as $\rho \rightarrow 0$.

EXERCISE 4.2.2. Let $\Delta := \{(x, y) \in \mathbb{R}^2 \mid x, y > 0\}$ be the positive quadrant. Then the Hilbert distance is given by

$$d((x, y), (x', y')) = \log \max \left\{ \frac{x}{x'}, \frac{x'}{x}, \frac{y}{y'}, \frac{y'}{y}, \frac{xy'}{x'y}, \frac{x'y}{xy'} \right\}.$$

- Show that the unit balls are hexagons.
- For any two points $p, p' \in \Delta$, show that there are infinitely many geodesics joining p to p' .
- Show that there are even non-smooth polygonal curves from p to p' having minimal length.

Daryl Cooper has called such a Finsler, where the unit balls are hexagons, a *hex-metric*.

4.3. Vey's semisimplicity theorem

The following theorem is due to Vey [277, 276].

THEOREM 4.3.1. Let V be a real vector space and $\Omega \subset V$ a divisible sharp convex cone. Then the action of $\text{Aut}(\Omega)$ is semisimple, that is, any $\text{Aut}(\Omega)$ -invariant linear subspace $W < V$, there exists an $\text{Aut}(\Omega)$ -invariant complementary linear subspace. In particular a unique decomposition

$$V = \bigoplus_{i=1}^r V_i$$

exists, with sharp convex cones $\Omega_i \subset V_i$ such that

$$\Omega = \prod_{i=1}^r \Omega_i$$

and the action of $\text{Aut}(\Omega_i)$ on V_i is irreducible.

COROLLARY 4.3.2. Suppose $\Omega \subset A^n$ is a properly convex divisible domain. Then Ω is a sharp convex cone.

The parabolic region $\{(x, y) \in \mathbb{A}^2 \mid y - x^2 > 0\}$ is an example of a properly convex domain with a group $\Gamma < \text{Aut}(\Omega)$ such that $\Gamma \backslash \Omega$ is compact but not Hausdorff (that is, the transformation group (Γ, Ω) is *syndetic*). In particular Ω is *not* a cone.

We don't prove all of Theorem 4.3.1 here, but just the special case when W is assumed to be a supporting hyperplane for Ω . This is all which is needed to deduce Corollary 4.3.2. Our treatment is based on Hoban [152].

PROOF OF COROLLARY 4.3.2. Suppose that $\Omega \subset \mathbb{A}$ is a properly convex domain with $\Gamma < \text{Aut}(\Omega)$ dividing Ω . Embed \mathbb{A} as an affine hyperplane in a vector space \mathbb{V} . Let $\psi \in \mathbb{V}^*$ be a linear functional such that $\mathbb{A} = \psi^{-1}(1)$. Then

$$\Omega' := \{\omega \in \mathbb{V} \mid \psi(\omega) \neq 0 \text{ and } \psi(\omega)^{-1}\omega \in \Omega\}$$

is sharp convex cone to which the action Γ on \mathbb{A} extends to a linear action on \mathbb{V} preserving Ω' . This linear action extends to $\Gamma \times \mathbb{R}^+$, where \mathbb{R}^+ acts by homotheties. The linear hyperplane

$$W := \text{Ker}(\psi) = \psi^{-1}(0)$$

supports Ω' in the sense of §4.1.1. Furthermore, taking $\lambda > 1$, the product $\Gamma' := \Gamma \times \langle \lambda \rangle$ divides Ω' and preserves W .

By the special case of Theorem 4.3.1 when W is a supporting hyperplane, $\exists L < \mathbb{V}$ which is Γ -invariant and

$$\mathbb{V} = W \oplus L.$$

Now L is a line which intersects the affine hyperplane \mathbb{A} in a point p_0 , and p_0 is fixed by Γ . Then Ω is a open convex cone with vertex p_0 . \square

PROOF OF SPECIAL CASE OF THEOREM 4.3.1. The proof uses the following general fact about metric spaces, whose proof is given in Appendix C.

LEMMA 4.3.3. *Let X be any compact metric space. Then any distance non-increasing homeomorphism of X is an isometry*

Let $\Omega^* \subset \mathbb{V}^*$ be the cone dual to Ω , as in §4.4. If W is a linear hyperplane which supports Ω , then the annihilator $\text{Ann}(W) < \mathbb{V}^*$ is a line lying in $\partial\Omega^*$. It suffices to find a Γ -invariant complementary linear hyperplane $H < \mathbb{V}^*$.

Let L be a line which is invariant under Γ and intersects nontrivially with $\overline{\Omega}$.

We define a Γ -invariant map $\Omega \xrightarrow{s} \partial\Omega$ as follows. For each $x \in \Omega$, let $\Omega_x := \Omega \cap (x + L)$. Since Ω is sharp, Ω_x is a ray, with endpoint $s(x)$ on the boundary of Ω .

Evidently s commutes with Γ .

Then, for $t \in \mathbb{R}$,

$$c_t(x) := s(x) + e^t(x - s(x))$$

defines a one parameter group of homeomorphisms of Ω .




FIGURE 4.1. Projection does not increase Hilbert distance

□

LEMMA 4.3.4. c_t does not increase Hilbert distance:

$$(13) \quad d^{\text{Hilb}}(c_t(x), c_t(y)) \leq d^{\text{Hilb}}(x, y)$$

PROOF. To prove (13), choose $p, q \in \Omega$ as in Figure 4.1. Clearly $c_t(p) - p$ and $c_t(q) - q$ are both in L . Let a and b denote the intersections of \overleftrightarrow{pq} with $\partial\Omega$. Let c and d denote the two points of the intersection of $\partial\Omega$ with

$$\overleftrightarrow{c_t(p) c_t(q)}.$$


Now the intersections of $a + L$ and $b + L$ respectively with $\overleftrightarrow{c_t(p) c_t(q)}$ are two points, denoted \widehat{a} and \widehat{b} . Since Ω is convex,

$$\overline{\widehat{a}\widehat{b}} \subset \overline{cd},$$

hence

$$[c, c_t(p), c_t(q), d] \leq [\widehat{a}, c_t(p), c_t(q), \widehat{b}] = [a, p, q, b].$$

The last equality is due to the invariance of the cross ratio under projective transformations (in particular this is a perspectivity). Since cross ratio is non-increasing, d^{Hilb} is non-increasing as well, concluding the proof of Lemma 4.3.4. □

FIGURE 4.2. c_t is an isometry of the cone

Since s commutes with Γ , we can pass the map c_t down to a map $M \xrightarrow{C_t} M$ on the quotient $M = \Gamma \backslash \Omega$. Furthermore C_t is distance non-increasing with respect to d_M^{Hilb} . Note that M is a metric space because Γ divides Ω . (In particular M is Hausdorff.) Thus C_t is a distance non-increasing homeomorphism of $M = \Gamma \backslash \Omega$, and by Lemma 4.3.3, it is an isometry. Thus c_t is a local isometry on Ω .

Note: Figure 4.1 is a bit misleading here since the domain pictured is not a cone, and c_t is actually strictly decreasing for this domain. Figure 4.2 below, depicts how c_t acts isometrically on the cone.

LEMMA 4.3.5. $s(\Omega)$ is convex.

PROOF. Suppose $\exists x, y \in \Omega$ such that the line segment

$$\overline{s(x) s(y)} \not\subset s(\Omega) \subset \partial\Omega.$$

Suppose

$$p \in \overline{s(x) s(y)} \setminus s(\Omega).$$

Let N be a neighborhood of p on which c_t is an isometry, and choose

$$q \in \overline{s(x) s(y)} \cap N.$$

Choose $t < 0$ such that both $c_t(p)$ and $c_t(q)$ lie in N . Let a, b be the intersections of $\overleftarrow{c_t(p) c_t(q)}$ with $\partial\Omega$. Write

$$\hat{a} = (a + L) \cap \overleftarrow{p q}, \quad \hat{b} = (b + L) \cap \overleftarrow{p q}.$$

Clearly $\hat{a} \in \overline{s(x) s(y)}$, but not equal to $s(x)$: otherwise $a = s(x)$ and $t = 0$. Similarly $\hat{b} \in \overline{s(x) s(y)}$, but not equal to $s(y)$.

PROOF. Clearly Ω^* is a convex cone. We show that Ω^* is properly convex and open. If Ω^* contains a line, then $\psi_0, \lambda \in \mathbf{V}^*$ exist such that $\lambda \neq 0$ and $\psi_0 + t\lambda \in \Omega^*$ for all $t \in \mathbb{R}$, that is, $\forall x \in \Omega$,

$$\psi_0(x) + t\lambda(x) > 0$$

for each $t \in \mathbb{R}$. Let $x \in \Omega$; then necessarily $\lambda(x) = 0$. Otherwise, if $\lambda(x) \neq 0$, then $t \in \mathbb{R}$ exists with

$$\psi_0(x) + t\lambda(x) \leq 0,$$

a contradiction. Thus Ω^* is properly convex. The openness of Ω^* follows from the proper convexity of Ω . Since Ω is properly convex, its projectivization $\mathbf{P}(\Omega)$ is a properly convex domain; in particular its closure lies in an open ball in an affine subspace \mathbf{A} of \mathbf{P} and thus the set of hyperplanes in \mathbf{P} disjoint from $\mathbf{P}(\Omega)$ is open. It follows that $\mathbf{P}(\Omega^*)$, and hence Ω^* , is open. \square

LEMMA 4.4.2. *The canonical isomorphism $\mathbf{V} \longrightarrow \mathbf{V}^{**}$ maps Ω onto Ω^{**} .*

PROOF. Identify \mathbf{V}^{**} with \mathbf{V} . Clearly, $\Omega \subset \Omega^{**}$. Since both Ω and Ω^{**} are open convex cones, either $\Omega = \Omega^{**}$ or $\partial\Omega \cap \Omega^{**} \neq \emptyset$. Let $y \in \partial\Omega \cap \Omega^{**}$. Let $H \subset \mathbf{V}$ be a supporting hyperplane for Ω at y . Then the covector $\psi \in \mathbf{V}^*$ defining H vanishes at y and is positive on Ω . Thus $\psi \in \Omega^*$. However, $y \in \Omega^{**}$ implies $\psi(y) > 0$, a contradiction. \square

THEOREM 4.4.3. *Let $\Omega \subset \mathbf{V}$ be a properly convex cone. Then there exists a real analytic $\text{Aff}(\Omega)$ -invariant closed 1-form α on Ω such that its covariant differential $\nabla\alpha$ is an $\text{Aff}(\Omega)$ -invariant Riemannian metric on Ω . Furthermore*

$$\alpha(\mathbf{R}_V) = -n < 0$$

where \mathbf{R}_V is the radiant vector field on \mathbf{V} .

4.4.1. The Vinberg characteristic function. Let $d\psi$ denote a parallel volume form on \mathbf{V}^* . Define the *characteristic function* f of the properly convex cone Ω by the following integral

$$(14) \quad \Omega \xrightarrow{f} \mathbb{R}$$

$$f(x) := \int_{\Omega^*} e^{-\psi(x)} d\psi$$

over the dual cone Ω^* . This function and its derivatives yields a canonical Riemannian geometry on Ω invariant under the automorphism group $\text{Aff}(\Omega)$. Furthermore it produces a canonical diffeomorphism $\Omega \longrightarrow \Omega^*$. (Note that replacing the parallel volume form $d\psi$ by another one $c d\psi$ replaces the characteristic function f by its constant

multiple $c f$. Thus $\Omega \xrightarrow{f} \mathbb{R}$ is well-defined only up to scaling.) For example in the one-dimensional case, where

$$\Omega = \mathbb{R}_+ \subset \mathbf{V} = \mathbb{R}$$

the dual cone equals $\Omega^* = \mathbb{R}_+$ and the characteristic function equals

$$f(x) = \int_0^\infty e^{-\psi x} d\psi = \frac{1}{x}.$$

We begin by showing the integral (14) converges for $x \in \Omega$. For $x \in \mathbf{V}$ and $t \in \mathbb{R}$ consider the hyperplane cross-section

$$\mathbf{V}_x^*(t) = \{\psi \in \mathbf{V}^* \mid \psi(x) = t\}$$

and let

$$\Omega_x^*(t) = \Omega^* \cap \mathbf{V}_x^*(t).$$

For each $x \in \Omega$ we obtain a decomposition

$$\Omega^* = \bigcup_{t>0} \Omega_x^*(t)$$

and for each $s > 0$ there is a diffeomorphism

$$\begin{aligned} \Omega_x^*(t) &\xrightarrow{h_s} \Omega_x^*(st) \\ \psi &\longmapsto s\psi \end{aligned}$$

and obviously $h_s \circ h_t = h_{st}$. We decompose the volume form $d\psi$ on Ω^* as

$$d\psi = d\psi_t \wedge dt$$

where $d\psi_t$ is an $(n-1)$ -form on $\mathbf{V}_x^*(t)$. Now the volume form $(h_s)^* d\psi_{st}$ on $\Omega_x^*(t)$ is a parallel translate of $t^{n-1} d\psi_t$. Thus:

$$\begin{aligned} f(x) &= \int_0^\infty \left(e^{-t} \int_{\Omega_x^*(t)} d\psi_t \right) dt \\ &= \int_0^\infty e^{-t} t^{n-1} \left(\int_{\Omega_x^*(1)} d\psi_1 \right) dt \\ (15) \quad &= (n-1)! \text{area}(\Omega_x^*(1)) < \infty \end{aligned}$$

since $\Omega_x^*(1)$ is a bounded subset of $\mathbf{V}_x^*(1)$. Since

$$\text{area}(\Omega_x^*(n)) = n^{n-1} \text{area}(\Omega_x^*(1)),$$

the formula in (15) implies:

$$(16) \quad f(x) = \frac{n!}{n^n} \text{area}(\Omega_x^*(n))$$

Let $\Omega_{\mathbb{C}}$ denote the *tube domain* $\Omega + \sqrt{-1}\mathbf{V} \subset \mathbf{V} \otimes_{\mathbb{R}} \mathbb{C}$. Then the integral defining $f(x)$ converges absolutely for every $x \in \Omega_{\mathbb{C}}$. Therefore

$\Omega \xrightarrow{f} \mathbb{R}$ extends to a holomorphic function $\Omega_{\mathbb{C}} \rightarrow \mathbb{C}$, from which it follows that f is real analytic on Ω .

LEMMA 4.4.4. *The function $f(x) \rightarrow +\infty$ as $x \rightarrow \partial\Omega$.*

PROOF. Consider a sequence $\{x_n\}_{n>0}$ in Ω converging to $x_\infty \in \partial\Omega$. Then the functions

$$\begin{aligned} \Omega^* &\xrightarrow{F_k} \mathbb{R} \\ \psi &\mapsto e^{-\psi(x_k)} \end{aligned}$$

are nonnegative functions converging uniformly to F_∞ on every compact subset of Ω^* so that

$$\liminf f(x_k) = \liminf \int_{\Omega^*} F_k(\psi) d\psi \geq \int_{\Omega^*} F_\infty(\psi) d\psi.$$

Suppose that $\psi_0 \in \mathbf{V}^*$ defines a supporting hyperplane to Ω at x_∞ ; then $\psi_0(x_\infty) = 0$. Let $K \subset \Omega^*$ be a closed ball; then $K + \mathbb{R}_+\psi_0$ is a cylinder in Ω^* with cross-section $K_1 = K \cap \psi_0^{-1}(c)$ for some $c > 0$.

$$\begin{aligned} \int_{\Omega^*} F_\infty(\psi) d\psi &\geq \int_{K + \mathbb{R}_+\psi_0} e^{-\psi(x_\infty)} d\psi \\ &\geq \int_{K_1} \left(\int_0^\infty dt \right) e^{-\psi(x_\infty)} d\psi_1 = \infty \end{aligned}$$

where $d\psi_1$ is a volume form on K_1 . □

LEMMA 4.4.5. *If $\gamma \in \text{Aff}(\Omega) \subset \text{GL}(\mathbf{V})$ is an automorphism of Ω , then*

$$(17) \quad f \circ \gamma = \det(\gamma)^{-1} \cdot f$$

In other words, if dx is a parallel volume form on \mathbf{A} , then $f(x) dx$ defines an $\text{Aff}(\Omega)$ -invariant volume form on Ω .

PROOF.

$$\begin{aligned} f(\gamma x) &= \int_{\Omega^*} e^{-\psi(\gamma x)} d\psi \\ &= \int_{\gamma^{-1}\Omega^*} e^{-\psi(x)} \gamma^* d\psi \\ &= \int_{\Omega^*} e^{-\psi(x)} (\det \gamma)^{-1} d\psi \\ &= (\det \gamma)^{-1} f(x) \end{aligned}$$

□

4.4.2. The covector field. Since $\det(\gamma)$ is a constant, it follows from (17) that $\log f$ transforms under γ by the additive constant $\log \det(\gamma)^{-1}$ and thus

$$\alpha = d \log f = f^{-1} df$$

is an $\text{Aff}(\Omega)$ -invariant closed 1-form on Ω . Furthermore, taking γ to be the homothety

$$x \xrightarrow{h_s} sx,$$

implies:

$$f \circ h_s = s^{-n} \cdot f,$$

which by differentiation with respect to s yields:

$$\alpha(\mathbf{R}_V) = -n.$$

Let $X \in \mathbb{T}_x \Omega \cong \mathbf{V}$ be a tangent vector; then $df(x) \in \mathbb{T}_x^* \Omega$ maps

$$X \mapsto - \int_{\Omega^*} \psi(X) e^{-\psi(x)} d\psi.$$

Using the identification $\mathbb{T}_x^* \Omega \cong \mathbf{V}^*$ we obtain a map

$$\begin{aligned} \Phi : \Omega &\longrightarrow \mathbf{V}^* \\ x &\mapsto -d \log f(x). \end{aligned}$$

As a linear functional, $\Phi(x)$ maps $X \in \mathbf{V}$ to

$$\frac{\int_{\Omega^*} \psi(X) e^{-\psi(x)} d\psi}{\int_{\Omega^*} e^{-\psi(x)} d\psi}$$

so if $X \in \Omega$, the numerator is positive and $\Phi(x) > 0$ on Ω . Thus $\Phi : \Omega \longrightarrow \Omega^*$. Furthermore by decomposing the volume form on Ω^* we obtain

$$\begin{aligned} \Phi(x) &= \frac{\int_0^\infty e^{-t} t^n \left(\int_{\Omega_x^*(1)} \psi_1 d\psi_1 \right) dt}{\int_0^\infty e^{-t} t^{n-1} \left(\int_{\Omega_x^*(1)} d\psi_1 \right) dt} \\ &= n \frac{\int_{\Omega_x^*(1)} \psi_1 d\psi_1 dt}{\int_{\Omega_x^*(1)} d\psi_1 dt} \\ &= n \text{centroid}(\Omega_x^*(1)). \end{aligned}$$

Since

$$\Phi(x) \in n \cdot \Omega_x^*(1) = \Omega_x^*(n),$$

that is, $\Phi(x) : x \mapsto n$,

$$(18) \quad \Phi(x) = \text{centroid}(\Omega_x^*(n)).$$

4.4.3. The metric tensor. For any function $\Omega \xrightarrow{f} \mathbb{R}$, the logarithmic Hessian

$$d^2 \log f = \nabla d \log f = \nabla \alpha$$

is an $\text{Aff}(\Omega)$ -invariant symmetric 2-form on Ω and equals:

$$d^2(\log f) = \nabla(f^{-1}df) = f^{-1}d^2f - (f^{-1}df)^2.$$

Furthermore the value of $d^2f(x) \in \text{Sym}^2\mathbb{T}_x^*\Omega$ on a pair

$$(X, Y) \in \mathbb{T}_x\Omega \times \mathbb{T}_x\Omega = \mathbb{V} \times \mathbb{V}$$

equals

$$\int_{\Omega^*} \psi(X)\psi(Y)e^{-\psi(x)} d\psi$$

PROPOSITION 4.4.6. $d^2 \log f$ is positive definite and defines an $\text{Aff}(\Omega)$ -invariant Riemannian metric on Ω .

PROOF.

$$\begin{aligned} f(x)^2 (d^2 \log f(x))(X, X) &= \int_{\Omega^*} e^{-\psi(x)} d\psi \int_{\Omega^*} \psi(X)^2 e^{-\psi(x)} d\psi \\ &\quad - \left(\int_{\Omega^*} \psi(X) e^{-\psi(x)} d\psi \right)^2 \\ &= \|e^{-\psi(x)/2}\|_2^2 \|\psi(X)e^{-\psi(x)/2}\|_2^2 \\ &\quad - \langle e^{-\psi(x)/2}, \psi(X)e^{-\psi(x)/2} \rangle_2^2 > 0 \end{aligned}$$

by the Schwartz inequality, since the functions

$$\begin{aligned} \psi &\longmapsto e^{-\psi(x)/2}, \\ \psi &\longmapsto \psi(X)e^{-\psi(x)/2} \end{aligned}$$

on Ω^* are not proportional. (Here $\langle \cdot, \cdot \rangle_2$ and $\|\cdot\|_2$ respectively denote the usual L^2 inner product and norm on $(\Omega^*, d\psi)$.) Thus $d^2 \log f$ is positive definite as claimed. \square

We characterize the linear functional $\Phi(x) \in \Omega^*$ quite simply as follows. Since $\Phi(x)$ is parallel to $df(x)$, each of its level hyperplanes is parallel to the tangent plane of the level set S_x of $\Omega \xrightarrow{f} \mathbb{R}$ containing x . Note that $\Phi(x)(x) = n$.

PROPOSITION 4.4.7. *The tangent space to the level set S_x of $\Omega \xrightarrow{f} \mathbb{R}$ at x equals $\Phi(x)^{-1}(n)$.*

This characterization yields the following result:

THEOREM 4.4.8. $\Omega \xrightarrow{\Phi} \Omega^*$ is bijective.

PROOF. Let $\psi_0 \in \Omega^*$ and let

$$Q_0 := \{z \in \mathbf{V} \mid \psi_0(z) = n\}.$$

Then the restriction of $\log f$ to the affine hyperplane Q_0 is a convex function which approaches $+\infty$ on $\partial(Q_0 \cap \Omega)$. Therefore the restriction $f|_{Q_0 \cap \Omega}$ has a unique critical point x_0 , which is necessarily a minimum. Then $T_{x_0} S_{x_0} = Q_0$ from which Proposition 4.4.7 implies that $\Phi(x_0) = \psi_0$. Furthermore, if $\Phi(x) = \psi_0$, then $f|_{Q_0 \cap \Omega}$ has a critical point at x so $x = x_0$. Therefore $\Omega \xrightarrow{\Phi} \Omega^*$ is bijective as claimed. \square

If $\Omega \subset \mathbf{V}$ is a properly convex cone and Ω^* is its dual, then let $\Phi_{\Omega^*} : \Omega^* \rightarrow \Omega$ be the diffeomorphism $\Omega^* \rightarrow \Omega^{**} = \Omega$ defined above. If $x \in \Omega$, then $\psi = (\Phi^*)^{-1}(x)$ is the unique $\psi \in \mathbf{V}^*$ such that:

- $\psi(x) = n$;
- The centroid of $\Omega \cap \psi^{-1}(n)$ equals x .

The duality isomorphism $\mathbf{GL}(\mathbf{V}) \rightarrow \mathbf{GL}(\mathbf{V}^*)$ (given by inverse transpose of matrices) defines an isomorphism $\text{Aff}(\Omega) \rightarrow \text{Aff}(\Omega^*)$. Let

$$\begin{aligned} \Omega &\xrightarrow{\Phi_{\Omega}} \Omega^*, \\ \Omega^* &\xrightarrow{\Phi_{\Omega^*}} \Omega^{**} = \Omega \end{aligned}$$

be the duality maps for Ω and Ω^* respectively. Vinberg points out in [278] that, in general, the composition

$$\Omega \xrightarrow{\Phi_{\Omega^*} \circ \Phi_{\Omega}} \Omega$$

is not the identity. However, if Ω is *homogeneous* (that is, $\text{Aff}(\Omega) \subset \mathbf{GL}(\mathbf{V})$ acts transitively on Ω), then $\Omega_{\Omega^*} \circ \Phi_{\Omega} = \mathbb{I}$:

PROPOSITION 4.4.9 (Vinberg [278]). *Let $\Omega \subset \mathbf{V}$ be a homogeneous properly convex cone. Then Φ_{Ω^*} and Φ_{Ω} are inverse maps $\Omega^* \longleftrightarrow \Omega$.*

PROOF. Let $x \in \Omega$ and $Y \in \mathbf{V} \cong T_x \Omega$ be a tangent vector. Denote the value of the canonical Riemannian metric $\nabla \alpha = d^2 \log f$ at x by:

$$T_x \Omega \times T_x \Omega \xrightarrow{g_x} \mathbb{R}$$

Then the differential of $\Omega \xrightarrow{\Phi_{\Omega}} \Omega^*$ at x equals the composition

$$T_x \Omega \xrightarrow{\tilde{g}} T_x^* \Omega \cong \mathbf{V}^* \cong T_{\Phi(x)} \Omega^*$$

where $\Omega \xrightarrow{\tilde{g}_x} T_x^* \Omega$ is the linear isomorphism corresponding to g_x and the isomorphisms

$$T_x^* \Omega \cong \mathbf{V}^* \cong T_{\Phi(x)} \Omega^*$$

are induced by parallel translation. Taking the directional derivative of the equation

$$\alpha_x(\mathbf{R}_x) = -n$$

with respect to $Y \in \mathbf{V} \cong \mathbf{T}_x\Omega$ yields:

$$\begin{aligned} 0 &= (\nabla_Y \alpha)(\mathbf{R}) + \alpha(\nabla_Y \mathbf{R}) \\ &= g_x(\mathbf{R}_x, Y) + \alpha_x(Y) \\ (19) \quad &= g_x(x, Y) - \Phi(x)(Y). \end{aligned}$$

Let $\Omega \xrightarrow{f_\Omega} \mathbb{R}$ and $\Omega^* \xrightarrow{f_{\Omega^*}} \mathbb{R}$ be the characteristic functions for Ω and Ω^* respectively. Then $f_\Omega(x) dx$ is a volume form on Ω invariant under $\text{Aff}(\Omega)$ and $f_{\Omega^*}(\psi) d\psi$ is a volume form on Ω^* invariant under the induced action of $\text{Aff}(\Omega)$ on Ω^* . Moreover $\Omega \xrightarrow{\Phi} \Omega^*$ is equivariant with respect to the isomorphism $\text{Aff}(\Omega) \rightarrow \text{Aff}(\Omega^*)$. Therefore the tensor field on Ω defined by

$$\begin{aligned} &f_\Omega(x) dx \otimes (f_{\Omega^*} \circ \Phi)(x) d\psi \\ &\in \wedge^n \mathbf{T}_x \Omega \otimes \wedge^n \mathbf{T}_{\Phi(x)} \Omega^* \\ &\cong \wedge^n \mathbf{V} \otimes \wedge^n \mathbf{V}^* \end{aligned}$$

is $\text{Aff}(\Omega)$ -invariant. Since the parallel tensor field $dx \otimes d\psi \in \wedge^n \mathbf{V} \otimes \wedge^n \mathbf{V}^*$ is invariant under all of $\text{Aff}(\mathbf{V})$, the coefficient

$$(20) \quad h(x) = f_\Omega(x) dx \otimes (f_{\Omega^*} \circ \Phi)(x) d\psi$$

is an $\text{Aff}(\Omega)$ -invariant function on Ω . Since Ω is homogeneous, h is constant.

Differentiating $\log h$ using (20),

$$0 = d \log h = d \log f_\Omega(x) + d \log (f_{\Omega^*} \circ \Phi)(x).$$

Since $d \log f_{\Omega^*}(\psi) = \Phi_{\Omega^*}(\psi)$,

$$\begin{aligned} 0 &= -\Phi(x)(Y) + \Phi_{\Omega^*}(d\Phi(Y)) \\ &= -\Phi(x)(Y) + g_x(Y, \Phi_{\Omega^*} \circ \Phi_\Omega(x)) \end{aligned}$$

Combining this equation with (19) yields:

$$\Phi_{\Omega^*} \circ \Phi_\Omega(x) = x$$

as desired. □

Thus, if Ω is a homogeneous cone, then $\Phi(x) \in \Omega^*$ is the centroid of the cross-section $\Omega_x^*(n) \subset \Omega^*$ in \mathbf{V}^* .

4.5. Benzécri's Compactness Theorem

Let $\mathbf{P} = \mathbf{P}(\mathbf{V})$ and $\mathbf{P}^* = \mathbf{P}(\mathbf{V}^*)$ be the associated projective spaces. Then the projectivization $\mathbf{P}(\Omega) \subset \mathbf{P}$ of Ω is by definition a *properly convex domain* and its closure $K = \overline{\mathbf{P}(\Omega)}$ a *convex body*. Then the *dual convex body* K^* equals the closure of the projectivization $\mathbf{P}(\Omega^*)$ consisting of all hyperplanes $H \subset \mathbf{P}$ such that $\overline{\Omega} \cap H = \emptyset$. A *pointed convex body* consists of a pair (K, x) where K is a convex body and $x \in \text{int}(K)$ is an interior point of K . Let $H \subset \mathbf{P}$ be a hyperplane and $\mathbf{A} = \mathbf{P} \setminus H$ its complementary affine space. We say that the pointed convex body (K, u) is *centered relative to* \mathbf{A} (or H) if u is the centroid of K in the affine geometry of \mathbf{A} .

PROPOSITION 4.5.1. *Let (K, u) be a pointed convex body in a projective space \mathbf{P} . Then there exists a hyperplane $H \subset \mathbf{P}$ disjoint from K such that in the affine space $\mathbf{A} = \mathbf{P} \setminus H$, the centroid of $K \subset \mathbf{A}$ equals u .*

PROOF. Let $\mathbf{V} = \mathbf{V}(\mathbf{P})$ be the vector space corresponding to the projective space \mathbf{P} and let $\Omega \subset \mathbf{V}$ be a properly convex cone whose projectivization is the interior of K . Let $x \in \Omega$ be a point corresponding to $u \in \text{int}(K)$. Let

$$\Omega^* \xrightarrow{\Phi_{\Omega^*}} \Omega$$

be the duality map for Ω^* and let $\psi = (\Phi_{\Omega^*})^{-1}(y)$. Then the centroid of the cross-section

$$\Omega_\psi(n) = \{x \in \Omega \mid \psi(x) = n\}$$

in the affine hyperplane $\psi^{-1}(n) \subset \mathbf{V}$ equals y . Let $H = \mathbf{P}(\text{Ker}(\psi))$ be the projective hyperplane in \mathbf{P} corresponding to ψ ; then projectivization defines an affine isomorphism

$$\psi^{-1}(n) \longrightarrow \mathbf{P} \setminus H$$

mapping $\Omega_\psi(n) \longrightarrow K$. Since affine maps preserve centroids, it follows that (K, u) is centered relative to H . \square

Thus every pointed convex body (K, u) is centered relative to a unique affine space containing K . In dimension one, this means the following: let $K \subset \mathbb{R}\mathbf{P}^1$ be a closed interval $[a, b] \subset \mathbb{R}$ and let $a < x < b$ be an interior point. Then x is the midpoint of $[a, b]$ relative to the “hyperplane” H obtained by projectively reflecting x with respect to the pair $\{a, b\}$:

$$H = R_{[a,b]}(x) = \frac{(a+b)x - 2ab}{2x - (a+b)}$$

An equivalent version of Proposition 4.5.1 involves using collineations to “move a pointed convex body” into affine space to center it:

PROPOSITION 4.5.2. *Let $K \subset E$ be a convex body in an affine space and let $x \in \text{int}(A)$ be an interior point. Let $\mathbf{P} \supset E$ be the projective space containing A . Then there exists a collineation $\mathbf{P} \xrightarrow{g} \mathbf{P}$ such that:*

- $g(K) \subset A$;
- $(g(K), g(x))$ is centered relative to A .

The one-dimensional version is really just the fundamental theorem of projective geometry: if $[a, b]$ is a closed interval with interior point x , then the unique collineation mapping

$$\begin{aligned} a &\mapsto -1 \\ x &\mapsto 0 \\ b &\mapsto 1 \end{aligned}$$

centers $[a, b]$ at x .

PROPOSITION 4.5.3. *Let $K_i \subset A$ be convex bodies ($i = 1, 2$) in an affine space A with centroids u_i , and suppose that $\mathbf{P} \xrightarrow{g} \mathbf{P}$ is a collineation such that $g(K_1) = K_2$ and $g(u_1) = u_2$. Then g is an affine automorphism of A , that is, $g(A) = A$.*

PROOF. Let V be a vector space containing A as an affine hyperplane and let Ω_i be the properly convex cones in V whose projective images are K_i . By assumption there exists a linear map $V \xrightarrow{\tilde{g}} V$ and points $x_i \in \Omega_i$ mapping to $u_i \in K_i$ such that $\tilde{g}(\Omega_1) = \Omega_2$ and $\tilde{g}(x_1) = x_2$. Let $S_i \subset \Omega_i$ be the level set of the characteristic function

$$\Omega_i \xrightarrow{f_i} \mathbb{R}$$

containing x_i . Since (K_i, u_i) is centered relative to A , it follows that the tangent plane

$$T_{x_i} S_i = E \subset V.$$

Since the construction of the characteristic function is linearly invariant, it follows that $\tilde{g}(S_1) = S_2$. Moreover

$$\tilde{g}(T_{x_1} S_1) = T_{x_2} S_2,$$

that is, $\tilde{g}(A) = E$ and $g \in \text{Aff}(A)$ as desired. \square

4.5.1. Convex bodies in projective space. Let $\mathfrak{C}(\mathbb{P})$ denote the set of all convex bodies in \mathbb{P} , with the topology induced from the Hausdorff metric on the set of all closed subsets of \mathbb{P} (which is induced from the Fubini-Study metric on \mathbb{P} , see §Let

$$\mathfrak{C}_*(\mathbb{P}) = \{(K, x) \in \mathfrak{C}(\mathbb{P}) \times \mathbb{P} \mid x \in \text{int}(K)\}$$

be the corresponding set of pointed convex bodies, with a topology induced from the product topology on $\mathfrak{C}(\mathbb{P}) \times \mathbb{P}$. The collineation group G acts continuously on $\mathfrak{C}(\mathbb{P})$ and on $\mathfrak{C}_*(\mathbb{P})$.

Recall that an action of a group Γ on a space X is *syndetic* if $\exists K \subset\subset X$ such that $\Gamma K = X$. Furthermore the action is *proper* if the corresponding map

$$\begin{aligned} \Gamma \times X &\longrightarrow X \times X \\ (\gamma, x) &\longmapsto (\gamma x, x) \end{aligned}$$

is a proper map (inverse images of compact subsets are compact). See §A.2 for discussion of elementary properties of group actions.

THEOREM 4.5.4 (Benzécri). *The collineation group G acts properly and syndetically on $\mathfrak{C}_*(\mathbb{P})$. In particular the quotient $\mathfrak{C}_*(\mathbb{P})/G$ is a compact Hausdorff space.*




FIGURE 4.4. A sequence of projectively equivalent convex domains with a corner converging to a triangle with the corner as a vertex.




FIGURE 4.5. A sequence of projectively equivalent convex domains with a flat part of the boundary converging to a triangle having the flat part as a side.

4.5.2. How convex bodies can degenerate. While the quotient $\mathfrak{C}_*(\mathbb{P})/G$ is Hausdorff, the space of equivalence classes of convex bodies $\mathfrak{C}(\mathbb{P})/G$ is generally *not* Hausdorff. Here are three basic examples.

4.5.2.1. *Corners.* Suppose that Ω is a properly convex domain whose boundary is not C^1 at a point x_1 . Then $\partial\Omega$ has a “corner” at x_1 and we may choose homogeneous coordinates so that $x_1 = [1 : 0 : 0]$ and $\bar{\Omega}$ lies in the domain

$$\Delta = \{[x : y : z] \in \mathbb{RP}^2 \mid x, y, z > 0\}$$

in such a way that $\partial\Omega$ is tangent to $\partial\Delta$ at x_1 . Under the one-parameter group of collineations defined by

$$g_t = \begin{bmatrix} e^{-t} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^t \end{bmatrix}$$

as $t \rightarrow +\infty$, the domains $g_t\Omega$ converge to Δ . (Compare Figure 4.4.) Then the G -orbit of $\bar{\Omega}$ in $\mathfrak{C}(\mathbb{P})$ is not closed. The corresponding equivalence class of $\bar{\Omega}$ is not a closed point in $\mathfrak{C}(\mathbb{P})/G$ unless Ω was already a triangle.

4.5.2.2. *Flats.* Similarly suppose that Ω is a properly convex domain which is not *strictly convex*, that is, its boundary contains a non-trivial line segment σ . (We suppose that σ is a maximal line segment contained in $\partial\Omega$.) As above, we may choose homogeneous coordinates so that $\Omega \subset \Delta$ and such that $\bar{\Omega} \cap \bar{\Delta} = \bar{\sigma}$ and σ lies on the line $\{[x : y : 0] \mid x, y \in \mathbb{R}\}$. As $t \rightarrow +\infty$ the image of Ω under the collineation

$$g_t = \begin{bmatrix} e^{-t} & 0 & 0 \\ 0 & e^{-t} & 0 \\ 0 & 0 & e^{2t} \end{bmatrix}$$

converges to a triangle region with vertices $\{[0 : 0 : 1]\} \cup \partial\sigma$. As above, the equivalence class of $\bar{\Omega}$ in $\mathfrak{C}(\mathbb{P})/G$ is not a closed point in $\mathfrak{C}(\mathbb{P})/G$ unless Ω is a triangle.

4.5.2.3. *Osculating conics.* As a final example, consider a properly convex domain Ω with C^1 boundary such that there exists a point $u \in \partial\Omega$ such that $\partial\Omega$ is C^2 at u . In that case a conic C osculates $\partial\Omega$ at u . Choose homogeneous coordinates such that $u = [1 : 0 : 0]$ and

$$C = \{[x : y : z] \mid xy + z^2 = 0\}.$$

Then as $t \rightarrow +\infty$ the image of Ω under the collineation

$$g_t = \begin{bmatrix} e^{-t} & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

converges to the convex region

$$\{[x : y : z] \mid xy + z^2 < 0\}$$

bounded by C . As above, the equivalence class of $\bar{\Omega}$ in $\mathfrak{C}(\mathbb{P})/G$ is not a closed point in $\mathfrak{C}(\mathbb{P})/G$ unless $\partial\Omega$ is a conic.

In summary:

PROPOSITION 4.5.5. *Suppose $\bar{\Omega} \subset \mathbb{RP}^2$ is a convex body whose equivalence class $[\bar{\Omega}]$ is a closed point in $\mathfrak{C}(\mathbb{P})/G$. Suppose that $\partial\Omega$ is neither a triangle nor a conic. Then $\partial\Omega$ is a C^1 strictly convex curve which is nowhere C^2 .*

The forgetful map $\mathfrak{C}_*(\mathbb{P}) \xrightarrow{P_i} \mathfrak{C}(\mathbb{P})$ which forgets the point of a pointed convex body is induced from Cartesian projection $\mathfrak{C}(\mathbb{P}) \times \mathbb{P} \rightarrow \mathfrak{C}(\mathbb{P})$.

THEOREM 4.5.6 (Benzécri). *Let $\Omega \subset \mathbb{P}$ is a properly convex domain such that there exists a subgroup $\Gamma \subset \text{Aut}(\Omega)$ which acts syndetically on Ω . Then the corresponding point $[\bar{\Omega}] \in \mathfrak{C}(\mathbb{P})/G$ is closed.*

In the following result, all but the continuous differentiability of the boundary in the following result was originally proved in Kuiper [192] using a somewhat different technique; the C^1 statement is due to Benzécri [35] as well as the proof given here.

COROLLARY 4.5.7. *Suppose that $M = \Omega/\Gamma$ is a convex \mathbb{RP}^2 -manifold such that $\chi(M) < 0$. Then either the \mathbb{RP}^2 -structure on M is a hyperbolic structure or the boundary $\partial\Omega$ of its universal covering is a C^1 strictly convex curve which is nowhere C^2 .*

PROOF. Apply Proposition 4.5.5 to Theorem 4.5.6. □

PROOF OF THEOREM 4.5.6 ASSUMING THEOREM 4.5.4. Let Ω be a properly convex domain with an automorphism group $\Gamma \subset \text{Aff}(\Omega)$ acting syndetically on Ω . It suffices to show that the G -orbit of $\{\bar{\Omega}\}$ in $\mathfrak{C}(\mathbb{P})$ is closed, which is equivalent to showing that the G -orbit of

$$\Pi^{-1}(\{\bar{\Omega}\}) = \{\bar{\Omega}\} \times \Omega$$

in $\mathfrak{C}_*(\mathbb{P})$ is closed. This is equivalent to showing that the image of

$$\{\bar{\Omega}\} \times \Omega \subset \mathfrak{C}_*(\mathbb{P})$$

under the quotient map $\mathfrak{C}_*(\mathbb{P}) \longrightarrow \mathfrak{C}_*(\mathbb{P})/G$ is closed. Let $K \subset \Omega$ be a compact subset such that $\Gamma K = \Omega$; then $\{\bar{\Omega}\} \times K$ and $\{\bar{\Omega}\} \times \Omega$ have the same image in $\mathfrak{C}_*(\mathbb{P})/\Gamma$ and hence in $\mathfrak{C}_*(\mathbb{P})/G$. Hence it suffices to show that the image of $\{\bar{\Omega}\} \times K$ in $\mathfrak{C}_*(\mathbb{P})/G$ is closed. Since K is compact and the composition

$$K \longrightarrow \{\bar{\Omega}\} \times K \hookrightarrow \{\bar{\Omega}\} \times \Omega \subset \mathfrak{C}_*(\mathbb{P}) \longrightarrow \mathfrak{C}_*(\mathbb{P})/G$$

is continuous, it follows that the image of K in $\mathfrak{C}_*(\mathbb{P})/G$ is compact. By Theorem 4.5.4, $\mathfrak{C}_*(\mathbb{P})/G$ is Hausdorff and hence the image of K in $\mathfrak{C}_*(\mathbb{P})/G$ is closed, as desired. The proof of Theorem 4.5.6 (assuming Theorem 4.5.4) is now complete. \square

Now we prove Theorem 4.5.4. Choose a fixed hyperplane $H_\infty \subset \mathbb{P}$ and let $\mathbb{A} = \mathbb{P} \setminus H_\infty$ be the corresponding affine patch and $\text{Aff}(\mathbb{A})$ the group of affine automorphisms of \mathbb{A} . Let $\mathfrak{C}(\mathbb{A}) \subset \mathfrak{C}(\mathbb{P})$ denote the set of convex bodies $K \subset E$, with the induced topology. (Note that the $\mathfrak{C}(\mathbb{A})$ is a complete metric space with respect to the Hausdorff metric induced from the Euclidean metric on \mathbb{E} and we may use this metric to define the topology on $\mathfrak{C}(\mathbb{A})$. The inclusion map $\mathfrak{C}(\mathbb{A}) \hookrightarrow \mathfrak{C}(\mathbb{P})$ is continuous, although not uniformly continuous.) We define a map

$$\mathfrak{C}(\mathbb{A}) \xrightarrow{\iota} \mathfrak{C}_*(\mathbb{P})$$

as follows. Let $K \in \mathfrak{C}(\mathbb{A})$ be a convex body in affine space \mathbb{A} ; let $\iota(K)$ to be the pointed convex body

$$\iota(K) = (K, \text{centroid}(K)) \in \mathfrak{C}_*(\mathbb{P});$$

clearly ι is equivariant respecting the embedding $\text{Aff}(\mathbb{A}) \hookrightarrow G$.

4.5.3. Reduction to the affine case.

THEOREM 4.5.8. *Let $\mathbb{A} \subset \mathbb{P}$ be an affine patch in projective space. Then the map*

$$\begin{aligned} \mathfrak{C}(\mathbb{A}) &\xrightarrow{\iota} \mathfrak{C}_*(\mathbb{P}) \\ K &\longmapsto (K, \text{centroid}(K)) \end{aligned}$$

is equivariant with respect to the inclusion $\text{Aff}(\mathbb{A}) \longrightarrow G$ and the corresponding homomorphism of topological transformation groupoids

$$(\mathfrak{C}(\mathbb{A}), \text{Aff}(\mathbb{A})) \xrightarrow{\iota} (\mathfrak{C}_*(\mathbb{P}), G)$$

is an equivalence of groupoids.

PROOF. The surjectivity of $\mathfrak{C}(\mathbb{A})/\text{Aff}(\mathbb{A}) \xrightarrow{\iota_*} \mathfrak{C}_*(\mathbb{P})/G$ follows immediately from Proposition 4.4.9 and the bijectivity of

$$\text{Hom}(a, b) \xrightarrow{\iota_*} \text{Hom}(\iota(a), \iota(b))$$

follows immediately from Proposition 4.5.2. \square

Thus the proof of Proposition 4.5.3 reduces (via Lemma A.3.1 and Theorem 4.5.8) to the following:

THEOREM 4.5.9. *Aff(A) acts properly and syndetically on $\mathfrak{C}(A)$.*

Let $\mathcal{E} \subset \mathfrak{C}(A)$ denote the subspace of ellipsoids in A ; the affine group $\text{Aff}(A)$ acts transitively on \mathcal{E} with isotropy group the orthogonal group — in particular this action is proper. If $K \in \mathfrak{C}(A)$ is a convex body, then there exists a unique ellipsoid $\text{ell}(K) \in \mathcal{E}$ (*the ellipsoid of inertia of K*) such that for each affine map $\psi : A \rightarrow \mathbb{R}$ such that $\psi(\text{centroid}(K)) = 0$ the moments of inertia satisfy:

$$\int_K \psi^2 dx = \int_{\text{ell}(K)} \psi^2 dx$$

PROPOSITION 4.5.10. *Taking the ellipsoid-of-inertia of a convex body*

$$\mathfrak{C}(A) \xrightarrow{\text{ell}} \mathcal{E}$$

defines an Aff(A)-invariant proper retraction of $\mathfrak{C}(A)$ onto \mathcal{E} .

PROOF OF THEOREM 4.5.9 ASSUMING PROPOSITION 4.5.10.

Since $\text{Aff}(A)$ acts properly and syndetically on \mathcal{E} and ell is a proper map, it follows that $\text{Aff}(A)$ acts properly and syndetically on $\mathfrak{C}(A)$. \square

PROOF OF PROPOSITION 4.5.10. ell is clearly affinely invariant and continuous. Since $\text{Aff}(A)$ acts transitively on \mathcal{E} , it suffices to show that a single fiber $\text{ell}^{-1}(e)$ is compact for $e \in \mathcal{E}$. We may assume that e is the unit sphere in E centered at the origin 0 . Since the collection of compact subsets of E which lie between two compact balls is compact subset of $\mathfrak{C}(A)$, Proposition 4.5.10 will follow from: \square

PROPOSITION 4.5.11. *For each n there exist constants $0 < r(n) < R(n)$ such that every convex body $K \subset \mathbb{R}^n$ whose centroid is the origin and whose ellipsoid-of-inertia is the unit sphere satisfies*

$$B_{r(n)}(O) \subset K \subset B_{R(n)}(O).$$

Cooper-Long-Tillmann [78] call such an affine chart a *Benzécri chart*, and in [79] provide an algorithm for computing a Benzécri chart.

The proof of Proposition 4.5.11 is based on:

LEMMA 4.5.12. *Let $K \subset A$ be a convex body with centroid O . Suppose that l is a line through O which intersects ∂K in the points X, X' . Then*

$$(21) \quad \frac{1}{n} \leq \frac{OX}{OX'} \leq n.$$

PROOF. Let $\psi \in \mathbf{A}^*$ be a linear functional such that $\psi(X) = 0$ and $\psi^{-1}(1)$ is a supporting hyperplane for K at X' ; then necessarily $0 \leq \psi(x) \leq 1$ for all $x \in K$. We claim that

$$(22) \quad \psi(O) \leq \frac{n}{n+1}.$$

For $t \in \mathbb{R}$ let

$$\begin{aligned} \mathbf{A} &\xrightarrow{h_t} \mathbf{A} \\ x &\mapsto t(x - X) + X \end{aligned}$$

be the homothety fixing X having strength t . We compare the linear functional ψ with the “polar coordinates” on K defined by the map

$$\begin{aligned} [0, 1] \times \partial K &\xrightarrow{F} K \\ (t, \mathbf{s}) &\mapsto h_t \mathbf{s} \end{aligned}$$

which is bijective on $(0, 1] \times \partial K$ and collapses $\{0\} \times \partial K$ onto X . Thus there is a well-defined function $K \xrightarrow{\mathbf{t}} \mathbb{R}$ such that for each $x \in K$, there exists $x' \in \partial K$ such that $x = F(\mathbf{t}, x')$. Since $0 \leq \psi(F(t, \mathbf{s})) \leq 1$, it follows that for $x \in K$,

$$0 \leq \psi(x) \leq \mathbf{t}(x)$$

Let $\mu = \mu_K$ denote the probability measure supported on K defined by

$$\mu(S) = \frac{\int_{S \cap K} dx}{\int_K dx}.$$

There exists a measure ν on ∂K such that for each measurable function $f : \mathbf{A} \rightarrow \mathbb{R}$

$$\int f(x) d\mu(x) = \int_{t=0}^1 \int_{\mathbf{s} \in \partial K} f(t\mathbf{s}) t^{n-1} d\nu(\mathbf{s}) dt,$$

that is, $F^* d\mu = t^{n-1} d\nu \wedge dt$.

The first moment of $K \xrightarrow{\mathbf{t}} [0, 1]$ is:

$$\bar{\mathbf{t}}(K) = \int_K \mathbf{t} d\mu = \frac{\int_K \mathbf{t} d\mu}{\int_K d\mu} = \frac{\int_0^1 t^n \int_{\partial K} d\nu dt}{\int_0^1 t^{n-1} \int_{\partial K} d\nu dt} = \frac{n}{n+1}$$

and since the value of the affine function ψ on the centroid equals the first moment of ψ on K , we have

$$0 < \psi(O) = \int_K \psi(x) d\mu(x) < \int_K \mathbf{t}(x) d\mu = \frac{n}{n+1}.$$

Now the distance function on the line $\overleftrightarrow{XX'}$ is affinely related to the linear functional ψ , that is, there exists a constant $c > 0$ such that for $x \in \overleftrightarrow{XX'}$ the distance $Xx = c|\psi(x)|$; since $\psi(X') = 1$ it follows that

$$\psi(x) = \frac{Xx}{XX'}$$

and since $OX + OX' = XX'$ it follows that

$$\frac{OX'}{OX} = \frac{XX'}{OX} - 1 \geq \frac{n+1}{n} - 1 = \frac{1}{n}.$$

This gives the second inequality of (21). The first inequality follows by reversing the roles of X, X' . \square

PROOF OF PROPOSITION 4.5.11. Let $X \in \partial K$ be a point at minimum distance from the centroid O ; then there exists a supporting hyperplane H at x which is orthogonal to \overleftrightarrow{OX} and let $\mathbf{A} \xrightarrow{\psi} \mathbb{R}$ be the corresponding linear functional of unit length. Let $a = \psi(X) > 0$ and $b = \psi(X') < 0$; Proposition 4.5.10 implies $-b < na$.

We claim that $0 < |\psi(x)| < na$ for all $x \in K$. To this end let $x \in K$; we may assume that $\psi(x) > 0$ since $-na < \psi(X') \leq \psi(x)$. Furthermore we may assume that $x \in \partial K$. Let $z \in \partial K$ be the other point of intersection of \overleftrightarrow{Ox} with ∂K ; then $\psi(z) < 0$. Now

$$\frac{1}{n} \leq \frac{Oz}{Ox} \leq n$$

implies that

$$\frac{1}{n} \leq \frac{|\psi(z)|}{|\psi(x)|} \leq n$$

(since the linear functional ψ is affinely related to signed distance along \overleftrightarrow{Ox}). Since $0 > \psi(z) \geq -a$, it follows that $|\psi(x)| \leq na$ as claimed.

Let w_n denote the moment of inertia of ψ for the unit sphere; then we have

$$w_n = \int_K \psi^2 d\mu \leq \int_K n^2 a^2 d\mu = n^2 a^2$$

whence $a \geq \sqrt{w_n}/n$. Taking $r(n) = \sqrt{w_n}/n$ we see that K contains the $r(n)$ -ball centered at O .

To obtain the upper bound, observe that if C is a right circular cone with vertex X , altitude h and base a sphere of radius ρ and $C \xrightarrow{t} [0, h]$ is the altitudinal distance from the base, then the integral

$$\int_C t^2 d\mu = \frac{2h^3 \rho^{n-1} v_{n-1}}{(n+2)(n+1)n}$$

where v_{n-1} denotes the $(n-1)$ -dimensional volume of the unit $(n-1)$ -ball. Let $X \in \partial K$ and C be a right circular cone with vertex X and base an $(n-1)$ -dimensional ball of radius $r(n)$. We have just seen that K contains $B_{r(n)}(O)$; it follows that $K \supset C$. Let $K \xrightarrow{t} \mathbb{R}$ be the unit-length linear functional vanishing on the base of C ; then

$$t(X) = h = OX.$$

Its second moment is

$$w_n = \int_K t^2 d\mu \geq \int_C t^2 d\mu = \frac{2h^3 r(n)^{n-1} v_{n-1}}{(n+2)(n+1)n}$$

and thus it follows that

$$OX = h \leq R(n)$$

where

$$R(n) = \left(\frac{(n+2)(n+1)nw_n}{2r(n)^{n-1}v_{n-1}} \right)^{\frac{1}{3}}$$

as desired. The proof is now complete. \square

EXERCISE 4.5.13. *The volume of the unit ball in \mathbb{R}^n equals:*

$$v_n = \begin{cases} \pi^{n/2}/(n/2)! & \text{for } n \text{ even} \\ 2^{(n+1)/2} \pi^{(n-1)/2} / (1 \cdot 3 \cdot 5 \cdots n) & \text{for } n \text{ odd} \end{cases}$$

Its moments of inertia are:

$$w_n = \begin{cases} v_n/(n+2) & \text{for } n \text{ even} \\ 2v_n/(n+2) & \text{for } n \text{ odd} \end{cases}$$

Based on Cooper-Long-Tillmann [78], explicit formulas are given in Casella-Tate-Tillmann[58] for the size of a *Benzécri chart*.

4.6. Quasi-homogeneous and divisible domains

Corollary 4.5.7 of Benzécri's Theorem ref provides sharp information on the geometry of a convex domain Ω on which $\text{Aut}(\Omega)$ acts syndetically. A convex domain $\Omega \subset \mathbb{P}$ is said to be *quasi-homogeneous* if $\text{Aut}(\Omega)$ acts syndetically, that is, the quotient $\Omega/\text{Aut}(\Omega)$ is compact (although not necessarily Hausdorff). If the action is *proper*, (that is, the quotient is Hausdorff), then the domain is said to be *divisible*. Although for simplicity we have only discussed the two-dimensional situation, much is known in this case, especially through a series of papers of Benoist, starting with his paper *Automorphismes des cônes convexes* [27], and continuing into his series of four *Convex divisible*

papers [30, 29, 31, 32]. He discusses these results, background and further advances (at the time) in his excellent survey paper [33] (which is written in English).

He begins by characterizing automorphisms of convex cones by the dynamical notion of *proximality*. A projective action of Γ on \mathbb{P} is *proximal* if $\exists \gamma \in \Gamma$ with a unique attracting fixed point. From this he finds restrictions on which Zariski closures Γ can occur.

A crucial idea is that when Ω is *strictly convex*, the natural geodesic flow defined by the Hilbert metric is an *Anosov flow*, that is, it preserves a decomposition of the tangent bundle as a direct sum of (1) the line bundle generating the flow; (2) a subbundle of tangent vectors exponential expanded by the flow (with respect to a fixed Riemannian structure); (3) a subbundle of tangent bundle exponentially contracted by the flow. From this he deduces that Γ is a hyperbolic group in the sense of Gromov. Furthermore $\partial\Omega$ is $C^{1+\alpha}$ for some $\alpha > 0$.

When Ω is *not* strictly convex (such as the triangle) much of this breaks down. However, as Benzécri noted, Theorem ?? implies that if Ω is not strictly convex, then Ω contains a properly embedded triangle. A simple example¹ is a deformation of a Coxeter group Γ_0 built on a regular ideal tetrahedron in $\mathbb{H}^3 \subset \mathbb{RP}^3$. Deformations Γ_t exist of this (noncompact) convex \mathbb{RP}^3 -orbifold where the cusps of Γ_t , where $t > 0$, are the regular (3, 3, 3)-triangle tessellations of a triangle discussed in §4.2.2 and depicted in Figure 2.3. In \mathbb{RP}^3 , these cusp groups preserve a projective hyperplane $H_t \subset \mathbb{RP}^3$. By adding reflections $R_1(t), R_2(t), R_3(t), R_4(t)$ in these hyperplanes one creates projective Coxeter groups Γ_t^* acting properly and syndetically on a properly convex domain Ω_t^* which is *not* strictly convex. (This is analogous to deforming the cusps in Thurston's hyperbolic Dehn surgery.) The quotient of Ω_t^* by a torsionfree finite-index subgroup of Γ_t^* is a closed 3-manifold with incompressible tori corresponding to the cusps of Γ_0 .

Benoist gives a comprehensive description of such divisible 3-domains in [32], and relates the non-strict convexity of the boundary to incompressible tori and the JSJ-decomposition of the quotient convex \mathbb{RP}^3 -manifolds. See Choi-Hodgson-Lee [72] for further discussion of divisible convex domains arising from Coxeter groups in \mathbb{RP}^3 .

In a series of papers [160, 159, 161, 162], Kyeonghee Jo investigates the differentiability of the boundary of a convex quasi-homogeneous domain. She shows that under various assumptions on the differentiability of $\partial\Omega$, such a domain must be homogeneous. For example, if Ω

¹The author gave a lecture on this example at a regional meeting of the American Mathematical Society on October 30, 1982.

is strictly convex quasi-homogeneous, then its boundary is at least C^1 , but if it is C^2 except at a finite set of points, then it must be an ellipsoid. (Other characterizations of ellipsoids among quasi-homogeneous convex domains are due to Socié-Méthou [256] and Colbois-Verovic [76].)

In [168] Kapovich gives examples of strictly convex divisible domains whose quotients are the non-locally symmetric negatively curved manifolds first discovered by Gromov and Thurston.

Part 2

Geometric manifolds

CHAPTER 5

Locally homogeneous geometric structures

Let M be a manifold and let X be a space with a transitive action of a Lie group G . Then in the spirit of Klein's Erlangen program [172], (G, X) defines a *geometry*: namely, the objects in X which are invariant under G . (A recent discussion of the Erlangen program is [158].)

We want to impart this geometry to M by a system of coordinate charts taking coordinate patches in M to open subsets of X , in such a way that the coordinate changes on overlapping patches are locally restrictions of transformations coming from G . Such a coordinate atlas defines a (G, X) -*structure* on M , and we call M a (G, X) -*manifold*.

The general notion of defining a structure on a manifold by an atlas of local charts is that of a *pseudogroup*. These are defined by a collection \mathcal{G} of homeomorphisms between open subsets of a topological space S satisfying several natural conditions: \mathcal{G} contains the identity \mathbb{I}_S and is closed under restrictions to open subsets, inversion and composition (where defined). Furthermore, if $U = \bigcup_{\alpha} U_{\alpha}$ and $g_{\alpha}, g_{\beta} \in \mathcal{G}$ are defined on U_{α} and U_{β} respectively, such that the restrictions

$$g_{\alpha}|_{U_{\alpha}} = g_{\beta}|_{U_{\beta}},$$

then $\exists g \in \mathcal{G}$ defined on U restricting to g_{α} on U_{α} . See Kobayashi-Nomizu [181], pp.1–2 for further discussion.

For example, if (G, X) is affine or projective geometry, the corresponding global object is an *affine structure* or *projective structure* on M . (Such structures are also called “affinely flat structures,” “flat affine structures,” “flat projective structures,” etc. An affine structure on a manifold is the same thing as a flat torsion-free affine connection, and a projective structure is the same thing as a flat normal projective connection (see Sharpe [249], Chern-Griffiths [64] Kobayashi [176] or Hermann [148] for the theory of projective connections). We shall refer to a projective structure modeled on $\mathbb{R}P^n$ as an $\mathbb{R}P^n$ -*structure*; a manifold with an $\mathbb{R}P^n$ -structure will be called an $\mathbb{R}P^n$ -*manifold*.

In many cases of interest, there may be a readily identifiable geometric object on X whose stabilizer is G , and modeling a manifold on (G, X) may be equivalent to a geometric object locally equivalent to the G -invariant geometric object on X . Perhaps the most important

such object is a locally homogeneous *Riemannian metric*. For example if X is a simply-connected Riemannian manifold of constant curvature K and G is its group of isometries, then locally modeling M on (G, X) is equivalent to giving M a Riemannian metric of curvature K . (This idea can be vastly extended, for example to cover indefinite metrics, locally homogeneous metrics whose curvature is not necessarily constant, etc.) In particular Riemannian metrics of constant curvature are special cases of (G, X) -structures on manifolds.

Thurston [266] gives a detailed discussion of some of the pseudogroups defining structures on 3-manifolds.

5.1. Geometric atlases

Let G be a Lie group acting transitively on a manifold X . Let $U \subset X$ be an open set and let $U \xrightarrow{f} X$ be a smooth map. We say that f is *locally- G* if for each component $U_i \subset U$, there exists $g_i \in G$ such that the restriction of g_i to $U_i \subset X$ equals the restriction of f to $U_i \subset U$. (Of course f will have to be a local diffeomorphism.) The collection of open subsets of X , together with locally- G maps defines a *pseudogroup* upon which can model structures on manifolds as follows.

A (G, X) -atlas on M is a pair (\mathcal{U}, Φ) where

$$\mathcal{U} := \{U_\alpha \mid \alpha \in A\},$$

is an open covering of M and

$$\Phi = \{U_\alpha \xrightarrow{\phi_\alpha} X\}_{U_\alpha \in \mathcal{U}}$$

is a collection of coordinate charts such that for each pair

$$(U_\alpha, U_\beta) \in \mathcal{U} \times \mathcal{U}$$

the restriction of $\phi_\alpha \circ (\phi_\beta)^{-1}$ to $\phi_\beta(U_\alpha \cap U_\beta)$ is locally- G . An (G, X) -structure on M is a maximal (G, X) -atlas and an (G, X) -manifold is a manifold together with an (G, X) -structure on it.

An (G, X) -manifold has an underlying real analytic structure, since the action of G on X is real analytic.

This notion of a map being *locally- G* has already been introduced for locally affine and locally projective maps.

Suppose that M and N are two (G, X) -manifolds and $M \xrightarrow{f} N$ is a map. Then f is an (G, X) -map if for each pair of charts

$$U_\alpha \xrightarrow{\phi_\alpha} X, \quad V_\beta \xrightarrow{\psi_\beta} X,$$

for M and N respectively, the restriction

$$\psi_\beta \circ f \circ \phi_\alpha^{-1} \Big|_{\phi_\alpha(U_\alpha \cap f^{-1}(V_\beta))}$$

is locally- G . In particular we only consider (G, X) -maps which are local diffeomorphisms. Clearly the set of (G, X) -automorphisms $M \rightarrow M$ forms a group, which we denote by $\mathbf{Aut}_{(G, X)}(M)$ or just $\mathbf{Aut}(M)$ when the context is clear.

EXERCISE 5.1.1. *Let N be an (G, X) -manifold and $M \xrightarrow{f} N$ a local diffeomorphism.*

- *There is a unique (G, X) -structure on M for which f is an (G, X) -map.*
- *Every covering space of an (G, X) -manifold has a canonical (G, X) -structure.*
- *Conversely suppose M is an (G, X) -manifold upon which a discrete subgroup $\Gamma \subset \mathbf{Aut}_{(G, X)}(M)$ acts properly and freely. Then M/Γ is an (G, X) -manifold and the quotient mapping*

$$M \rightarrow M/\Gamma$$

is a (G, X) -covering space.

5.1.1. The pseudogroup of local mappings. The fundamental example of an (G, X) -manifold is X itself. Evidently any open subset $\Omega \subset X$ has an (G, X) -structure (with only one chart—the inclusion $\Omega \hookrightarrow X$). Locally- G maps satisfy the following *Unique Extension Property*: If $U \subset X$ is a connected nonempty open subset, and $U \xrightarrow{f} X$ is locally- G , then there exists a unique element $g \in G$ restricting to f .

Here is another perspective on a (G, X) -atlas. First regard M as a quotient space of the disjoint union

$$\mathfrak{U} = \coprod_{\alpha \in A} U_\alpha$$

by the equivalence relation \sim defined by intersection of patches. A point $u \in U_\alpha \cap U_\beta$ determines corresponding elements

$$\begin{aligned} u_\alpha &\in U_\alpha \subset \mathfrak{U} \\ u_\beta &\in U_\beta \subset \mathfrak{U} \end{aligned}$$

and we define the equivalence relation on \mathfrak{U} by: $u_\alpha \sim u_\beta$.

Now the *coordinate change*

$$\phi_\beta(U_\alpha \cap U_\beta) \xrightarrow{\phi_\alpha \circ (\phi_\beta)^{-1}} \phi_\alpha(U_\alpha \cap U_\beta)$$

is locally- G . By the unique extension property, it agrees with the action of a unique element of G restricted to each connected component of $\phi_\beta(U_\alpha \cap U_\beta)$. Thus it corresponds to a *locally constant map*:

$$(23) \quad U_\alpha \cap U_\beta \xrightarrow{g_{\alpha\beta}} G$$

We can alternatively define the (G, X) -manifold M as the quotient of the disjoint union

$$\mathfrak{U}_\Phi := \coprod_{\alpha \in A} \phi_\alpha(U_\alpha)$$

by the equivalence relation \sim_Φ defined as:

$$\phi_\alpha(u_\alpha) \sim_\Phi g_{\alpha\beta}(\phi_\beta(u_\beta))$$

for $u \in U_\alpha \cap U_\beta$ notated as above. That \sim_Φ is an equivalence relation follows from the *cocycle identities*

$$(24) \quad \begin{aligned} g_{\alpha\alpha}(u_\alpha) &= 1 \\ g_{\alpha\beta}(u_\beta)g_{\beta\alpha}(u_\alpha) &= 1 \\ g_{\alpha\beta}(u_\beta)g_{\beta\gamma}(u_\gamma)g_{\gamma\alpha}(u_\alpha) &= 1 \end{aligned}$$

whenever $u \in U_\alpha$, $u \in U_\alpha \cap U_\beta$, $u \in U_\alpha \cap U_\beta \cap U_\gamma$, respectively.

This rigidity property is a distinguishing feature of the kind of geometric structures considered here. However, many familiar pseudogroup structures lack this kind of rigidity:

EXERCISE 5.1.2. *Show that the following pseudogroups do not satisfy the unique extension property:*

- C^r local diffeomorphisms between open subsets of \mathbb{R}^n , when $r = 0, 1, \dots, \infty, \omega$.
- Local biholomorphisms between open subsets of \mathbb{C}^n .
- Smooth diffeomorphisms between open subsets of a domain $\Omega \subset \mathbb{R}^n$ preserving an exterior differential form on Ω .

5.1.2. (G, X) -automorphisms. Now we discuss the *automorphisms* of a structure locally modeled on (G, X) .

If $\Omega \subset X$ is a domain, an (G, X) -automorphism $\Omega \xrightarrow{f} \Omega$ is the restriction of a unique element $g \in G$ preserving Ω , that is::

$$\text{Aut}_{(G, X)}(\Omega) \cong \text{Stab}_G(\Omega) = \{g \in G \mid g(\Omega) = \Omega\}$$

Now suppose that $M \xrightarrow{\phi} \Omega$ is a local diffeomorphism onto a domain $\Omega \subset X$. There is a homomorphism

$$\text{Aut}_{(G, X)}(M) \xrightarrow{\phi_*} \text{Aut}_{(G, X)}(\Omega)$$

whose kernel consists of all maps $M \xrightarrow{f} M$ making the diagram

$$\begin{array}{ccc} M & \xrightarrow{\phi} & \Omega \\ f \downarrow & & \parallel \\ M & \xrightarrow[\phi]{} & \Omega \end{array}$$

commute.

EXERCISE 5.1.3. *Find examples where:*

- ϕ_* is surjective but not injective;
- ϕ_* is injective but not surjective.

5.2. Development, holonomy

There is a useful globalization of the coordinate charts of a geometric structure in terms of the universal covering space and the fundamental group. The coordinate atlas $\{U_\alpha\}_{\alpha \in A}$ is replaced by a universal covering space $\widetilde{M} \rightarrow M$ with its group of deck transformations π . In the first approach, M is the quotient space of the disjoint union $\coprod_{\alpha \in A} U_\alpha$, and in the second it is the quotient space of \widetilde{M} by the group action π . The coordinate charts $U_\alpha \xrightarrow{\psi_\alpha} X$ are replaced by a globally defined map $\widetilde{M} \xrightarrow{\text{dev}} X$.

This process of development originated with Élie Cartan and generalizes the notion of a developable surface in \mathbf{E}^3 . If $S \hookrightarrow \mathbf{E}^3$ is an embedded surface of zero Gaussian curvature, then for each $p \in S$, the exponential map at p defines an isometry of a neighborhood of 0 in the tangent plane $\mathbb{T}_p S$, and corresponds to rolling the tangent plane $\mathbb{A}_p(S)$ on S without slipping. In particular every curve in S starting at p lifts to a curve in $\mathbb{T}_p S$ starting at $0 \in \mathbb{T}_p S$. For a Euclidean manifold, this globalizes to a local isometry of the universal covering $\widetilde{S} \rightarrow \mathbf{E}^2$, called by Élie Cartan the *development* of the surface (along the curve). The metric structure is actually subordinate to the affine connection, as this notion of development really only involves the construction of *parallel transport*.

Later this was incorporated into the notion of a *fiber space*, as discussed in the 1950 conference [263]. The collection of coordinate changes of a (G, X) -manifold M defines a fiber bundle $\mathcal{E}_M \rightarrow M$ with fiber X and structure group G . The fiber over $p \in M$ of the associated principal bundle

$$\mathfrak{P}_M \xrightarrow{\Pi_{\mathfrak{P}}} M$$

consists of all possible germs of (G, X) -coordinate charts at p . The fiber over $p \in M$ of \mathcal{E}_M consists of all possible *values* of (G, X) -coordinate charts at p . Assigning to the germ at p of a coordinate chart $U \xrightarrow{\psi} X$ its value

$$x = \psi(p) \in X$$

defines a mapping

$$(\mathfrak{P}_M)_p \longrightarrow (\mathcal{E}_M)_p.$$

Working in a local chart, the fiber over a point in $(\mathcal{E}_M)_p$ corresponding to $x \in X$ consists of all the different germs of coordinate charts ψ taking $p \in M$ to $x \in X$. This mapping identifies with the quotient mapping of the natural action of the stabilizer $\mathbf{Stab}(G, x) \subset G$ of $x \in X$ on the set of germs.

For Euclidean manifolds, $(\mathfrak{P}_M)_p$ consists of all *affine orthonormal frames*, that is, pairs (x, F) where $x \in \mathbf{E}^n$ is a point and F is an orthonormal basis of the tangent space $\mathbb{T}_x \mathbf{E}^n \cong \mathbb{R}^n$. For an affine manifold, $(\mathfrak{P}_M)_p$ consists of all *affine frames*: pairs (x, F) where now F is *any* basis of \mathbb{R}^n .

5.2.1. Construction of the developing map. Let M be an (G, X) -manifold. Choose a universal covering space

$$\tilde{M} \xrightarrow{\mathbf{\Pi}} M$$

and let $\pi = \pi_1(M)$ be the corresponding fundamental group. The covering projection $\mathbf{\Pi}$ induces an (G, X) -structure on \tilde{M} upon which π acts by (G, X) -automorphisms. The Unique Extension Property has the following important consequence.

PROPOSITION 5.2.1. *Let M be a simply connected (G, X) -manifold. Then there exists a (G, X) -map $M \xrightarrow{f} X$.*

It follows that the (G, X) -map f completely determines the (G, X) -structure on M , that is, the geometric structure on a simply-connected manifold is “pulled back” from the model space X . The (G, X) -map f is called a *developing map* for M and enjoys the following uniqueness property. If $M \xrightarrow{f'} X$ is another (G, X) -map, then there exists an (G, X) -automorphism ϕ of M and an element $g \in G$ such that

$$\begin{array}{ccc} M & \xrightarrow{f'} & X \\ \phi \downarrow & & \downarrow g \\ M & \xrightarrow{f} & X \end{array}$$




FIGURE 5.1. Extending a coordinate chart to a developing map

PROOF OF PROPOSITION 5.2.1. Choose a basepoint $x_0 \in M$ and a coordinate patch U_0 containing x_0 . For $x \in M$, we define $f(x)$ as follows. Choose a path $\{x_t\}_{0 \leq t \leq 1}$ in M from x_0 to $x = x_1$. Cover the path by coordinate patches U_i (where $i = 0, \dots, n$) such that $x_t \in U_i$ for $t \in (a_i, b_i)$ where

$$\begin{aligned} a_0 < 0 < a_1 < b_0 < a_2 < b_1 < a_3 < b_2 < \\ \dots < a_{n-1} < b_{n-2} < a_n < b_{n-1} < 1 < b_n \end{aligned}$$

Let $U_i \xrightarrow{\psi_i} X$ be an (G, X) -chart and let $g_i \in G$ be the unique transformation such that $g_i \circ \psi_i$ and ψ_{i-1} agree on the component of $U_i \cap U_{i-1}$ containing the curve $\{x_t\}_{a_i < t < b_{i-1}}$. Let

$$f(x) = g_1 g_2 \dots g_{n-1} g_n \psi_n(x);$$

we show that f is indeed well-defined. The map f does not change if the cover is refined. Suppose that a new coordinate patch U' is “inserted between” U_{i-1} and U_i . Let $\{x_t\}_{a' < t < b'}$ be the portion of the curve lying

inside U' so

$$a_{i-1} < a' < a_i < b_{i-1} < b' < b_i.$$

Let $U' \xrightarrow{\psi'} X$ be the corresponding coordinate chart and let $h_{i-1}, h_i \in G$ be the unique transformations such that ψ_{i-1} agrees with $h_{i-1} \circ \psi'$ on the component of $U' \cap U_{i-1}$ containing $\{x_t\}_{a' < t < b_{i-1}}$ and ψ' agrees with $h_i \circ \psi_i$ on the component of $U' \cap U_i$ containing $\{x_t\}_{a_i < t < b'}$. By the unique extension property $h_{i-1}h_i = g_i$ and it follows that the corresponding developing map

$$\begin{aligned} f(x) &= g_1 g_2 \cdots g_{i-1} h_{i-1} h_i g_{i+1} \cdots g_{n-1} g_n \psi_n(x) \\ &= g_1 g_2 \cdots g_{i-1} g_i g_{i+1} \cdots g_{n-1} g_n \psi_n(x) \end{aligned}$$

is unchanged. Thus the developing map as so defined is independent of the coordinate covering, since any two coordinate coverings have a common refinement.

Next we claim the developing map is independent of the choice of path. Since M is simply connected, any two paths from x_0 to x are homotopic. Every homotopy can be broken up into a succession of “small” homotopies, that is, homotopies such that there exists a partition

$$0 = c_0 < c_1 < \cdots < c_{m-1} < c_m = 1$$

such that during the course of the homotopy the segment $\{x_t\}_{c_i < t < c_{i+1}}$ lies in a coordinate patch. It follows that the expression defining $f(x)$ is unchanged during each of the small homotopies, and hence during the entire homotopy. Thus f is independent of the choice of path.

Since f is a composition of a coordinate chart with a transformation $X \rightarrow X$ coming from G , it follows that f is an (G, X) -map. The proof of Proposition 5.2.1 is complete. \square

If M is an arbitrary (G, X) -manifold, then we may apply Proposition 5.2.1 to a universal covering space \tilde{M} . We obtain the following basic result:

THEOREM 5.2.2 (Development Theorem). *Let M be an (G, X) -manifold with universal covering space $\tilde{M} \xrightarrow{\mathbf{\Pi}} M$ and group of deck transformations*

$$\pi = \pi_1(M) \subset \text{Aut}\left(\tilde{M} \xrightarrow{\mathbf{\Pi}} M\right)$$

Then \exists a pair (dev, h) consisting of an (G, X) -map $\widetilde{M} \xrightarrow{\text{dev}} X$ and a homomorphism $\pi \xrightarrow{h} G$ such that for each $\gamma \in \pi$,

$$\begin{array}{ccc} \widetilde{M} & \xrightarrow{\text{dev}} & X \\ \gamma \downarrow & & \downarrow h(\gamma) \\ \widetilde{M} & \xrightarrow{\text{dev}} & X \end{array}$$

commutes. Furthermore if (dev', h') is another such pair, then $\exists g \in G$ such that $\text{dev}' = g \circ \text{dev}$ and $h'(\gamma) = \text{Inn}(g) \circ h(\gamma)$ for all $\gamma \in \pi$. That is, the diagram

$$\begin{array}{ccccc} \widetilde{M} & \xrightarrow{\text{dev}} & X & \xrightarrow{g} & X \\ \gamma \downarrow & & \downarrow \text{hol}(\gamma) & & \downarrow \text{hol}'(\gamma) \\ \widetilde{M} & \xrightarrow{\text{dev}} & X & \xrightarrow{g} & X \end{array}$$

commutes.

We call such a pair (dev, h) a *development pair*, and the homomorphism h the *holonomy representation*. (It is the holonomy of a flat connection on a principal G -bundle over M associated to the (G, X) -structure.) The developing map globalizes the coordinate charts of the manifold and the holonomy representation globalizes the coordinate changes. In this generality the Development Theorem is due to C. Ehresmann [96] in 1936.

5.2.2. Role of the holonomy group. The image of the holonomy representation is the “smallest” subgroup $\Gamma \subset G$ such that M admits a (Γ, X) -structure:

EXERCISE 5.2.3. Let M be an (G, X) -manifold with development pair (dev, h) .

- Find a (G, X) -atlas for M such that the coordinate changes $g_{\alpha\beta}$ lie in Γ .
- Suppose that $N \rightarrow M$ is a covering space. Show that there exists a (G, X) -map $N \rightarrow X$ if and only if the holonomy representation restricted to $\pi_1(N) \hookrightarrow \pi_1(M)$ is trivial.

Thus the *holonomy covering space* $\widehat{M} \rightarrow M$ — the covering space of M corresponding to the kernel of h — is the “smallest” covering space of M for which a developing map is “defined.”

The holonomy group

$$\text{hol}(\pi) = \Gamma \subset G$$

is the “smallest” subgroup of G for which there is a compatible (G, X) -atlas, where the coordinate changes lie in Γ .

EXERCISE 5.2.4. *Let M be an (G, X) -manifold. Find a (G, X) -atlas such that all the coordinate changes are restrictions of transformations in Γ .*

EXERCISE 5.2.5. *Suppose that (G, X) and (G', X') represent a pair of geometries for which there exists a pair (Φ, ϕ) as in §5.2.3. Show that if M is a (G, X) -manifold with development pair (dev, h) , then*

$$(\Phi \circ \text{dev}, \phi \circ h)$$

is a development pair for the induced (X', G') -structure on M .

5.2.3. Extending geometries. A geometry may *contain* or *refine* another geometry. In this way one can pass from structures modeled on one geometry to structures modeled on a geometry containing it. Let (X, G) and (X', G') be homogeneous spaces and let $X \xrightarrow{\Phi} X'$ be a local diffeomorphism which is equivariant with respect to a homomorphism $\phi : G \rightarrow G'$ in the following sense: for each $g \in G$ the diagram

$$\begin{array}{ccc} X & \xrightarrow{\Phi} & X' \\ g \downarrow & & \downarrow \phi(g) \\ X & \xrightarrow[\phi]{} & X' \end{array}$$

commutes. Hence locally- G maps determine locally- (X', G') -maps and an (G, X) -structure on M induces an (X', G') -structure on M in the following way. Let $U_\alpha \xrightarrow{\psi_\alpha} X$ be an (G, X) -chart; the composition

$$U_\alpha \xrightarrow{\Phi \circ \psi_\alpha} X'$$

defines an (X', G') -chart.

EXERCISE 5.2.6. *Explain the extension of geometries in terms of the development pair.*

5.2.4. Simple applications of the developing map.

EXERCISE 5.2.7. *Suppose that M is a closed manifold with finite fundamental group.*

- *If X is noncompact then M admits no (G, X) -structure.*
- *If X is compact and simply-connected show that every (G, X) -manifold is (G, X) -isomorphic to a quotient of X by a finite subgroup of G .*

(Hint: if M and N are manifolds of the same dimension, $M \xrightarrow{f} N$ is a local diffeomorphism and M is closed, show that f must be a covering space.)

As a consequence a closed affine manifold must have infinite fundamental group and every $\mathbb{R}\mathbb{P}^n$ -manifold with finite fundamental group is a quotient of S^n by a finite group (and hence a spherical space form).

EXERCISE 5.2.8. Suppose $\Omega \subset X$ is a Γ -invariant open subset.

- $\text{dev}^{-1}(\Omega)$ is a π -invariant open subset of \widetilde{M} ;
- Its image

$$M_\Omega := \Pi(\text{dev}^{-1}(\Omega))$$

is an open subset of M ;

- Each connected component of $\Pi^{-1}(M_\Omega) \subset \widetilde{M}$ is a connected component of $\text{dev}^{-1}(\Omega)$.
- $M_\Omega \subset M$ depends only on the pair (Γ, Ω) and is independent of the choice of universal covering space $\widetilde{M} \rightarrow M$ and developing map $\widetilde{M} \xrightarrow{\text{dev}} X$.

This will be used later in §14.2.

5.3. The graph of a geometric structure

This can be put in an even “more global” context using the *fiber bundle associated to a (G, X) -structure*. This is a fiber bundle $\mathcal{E}_M \rightarrow M$ with fiber X , structure group G in the sense of Steenrod[259]. It plays a role analogous to the tangent bundle of a smooth manifold. It admits a flat structure, that is a foliation \mathcal{F} transverse to the fibration, and a section \mathcal{D}_M which is transverse to \mathcal{F} as well as the fibration. The section \mathcal{D}_M plays the role of the zero-section of the tangent bundle. Indeed, its normal bundle inside \mathcal{E}_M is isomorphic to the tangent bundle TM of M . It is obtained as the *graph* of the collection Φ of coordinate charts. The flat bundle \mathcal{E}_M is the natural “home” in which \mathcal{D}_M lives.

5.3.1. The tangent (G, X) -bundle. The total space \mathcal{E}_M of this bundle is obtained from the disjoint union

$$\mathfrak{U}_X := \coprod_{\alpha \in A} U_\alpha \times X$$

of trivial X -bundles.

Now suppose $U_\alpha, U_\beta \in \mathcal{U}$ are coordinate patches. Introduce an equivalence relation \sim_X on \mathfrak{U}_X by:

$$(u_\alpha, x) \sim_X (u_\beta, g_{\alpha\beta}(u_\beta)x)$$

where $g_{\alpha\beta}$ is the cocycle introduced in (23). The cocycle identities (24) imply that \sim_X is an equivalence relation. The projections

$$U_\alpha \times X \longrightarrow U_\alpha$$

are trivial X -bundles and define a trivial X -bundle

$$\mathfrak{U}_X \longrightarrow \mathfrak{U}$$

compatible with the equivalence relations \sim_X, \sim . The corresponding mapping of quotient spaces

$$\begin{array}{ccc} \mathcal{E}_M & \xlongequal{\quad} & \mathfrak{U}_X / \sim_X \\ & & \downarrow \Pi \\ M & \xlongequal{\quad} & \mathfrak{U} / \sim \end{array}$$

is a locally trivial X -bundle with structure group G .

Furthermore the structure group is really G with the discrete topology, since the transition functions

$$U_\alpha \cap U_\beta \xrightarrow{g_{\alpha\beta}} G$$

are locally constant. This implies that the foliation of the total space \mathcal{E}_M with local leaves (sometimes called *plaques*) $U_\alpha \times \{x\}$ piece together to define the leaves of a foliation \mathcal{F} of \mathcal{E}_M . (Compare Steenrod [259, §13].)

EXERCISE 5.3.1.

- Show that for every leaf $L \subset \mathcal{E}_M$ of \mathcal{F} , the restriction $\Pi|_L$ is a covering space $L \longrightarrow M$.
- If M is simply connected, then $(\mathcal{E}_M, \mathcal{F})$ is trivial, that is, isomorphic to $M \times X$ with the trivial foliation, namely the one with leaves $M \times \{x\}$, where $x \in X$.

It follows that the flat (G, X) -bundle $(\mathcal{E}_M, \mathcal{F})$ arises from a representation $\pi_1(M) \xrightarrow{h} G$ as follows. The group $\pi_1(M)$ admits a (left-)action on the trivial bundle $\widetilde{M} \times X$ by:

$$(\tilde{p}, x) \mapsto (\tilde{p}\gamma^{-1}, h(\gamma)x)$$

where

$$\begin{aligned} \widetilde{M} \times \pi_1(M) &\longrightarrow \widetilde{M} \\ (\tilde{p}, \gamma) &\longmapsto \tilde{p}\gamma \end{aligned}$$

denotes the (right-) action of $\pi_1(M)$ by deck transformations. Then \mathcal{E}_M identifies as the quotient $(\widetilde{M} \times X) / \pi_1(M)$, that is as the *fiber product*




FIGURE 5.2. The graph of a developing map is an \mathcal{F} -transverse section

$\widetilde{M} \times_{\mathfrak{h}} X$. Furthermore \mathfrak{h} is unique up to the action of $\text{Inn}(G)$ by left-composition. We call $\mathfrak{h} \in \text{Hom}(\pi_1(M), G)$ the *holonomy representation* of the flat (G, X) -bundle $(\mathcal{E}_M, \mathcal{F})$.

5.3.2. Developing sections. Just as $(\mathcal{E}_M, \mathcal{F})$ globalizes the coordinate changes, its *transverse section* \mathcal{D}_M globalizes the coordinate atlas Φ .

When M is a single coordinate patch, then \mathcal{E}_M is just the product $M \times X$ and $\mathcal{E}_M \rightarrow M$ is just the Cartesian projection $M \times X \rightarrow M$. A section of $\mathcal{E}_M \rightarrow M$ is just the graph of a map $M \xrightarrow{f} X$:

$$\begin{aligned} M &\xrightarrow{\text{graph}(f)} M \times X \cong \mathcal{E}_M \\ p &\longmapsto (p, f(p)) \end{aligned}$$

5.3.3. The associated principal bundle. The fiber over $p \in M$ of the associated principal bundle

$$\mathfrak{P}_M \xrightarrow{\Pi_{\mathfrak{P}}} M$$

consists of all possible germs of (G, X) -coordinate charts at p . The fiber over $p \in M$ of \mathcal{E}_M consists of all possible *values* of (G, X) -coordinate




FIGURE 5.3. The isotopy between nearby \mathcal{F} -transverse sections

charts at p . Assigning to the germ at p of a coordinate chart $U \xrightarrow{\psi} X$ its value

$$x = \psi(p) \in X$$

defines a mapping

$$(\mathfrak{P}_M)_p \longrightarrow (\mathcal{E}_M)_p.$$

Working in a local chart, the fiber over a point in $(\mathcal{E}_M)_p$ corresponding to $x \in X$ consists of all the different germs of coordinate charts ψ taking $p \in M$ to $x \in X$. This mapping identifies with the quotient mapping of the natural action of the stabilizer $\mathbf{Stab}(G, x) \subset G$ of $x \in X$ on the set of germs.

For Euclidean manifolds, $(\mathfrak{P}_M)_p$ consists of all *affine orthonormal frames*, that is, pairs (x, F) where $x \in \mathbf{E}^n$ is a point and F is an orthonormal basis of the tangent space $\mathbf{T}_x \mathbf{E}^n \cong \mathbb{R}^n$. For an affine manifold, $(\mathfrak{P}_M)_p$ consists of all *affine frames*: pairs (x, F) where now F is *any* basis of \mathbb{R}^n .

The coordinate atlas/developing map defines a section of $\mathcal{E}_M \rightarrow M$ which is transverse to the two complementary foliations of \mathcal{E}_M :

- As a section, it is necessarily transverse to the foliation of \mathcal{E}_M by fibers;
- The nonsingularity of the coordinate charts/developing map implies this section is transverse to the horizontal foliation \mathcal{F}_M of \mathcal{E}_M defining the flat structure.

This picture of an Ehresmann structure will be used in defining the deformation space $\text{Def}_{(G,X)}(\Sigma)$ in Chapter 7, §7.2.

Figure 5.4, Figure 5.5, and Figure 5.6 depict developing sections for various $\mathbb{R}\mathbb{P}^1$ -manifolds. M and X are both homeomorphic to S^1 , and we represent M as a horizontal closed interval with endpoints identified. Similarly X is represented as a vertical closed interval with endpoints identified. Thus the total space \mathcal{E}_M is represented by a square, where the left and right edges are identified by parallel (horizontal) translation. The projection Π is just horizontal projection, with fibers are vertical line segments. The leaves of \mathcal{F}_M of \mathcal{E}_M are drawn so that they are identified by the parallel translation.

Figure 5.4 and Figure 5.5) depict structures with trivial holonomy. The leaves, represented by horizontal lines (lines of slope 0), are all closed sections, corresponding to the singular structure with “constant developing map”. Figure 5.4 depicts the canonical structure $\mathbb{R}\mathbb{P}^1$; the developing section is the line of slope 1, the graph of the identity map $\mathbb{R}\mathbb{P}^1 \rightarrow \mathbb{R}\mathbb{P}^1$.

For any $m \in \mathbb{Z}$ a line segment of slope m (and some of its horizontal translates) describes a section s . We have already discussed the cases $m = 0, 1$. If $m \neq 0$, the section is transverse to \mathcal{F} . Replacing m by $-m$ gives a section inducing the opposite orientation, so that the line of slope -1 (the other diagonal of the square) depicts the developing section for an oppositely oriented manifold; explicitly, the developing section is the graph of a reflection of $\mathbb{R}\mathbb{P}^1$ (an involution which reverses orientation).

Figure 5.5) depicts the developing section for the double covering of $\mathbb{R}\mathbb{P}^1$.

Figure 5.6 depicts a structure with elliptic holonomy. In this case the foliation is a linear foliation of the torus. In this case the leaves are drawn as lines of positive slope (explicitly $m = 1/3$ and each leaf projects to M by a triple covering. The “diagonal” section s_1 and the “horizontal” section s_2 are both \mathcal{F} -transverse and define $\mathbb{R}\mathbb{P}^1$ -structures.

Figure 5.7 depicts structures with hyperbolic holonomy η . The two fixed points of η on $\mathbb{R}\mathbb{P}^1$ determine two “horizontal” sections (“constant developing maps” of singular structures). In the picture, these are

represented by the top/bottom edges of the square and the diameter halfway up. The depicted section s_1 is horizontal and misses these two horizontal sections; the corresponding developing map misses $\text{Fix}(\eta)$ and corresponds to the (Hopf) affine structure. The depicted section s_2 crosses both constant horizontal sections, and the corresponding developing map is onto.

Figure 5.8 is similar, except now the holonomy η is parabolic. Corresponding to the single fixed point of η is a horizontal closed leaf, represented in this picture as the top/bottom sides of the square. The complete Euclidean structure is represented by the \mathcal{F} -transverse horizontal section s_1 and misses this "constant" section. The section s_2 is \mathcal{F} -transverse and corresponds to a structure with surjective developing map.




FIGURE 5.4. Developing section for the canonical projective closed 1-manifold \mathbb{RP}^1




FIGURE 5.5. Developing section for other projective closed 1-manifolds with trivial holonomy




FIGURE 5.6. Developing sections for projective closed 1-manifolds with elliptic holonomy




FIGURE 5.7. Developing sections for projective closed 1-manifolds with hyperbolic holonomy




FIGURE 5.8. Developing sections for projective closed 1-manifolds with parabolic holonomy

5.4. The classification of geometric 1-manifolds

The basic general question concerning geometric structures on manifolds is, given a topological manifold M and a geometry (G, X) , whether an (G, X) -structure on M exists, and if so, to classify all (G, X) -structures on M . Ideally, one would like a *deformation space*, a topological space whose points correspond to isomorphism classes of (G, X) -manifolds.

As an exercise to illustrate these general ideas, we classify geometric manifolds in dimension one. We consider the three geometries

$$\mathbb{E}^1 \xrightarrow{\cong} \mathbb{A}^1 \hookrightarrow \mathbb{P}^1$$

in increasing order. Euclidean manifolds are affine manifolds, which in turn are projective manifolds. Thus we classify $\mathbb{R}\mathbb{P}^1$ -manifolds. (Compare Kuiper [190], Goldman [115], Baues [23].)

Let M be a connected 1-manifold. There are two cases:

- M is noncompact, in which case M is homeomorphic (diffeomorphic) to a line ($M \approx \mathbb{R}$);
- M is compact, in which case M homeomorphic (diffeomorphic) to a circle ($M \approx S^1$).

In particular M is simply connected $\iff M$ is noncompact and otherwise $\pi_1(M) \cong \mathbb{Z}$.

5.4.1. Compact Euclidean 1-manifolds and flat tori. The cyclic group \mathbb{Z} acts by translations on $\mathbb{E}^1 \cong \mathbb{R}$. The quotient

$$E_1 := \mathbb{E}^1/\mathbb{Z} \cong \mathbb{R}/\mathbb{Z}$$

is a compact Euclidean 1-manifold. The Euclidean metric on \mathbb{R} induces a flat Riemannian structure on the quotient \mathbb{R}/\mathbb{Z} which has length 1.

More generally, choose $\ell > 0$. Then the quotient

$$E_\ell := \mathbb{E}^1/\ell\mathbb{Z} \cong \mathbb{R}/\ell\mathbb{Z}$$

is a compact Euclidean 1-manifold which has length ℓ . Different choices of ℓ determine different isometry classes of Euclidean 1-manifolds but E_1 is *affinely isomorphic* to E_ℓ by the affine map $x \mapsto \ell x$. In other words, if $\ell \neq 1$, then E_1 and E_ℓ are inequivalent Euclidean manifolds but equivalent affine manifolds.

EXERCISE 5.4.1. *Let $M = E_\ell$. Show that the total space of \mathcal{E}_M identifies with the quotient of \mathbb{R}^2 by the diagonally embedded \mathbb{Z} acting by translations:*

$$(x, y) \mapsto (x + n, y + n\ell)$$

for $n \in \mathbb{Z}$, the fibration is induced by the projection

$$(x, y) \mapsto x,$$

the foliation induced by horizontal lines $\mathbb{R} \times \{y\}$, and the developing section by the diagonal

$$\Delta(x) := (x, x).$$

When these structures are regarded as $\mathbb{R}\mathbb{P}^1$ -manifolds, \mathcal{E}_M acquires an extra (horizontal) closed leaf. This section (corresponding to the ideal point of $\mathbb{R}\mathbb{P}^1$) is disjoint from the developing section Δ .

These manifolds generalize to one of the most basic classes of closed geometric manifolds, namely the *flat tori*. Let $\Lambda \subset \mathbb{R}^n$ be a *lattice*, that is the additive subgroup of \mathbb{R}^n generated by a basis. Then Λ acts by translations, so the quotient \mathbb{R}^n/Λ is a compact Euclidean manifold. Bieberbach proved that *every* compact Euclidean manifold is finitely covered by a flat torus.

EXERCISE 5.4.2. *Since Λ is a normal subgroup of \mathbb{R}^n , a flat torus is also an abelian Lie group. Show that this algebraic structure is compatible with the geometric structure: the Euclidean structure on \mathbb{R}^n/Λ is invariant under multiplications. (Since \mathbb{R}^n is commutative, left-multiplications and right-multiplications coincide.)*

5.4.2. Compact affine 1-manifolds and Hopf circles. A compact affine manifold is either a Euclidean manifold as above, or given by the following construction. Let $\lambda > 1$ and consider the cyclic group $\langle \lambda \rangle \cong \mathbb{Z}$ acting by homotheties on \mathbf{A}^1 :

$$x \mapsto \lambda^n x$$

Then

$$A_\lambda := \mathbb{R}^+ / \langle \lambda \rangle$$

is a compact affine 1-manifold.

EXERCISE 5.4.3. *Show that different values of λ yield inequivalent affine structures, and no A_λ is affinely equivalent to E_ℓ . However show that, for λ, λ' the developing maps for A_λ and $A_{\lambda'}$ are topologically conjugate by a homeomorphism $\mathbf{A}^1 \rightarrow \mathbf{A}^1$ and the developing maps for A_λ and E_ℓ are topologically semi-conjugate by a homeomorphism $\mathbb{R}^+ \rightarrow \mathbb{R} \cong \mathbf{E}^1$.*

We call these latter affine 1-manifolds *Hopf circles*, since these are the 1-dimensional cases of *Hopf manifolds* discussed in §6.4.

5.4.2.1. *Geodesics.* Hopf circles model incomplete closed geodesics on affine manifolds. The affine parameter on a Hopf circle is paradoxical. A particle moving with zero acceleration seems to be accelerating so rapidly that in finite time it “runs off the edge of the manifold.” Here is an explicit calculation:

The geodesic on \mathbf{A}^1 defined by

$$t \mapsto 1 + t(\lambda^{-1} - 1)$$

begins at 1 and in time

$$t_\infty := 1 + \lambda^{-1} + \lambda^{-2} + \dots = (1 - \lambda^{-1})^{-1} > 0$$

reaches 0. It defines a closed incomplete closed geodesic $p(t)$ on M starting at $p(0) = p_0$. The lift

$$(-\infty, t_\infty) \xrightarrow{\tilde{p}} \widetilde{M}$$

satisfies

$$\text{dev}(\tilde{p}(t)) = 1 + t(\lambda^{-1} - 1),$$

which uniquely specifies the geodesic $p(t)$ on M . It is a geodesic since its velocity

$$p'(t) = (\lambda^{-1} - 1)\partial_x$$

is constant (parallel). However $p(t_n) = p_0$ for

$$t_n := \frac{1 - \lambda^{-n}}{1 - \lambda^{-1}} = 1 + \lambda^{-1} + \dots + \lambda^{1-n}$$

and as viewed in M , seems to go “faster and faster” through each cycle. By time $t_\infty = \lim_{n \rightarrow \infty} t_n$, it seems to “run off the manifold:” the geodesic is only defined for $t < t_\infty$. The apparent paradox is that $p(t)$ has *zero acceleration*: it would have “constant speed” if “speed” were only defined.

EXERCISE 5.4.4. *Show that these affine structures are invariant affine structures on the Lie group S^1 , namely, that the translation on the group S^1 is affine. (Since S^1 is abelian, both left- and right-translation agree.)*

These are the only examples of compact affine 1-manifolds, although there are projective manifolds which have the “same” holonomy homomorphisms, defined by *grafting*; see §5.4.5.

5.4.3. Classification of projective 1-manifolds. To simplify matters, we pass to the universal covering $X = \widetilde{\mathbb{R}P^1}$, which is homeomorphic to \mathbb{R} and the corresponding covering group $G = \widetilde{\mathrm{PGL}(2, \mathbb{R})}$ which acts on X . Suppose that M is a connected noncompact $\mathbb{R}P^1$ -manifold (and thus diffeomorphic to an open interval). Then a developing map

$$M \approx \mathbb{R} \xrightarrow{\mathrm{dev}} \mathbb{R} \approx X$$

is necessarily an embedding of M onto an open interval in X . Given two such embeddings

$$M \xrightarrow{f} X, \quad M \xrightarrow{f'} X$$

whose images are equal, then $f' = j \circ f$ for a diffeomorphism $M \xrightarrow{f} M$. Thus two $\mathbb{R}P^1$ -structures on M which have equal developing images are isomorphic. Thus the classification of $\mathbb{R}P^1$ -structures on M is reduced to the classification of G -equivalence classes of intervals $J \subset X$. Choose a diffeomorphism

$$X \approx \mathbb{R} \approx (-\infty, \infty);$$

an interval in X is determined by its pair of endpoints in $[-\infty, \infty]$. Since G acts transitively on X , an interval J is either bounded in X or projectively equivalent to X itself or one component of the complement of a point in X . Suppose that J is bounded. Then either the endpoints of J project to the same point in $\mathbb{R}P^1$ or to different points. In the first case, let $N > 0$ denote the degree of the map

$$J/\partial J \longrightarrow \mathbb{R}P^1$$

induced by dev ; in the latter case choose an interval J^+ such that the restriction of the covering projection $X \longrightarrow \mathbb{R}P^1$ to J^+ is injective

and the union $J \cup J^+$ is an interval in X whose endpoints project to the same point in $\mathbb{R}P^1$. Let $N > 0$ denote the degree of the restriction of the covering projection to $J \cup J^+$. Since G acts transitively on pairs of distinct points in $\mathbb{R}P^1$, it follows easily that bounded intervals in X are determined up to equivalence by G by the two discrete invariants: whether the endpoints project to the same point in $\mathbb{R}P^1$ and the positive integer N . It follows that every (G, X) -structure on M is (G, X) -equivalent to one of the following types. We shall identify X with the real line and group of deck transformations of $X \rightarrow \mathbb{R}P^1$ with the group of integer translations.

- A complete (G, X) -manifold (that is, $M \xrightarrow{\text{dev}} X$ is a diffeomorphism);
- $M \xrightarrow{\text{dev}} X$ is a diffeomorphism onto one of two components of the complement of a point in X , for example, $\mathbb{R}^+ = (0, \infty)$.
- dev is a diffeomorphism onto an interval $(0, N)$ where $N > 0$ is a positive integer;
- dev is a diffeomorphism onto an interval $(0, N + \frac{1}{2})$.

Next consider the case that M is a compact 1-manifold; choose a basepoint $x_0 \in M$. Let

$$\pi = \pi_1(M, x_0) \cong \mathbb{Z}$$

be the corresponding fundamental group of M and let $\gamma \in \pi$ be a generator. We claim that the conjugacy class of $h(\gamma) \in G$ completely determines the structure. Choose a lift J of $M \setminus \{x_0\}$ to \widetilde{M} to serve as a fundamental domain for π . Then J is an open interval in \widetilde{M} with endpoints y_0 and y_1 . After choosing a developing map $\widetilde{M} \xrightarrow{\text{dev}} X$, a holonomy representation $\pi \xrightarrow{h} G$,

$$\text{dev}(y_1) = h(\gamma)\text{dev}(y_0).$$

Now suppose that dev' is a developing map for another structure with the same holonomy. By applying an element of G we may assume that $\text{dev}(y_0) = \text{dev}'(y_0)$ and that $\text{dev}(y_1) = \text{dev}'(y_1)$. Furthermore a diffeomorphism $J \xrightarrow{\phi} J$ exists such that

$$\text{dev}' = \phi \circ \text{dev}.$$

This diffeomorphism lifts to a diffeomorphism $\widetilde{M} \xrightarrow{\tilde{\phi}} \widetilde{M}$ taking dev to dev' . Conversely suppose that $\eta \in G$ is orientation-preserving (this means simply that η lies in the identity component of G) and is not the identity. Then $\exists x_0 \in X$ which is not fixed by η ; let $x_1 = \eta x_0$. There exists a diffeomorphism $J \rightarrow X$ taking the endpoints y_i of J

to x_i for $i = 0, 1$. This diffeomorphism extends to a developing map $\widetilde{M} \xrightarrow{\text{dev}} X$. In summary:

THEOREM 5.4.5. *A compact \mathbb{RP}^1 -manifold is either projectively equivalent to:*

- *A Hopf circle $\mathbb{R}^+ / \langle \lambda \rangle$;*
- *A Euclidean 1-manifold \mathbb{R}/\mathbb{Z} ;*
- *A quotient of the universal covering of \mathbb{RP}^1 by a cyclic group.*

The first two cases are the affine 1-manifolds, and are homogeneous. The last case contains homogeneous structures if the holonomy is elliptic.

EXERCISE 5.4.6. *Determine all automorphisms of each of the above list of \mathbb{RP}^1 -manifolds.*

COROLLARY 5.4.7. *Let G^0 denote the identity component of the universal covering group G of $\text{PGL}(2, \mathbb{R})$. Let M be a closed 1-manifold. Then the set of isomorphism classes of \mathbb{RP}^1 -structures on M is in bijective correspondence with the set*

$$(G^0 \setminus \{1\}) / \text{Inn}(G)$$

of G -conjugacy classes in the set $G^0 \setminus \{1\}$ of elements of G^0 not equal to the identity.

EXERCISE 5.4.8. *Show that the quotient topology on $(G^0 \setminus \{1\}) / \text{Inn}(G)$ is not Hausdorff.*

5.4.4. Homogeneous affine structures. As observed in Exercise 5.4.4, a closed one-dimensional affine manifold M has the extra structure as an *affine Lie group*: M is a Lie group isomorphic to the circle \mathbb{R}/\mathbb{Z} and the operations of left-translation and right-translation are affine. (Since M is abelian, these two operations are identical.) In particular the universal covering \widetilde{M} inherits an affine Lie group structure (isomorphic to \mathbb{R}). By forming products one obtains affine Lie group structures on the two-dimensional abelian Lie group \mathbb{R}^2 .

EXERCISE 5.4.9. *Affine Lie group structures on \mathbb{R}^2 .*

- *Show that the products of affine Lie group structures on \mathbb{R} give three nonequivalent affine Lie group structures on $G = \mathbb{R}^2$. If $\Lambda < G$ is a lattice, then G/Λ is an affine Lie group isomorphic to the 2-torus $\mathbb{R}^2/\mathbb{Z}^2$. Find such a structure which is not affinely equivalent to a product of closed affine 1-manifolds.*
- *Find two other affine Lie group structures on G .*
- *Prove that these five structures are the only affine Lie group structures on G .*

In § 10.2, these structures will be identified with 2-dimensional commutative associative algebras over \mathbb{R} and will be generalized to left-invariant affine structures on (possibly noncommutative) Lie groups.

Every *homogeneous* affine structure on T^2 is obtained by this construction. The other affine structures are obtained by the radiant suspension construction of Exercise 6.5.10; compare Baues [23] for more information on the affine structures on T^2 .





FIGURE 5.9. Some incomplete complex-affine structures on T^2

FIGURE 5.10. Some hyperbolic affine structures on T^2




FIGURE 5.11. Radiant affine structures on T^2 developing to a halfplane




FIGURE 5.12. Nonradiant affine structures on T^2 developing to a halfplane

5.4.5. Grafting. Another approach to the classification is through the operation of *grafting*, developed in Goldman [120] in this generality. Let M_1, M_2 be two (G, X) -manifolds with two-sided hypersurfaces $V_i \subset M_i$ respectively. Suppose that each V_i has a tubular neighborhood U_i and with an (G, X) -isomorphism $U_1 \xrightarrow{f} U_2$. Then the complement $M_i \setminus V_i$ is the interior of a manifold-with-boundary $M_i|V_i$ with two boundary components V_i', V_i'' and an identification map $M_i|V_i \rightarrow M_i$ which identify $V_i' \longleftrightarrow V_i''$ to V_i

EXERCISE 5.4.10. *The restriction of the isomorphism f to $V_i \subset M_i$ induces identifications $V_1' \longleftrightarrow V_2''$ and $V_2' \longleftrightarrow V_1''$ which defines an equivalence relation \sim on the disjoint union $M_1|V_1 \sqcup M_2|V_2$. Then the quotient space*

$$M := \left(M_1|V_1 \sqcup M_2|V_2 \right) / \sim$$

inherits a unique (G, X) -structure such that the natural inclusions $M_i \setminus V_i \hookrightarrow M$ are (G, X) -maps.

This construction applies in dimension one, to give all compact \mathbb{RP}^1 -manifolds.

EXERCISE 5.4.11. *If M is a closed \mathbb{RP}^1 -manifold with hyperbolic or parabolic holonomy, the following conditions are equivalent:*

- *dev is surjective;*
- *M is not homogeneous;*
- *The developing image $\text{dev}(\widetilde{M})$ contains at least one fixed point of the holonomy;*
- *M is obtained by grafting a homogeneous (affine) 1-manifold with the model \mathbb{RP}^1 -manifold M_0 (given by an isomorphism $M_0 \cong \mathbb{RP}^1$).*

CHAPTER 6

Examples of Geometric Structures

This section introduces examples of geometric manifolds in dimensions greater than one. The theory of Lie groups and their homogeneous spaces organized the abundance of classical geometries, and this *algebraicization of geometry* clarifies the relationship between various geometric structures. We give several general constructions to pass from one geometric structure to another. This provides a rich class of geometric structures on manifolds.

We begin with general remarks on these constructions, which include the inclusion of homogeneous subdomains, Cartesian products, mapping tori and homogeneous fibrations. Then we study parallel structures in affine geometry, generalizing the construction of Euclidean geometry as (flat) Riemannian geometry. From our viewpoint, a Euclidean structure is just a *parallel Riemannian structure* on an affine manifold. This is the first example of *extending* a geometry, where the model space X is fixed (in this case an affine space) but the automorphism group G is reduced or enlarged. We digress to discuss Bieberbach's theorem structure of Euclidean manifolds, (see Charlap [63] or Wolf [286]), and give some examples of closed Euclidean manifolds, some of which are mapping tori constructed as *parallel suspensions*.

We then discuss other important cases, arising when one model space X' embeds in the other model space as a subdomain:

- The inclusion of affine space in projective space; in this way every affine structure inherits a projective structure.
- The (real-) projective models of hyperbolic geometry and elliptic geometry, whereby every elliptic or hyperbolic manifold has an $\mathbb{R}P^n$ -structure.
- The (complex-) projective models of hyperbolic geometry and elliptic geometry; in real dimension 2, every elliptic or hyperbolic surface has a natural $\mathbb{C}P^1$ -structure.
- The complement $A^n \setminus \{p\}$ of a point p in affine n -space naturally identifies with the complement $\mathbb{R}^n \setminus \{\mathbf{0}\}$ which has automorphism group $\mathrm{GL}(n+1, \mathbb{R})$ and every $(\mathrm{GL}(n+1, \mathbb{R}), \mathbb{R}^n \setminus \{\mathbf{0}\})$ -manifold has a special type of affine structure (called *radiant*).

Hopf manifolds — quotients of $\mathbb{A}^n \setminus \{p\}$ by cyclic groups are basic examples of radiant affine manifolds, closely related to $\mathbb{R}\mathbb{P}^{n-1}$ -manifolds.

Properties of radiant affine manifolds are discussed, and we use the *radiant suspension* construction to produce affine structures on products $\Sigma \times S^1$, for any surface Σ .

6.1. The hierarchy of geometries

We begin by stating the general construction for enlarging one geometry to another. Cartesian products are closely related and we describe taking products of affine and projective structures. These are special cases of *homogeneous fibrations*. The general discussion ends with a brief review of the mapping torus construction, and how suspensions of affine and projective automorphisms lead to new structures.

6.1.1. Enlarging and refining. Here is the general construction. Suppose that $X \xrightarrow{\Phi} X'$ is a universal covering space and G is the group of lifts of transformations $X' \xrightarrow{g'} X'$ in G' to X . Let $G \xrightarrow{\phi} G'$ be the corresponding homomorphism.

EXERCISE 6.1.1. *Show that (Φ, ϕ) induces an isomorphism between the categories of (G, X) -manifolds/maps and (X', G') -manifolds/maps.*

For this reason we may always assume (when convenient) that our model space X is simply-connected.

In many cases, we wish to consider maps between different manifolds with geometric structures modeled on different geometries. To this end we consider the following general situation. Let (G, X) and (X', G') be two homogeneous spaces representing different geometries and consider a family \mathfrak{M} of maps $X \rightarrow X'$ such that if $f \in \mathfrak{M}, g \in G, g' \in G'$, then the composition

$$g' \circ f \circ g \in \mathfrak{M}.$$

If $U \subset X$ is a domain, a map $U \xrightarrow{f} X'$ is *locally- \mathfrak{M}* if for each component $U_i \subset U$ there exists $f_i \in \mathfrak{M}$ such that the restriction of f to $U_i \subset U$ equals the restriction of f_i to $U_i \subset X$. Let M be an (G, X) -manifold and N an (X', G') -manifold. Suppose that $f : M \rightarrow N$ is a smooth map. We say that f is an *\mathfrak{M} -map* if for each pair of charts

$$\begin{aligned} U_\alpha &\xrightarrow{\phi_\alpha} X \text{ (for } M) \\ V_\beta &\xrightarrow{\psi_\beta} X' \text{ (for } N) \end{aligned}$$

the restriction of the composition $\psi_\beta \circ f \circ \phi_\alpha^{-1}$ to $\phi_\alpha(U_\alpha \cap f^{-1}(V_\beta))$ is locally- \mathfrak{M} .

The basic examples are affine and projective maps between affine and projective manifolds: For affine maps we take

$$(X, G) = (\mathbb{R}^m, \text{Aff}(\mathbb{R}^m)), \quad (X', G') = (\mathbb{R}^n, \text{Aff}(\mathbb{R}^n)), \\ \mathfrak{M} = \text{aff}(\mathbb{R}^m, \mathbb{R}^n).$$

For example if M, N are affine manifolds, and $M \times N$ is the product affine manifold (see §4.17), then the projections $M \times N \rightarrow M$ and $M \times N \rightarrow N$ are affine. Similarly if $x \in M$ and $y \in N$, the inclusions

$$\{x\} \times N \hookrightarrow M \times N, \\ M \times \{y\} \hookrightarrow M \times N$$

are each affine.

For projective maps we take

$$(X, G) = (\mathbb{P}^m, \text{Proj}(\mathbb{P}^m)), \\ (X', G') = (\mathbb{P}^n, \text{Proj}(\mathbb{P}^n)),$$

and $\mathfrak{M} = \text{Proj}(\mathbb{P}^m, \mathbb{P}^n)$, the set of projective maps $\mathbb{P}^m \rightarrow \mathbb{P}^n$ (or more generally the collection of locally projective maps defined on open subsets of \mathbb{P}^m). Important examples of this correspondence abound, many of which occur when Φ is an embedding. For example when Φ is the identity map and $G \subset G'$ is a subgroup, then every (G, X) -structure is a fortiori an (X', G') -structure. Thus every Euclidean structure is a similarity structure which in turn is an affine structure. Similarly every affine structure determines a projective structure, using the embedding

$$(\mathbb{R}^n, \text{Aff}(\mathbb{R}^n)) \hookrightarrow (\mathbb{P}^n, \text{Proj}(\mathbb{P}^n))$$

of affine geometry in projective geometry.

The polarities discussed in § 3.2.3 provide further examples. For example, elliptic-geometry structures arise as projective structures whose holonomy preserve an elliptic polarity — these identify with Riemannian structures of constant positive curvature. Hyperbolic structures arise from projective structures whose holonomy preserves a hyperbolic polarity of index 1. In each case these correspond to the respective subgroups $\text{PO}(n+1)$ and $\text{PO}(n, 1)$ of $\text{PGL}(n+1, \mathbb{R})$. *Contact projective structures* arise on manifolds M^{2m+1} when the holonomy preserves a null polarity, as in Exercise 3.2.8; these structure correspond to the subgroup $\text{PSp}(2d, \mathbb{R}) < \text{PGL}(2d+1)$.

EXERCISE 6.1.2. *A null polarity θ on \mathbb{P} defines a contact structure ξ on \mathbb{P} , namely the contact hyperplane $\xi_p \subset T_p\mathbb{P}$ at $p \in \mathbb{P}$ is just*

the tangent plane $T_p\theta(p)$. Conversely, the polar hyperplane $\theta(p)$ is the unique projective hyperplane in \mathbf{P} whose tangent space at p equals ξ_p .

A manifold modeled on the projective space with a contact structure arising from a null polarity may be called a *contact projective manifold*. Gray's stability theorem (see for example Eliashberg-Mishachev [98], §9.5.2, .95) asserts that contact structures fall into a discrete set of isotopy classes. Perhaps a more appropriate classification problem involves fixing a closed contact manifold (N^{2d+1}, ξ) and classifying the marked contact $\mathbb{R}\mathbf{P}^{2d+1}$ -structures compatible with ξ .

6.1.2. Cartesian products. The following is due to Benzécri [35].

EXERCISE 6.1.3 (Products of affine manifolds). *Let M^m, N^n be affine manifolds.*

- (1) *Show that the Cartesian product $M^m \times N^n$ has a natural affine structure.*
- (2) *Show that $M \times N$ is complete if and only if both M and N are complete.*
- (3) *Show that $M \times N$ is radiant if and only if both M and N are radiant.*

For projective structures, the situation is somewhat different:

EXERCISE 6.1.4. *On the other hand, find compact manifolds M, N each of which has a projective structure but $M \times N$ does not admit a projective structure.*

- (1) *If M_1, \dots, M_r are manifolds with real projective structures, show that the Cartesian product $M_1 \times \dots \times M_r \times T^{r-1}$ admits a projective structure.*

6.1.3. Fibrations. One can also *pull back* geometric structures by *fibrations* of geometries as follows. Let (G, X) be a homogeneous space and suppose that $X' \xrightarrow{\Phi} X$ is a fibration with fiber F and that $G' \xrightarrow{\phi} G$ is a homomorphism such that for each $g' \in G'$ the diagram

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X' \\ \Phi \downarrow & & \downarrow \Phi \\ X' & \xrightarrow{\phi(g')} & X' \end{array}$$

commutes.

Suppose that M is an (G, X) -manifold, with a universal covering space $\widetilde{M} \xrightarrow{\Pi} M$ with group of deck transformations π and developing

pair (dev, h) . Then the pullback $\text{dev}^*\Phi$ is an F -fibration \widetilde{M}' over \widetilde{M} and the induced map $M' \xrightarrow{\text{dev}'} X'$ is a local diffeomorphism and thus a developing map for an (G', X') -structure on \widetilde{M}' . We summarize these maps in the following commutative diagram:

$$\begin{array}{ccc} \widetilde{M}' & \xrightarrow{\text{dev}'} & X' \\ \downarrow & & \downarrow \Phi \\ \widetilde{M} & \xrightarrow{\text{dev}} & X \end{array}$$

Suppose that the holonomy representation $\pi \xrightarrow{h} G$ lifts to $\pi \xrightarrow{h'} G'$. (In general the question of whether h lifts will be detected by certain invariants in the cohomology of M .) Then h' defines an extension of the action of π on \widetilde{M} to \widetilde{M}' by (G', X') -automorphisms. Since the action of π on \widetilde{M}' is proper and free, the quotient $M' = \widetilde{M}'/\pi$ is an (G', X') -manifold. Moreover the fibration $\widetilde{M}' \rightarrow \widetilde{M}$ descends to an F -fibration $M' \rightarrow M$.

6.1.4. Suspensions. Before discussing Benzécri’s theorem and the classification of 2-dimensional affine manifolds, we describe several constructions for affine structures from affine structures and projective structures of lower dimension. Namely, let Σ be a smooth manifold and $\Sigma \xrightarrow{f} \Sigma$ a diffeomorphism. The mapping torus of f is defined to be the quotient $M = \mathbf{M}_f(\Sigma)$ of the product $\Sigma \times \mathbb{R}$ by the \mathbb{Z} -action defined by

$$(x, t) \mapsto (f^{-n}x, t + n)$$

It follows that dt defines a nonsingular closed 1-form ω on M tangent to the fibration

$$M \xrightarrow{t} S^1 = \mathbb{R}/\mathbb{Z}.$$

Furthermore the vector field $\frac{\partial}{\partial t}$ on $\Sigma \times \mathbb{R}$ defines a vector field S_f on M , the *suspension* of the diffeomorphism $\Sigma \xrightarrow{f} \Sigma$. The dynamics of f is mirrored in the dynamics of S_f : there is a natural correspondence between the orbits of f and the trajectories of S_f . The embedding $\Sigma \hookrightarrow \Sigma \times \{t\}$ is transverse to the vector field S_f and each trajectory of S_f meets Σ . Such a hypersurface is called a *cross-section* to the vector field. Given a cross-section Σ to a flow $\{\xi_t\}_{t \in \mathbb{R}}$, then (after possibly reparametrizing $\{\xi_t\}_{t \in \mathbb{R}}$), the flow can be recovered as a suspension. Namely, given $x \in \Sigma$, let $f(x)$ equal $\xi_t(x)$ for the smallest $t > 0$ such that $\xi_t(x) \in \Sigma$, that is, the first-return map or Poincaré map for $\{\xi_t\}_{t \in \mathbb{R}}$ on Σ . For the theory of cross-sections to flows see Fried [107].

6.2. Parallel structures in affine geometry

Perhaps the simplest case occurs when the model spaces are equal: $X = X'$ and $G < G'$. We saw this already in §1.4, for affine structures, where $X' = \mathbf{A}^n$ and $G' = \text{Aff}(\mathbf{A}^n)$. By imposing conditions on the linear holonomy $L(G) < \text{GL}(\mathbb{R}^n)$, one obtains refinements of affine geometry involving parallel structures. For example, when $G = L^{-1}\text{O}(n)$, one obtains Euclidean geometry as *affine geometry with a parallel Riemannian metric*. Replacing the Euclidean inner product on $V = \mathbb{R}^n$ with other bilinear forms \mathbf{B} and taking $G = L^{-1}(\text{O}(V; \mathbf{B}))$ yields affine structures with parallel tensor fields. Taking \mathbf{B} to be a Lorentzian inner product, yields a category of Ehresmann structures which correspond to *flat Lorentzian manifolds*, where $G = \text{Isom}(\mathbf{E}^{n-1,1}) = L^{-1}(\text{O}(n-1, 1))$.

If $n = 2m$, then $G = L^{-1}(\text{GL}(m, \mathbb{C}))$ realizes *complex affine geometry* as affine geometry with a *parallel almost complex structure* as in §1.4.3.

6.2.1. Flat tori and Euclidean structures. Recall that a *flat torus* is a Euclidean manifold of the form \mathbf{E}^n/Γ , where Γ is a *lattice* of translations. We can regard flat tori as (G, X) -manifolds where both X and G are the same vector space, and G is acting on X by translation. In fact, every closed (G, X) -manifold is a flat torus.

Bieberbach's structure theorem is essentially a qualitative structure theorem *classifying* closed Euclidean manifolds. It states that every closed Euclidean manifold is *finitely covered* by a flat torus. That is, given a closed Euclidean manifold M , there is a flat torus N and a finite subgroup $F \subset \text{Isom}(N)$ such that F acts freely on N and M is isometric to the quotient manifold N/F . For an extensive discussion see Charlap [63].

From the viewpoint of enlarging and refining geometric structures, this result may be stated as follows. Corresponding to F is a finite subgroup $\Phi \subset \text{O}(n)$, the linear holonomy group of M . Let $\mathbf{V}\Phi$ be the subgroup of $\text{Isom}(\mathbf{E}^n)$ generated by the translation group \mathbf{V} and Φ . Then Bieberbach's structure theorem can be restated as follows:

THEOREM 6.2.1. *Every closed $(\text{Isom}(\mathbf{E}^n), \mathbf{E}^n)$ -manifold has a $(\mathbf{V}\Phi, \mathbf{E}^n)$ structure for some finite subgroup $\Phi \subset \text{O}(n)$.*

6.2.2. Parallel suspensions. Let Σ be an affine manifold and $f \in \text{Aff}(\Sigma)$ an automorphism. We shall define an affine manifold M with a parallel vector field S_f and cross-section $\Sigma \hookrightarrow M$ such that the corresponding Poincaré map is f . (Compare §6.1.4.) We proceed as follows. Let $\Sigma \times \mathbf{A}^1$ be the Cartesian product with the product affine

structure and let $\Sigma \times \mathbf{A}^1 \xrightarrow{t} \mathbf{A}^1$ be an affine coordinate on the second factor. Then the map

$$\begin{aligned} \Sigma \times \mathbf{A}^1 &\xrightarrow{\tilde{f}} \Sigma \times \mathbf{A}^1 \\ (x, t) &\longmapsto (f^{-1}(x), t + 1) \end{aligned}$$

is affine and generates a free proper \mathbb{Z} -action on $\Sigma \times \mathbf{A}^1$, which t -covers the action of \mathbb{Z} on $\mathbf{A}^1 \cong \mathbb{R}$ by translation. Let M be the corresponding quotient affine manifold. Then d/dt is a parallel vector field on $\Sigma \times \mathbf{A}^1$ invariant under \tilde{f} and thus defines a parallel vector field S_f on M . Similarly the parallel 1-form dt on $\Sigma \times \mathbf{A}^1$ defines a parallel 1-form ω_f on M for which $\omega_f(S_f) = 1$. For each $t \in \mathbf{A}^1/\mathbb{Z}$, the inclusion $\Sigma \times \{t\} \hookrightarrow M$ defines a cross-section to S_f . We call (M, S_f) the *parallel suspension* or *affine mapping torus* of the affine automorphism (Σ, f) .

EXERCISE 6.2.2. *Suppose that N and Σ are affine manifolds and that*

$$\pi_1(\Sigma) \xrightarrow{\phi} \text{Aff}(N)$$

is an action of $\pi_1(\Sigma)$ on N by affine automorphisms. The flat N -bundle over Σ with holonomy ϕ is defined as the quotient of $\tilde{\Sigma} \times N$ by the diagonal action of $\pi_1(\Sigma)$ given by deck transformations on $\tilde{\Sigma}$ and by ϕ on N . Show that the total space M is an affine manifold such that the fibration $M \rightarrow \Sigma$ is an affine map and the the flat structure (the foliation of M induced by the foliation of $\tilde{\Sigma} \times N$ by leaves $\tilde{\Sigma} \times \{y\}$, for $y \in N$) is an affine foliation.

6.2.3. Closed Euclidean manifolds. The first example of a closed Euclidean manifold which is not a flat torus is a *Euclidean Klein bottle*. One can easily construct it as the *parallel suspension* of a free isometric involution of the *Euclidean circle* \mathbf{E}^1/\mathbb{Z} .

EXERCISE 6.2.3. *Compute the affine holonomy group of this complete affine surface. Show that it has the same rational homology as S^1 .*

6.2.3.1. A Euclidean \mathbb{Q} -homology 3-sphere. Here is an interesting example in dimension 3, which we denote by $\mathcal{S}_{\mathbb{Q}}^3$.

Here is the construction of $\mathcal{S}_{\mathbb{Q}}^3$ as a closed Euclidean 3-manifold.

Consider the group $\Gamma \subset \text{Isom}(\mathbb{E}^3)$ generated by the three isometries

$$A = \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array} \right]$$

$$B = \left[\begin{array}{ccc|c} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{array} \right]$$

$$C = \left[\begin{array}{ccc|c} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

and Γ is a discrete group of Euclidean isometries which acts properly and freely on \mathbb{R}^3 with quotient a compact 3-manifold M . Furthermore there is a short exact sequence

$$\mathbb{Z}^3 \cong \langle A^2, B^2, C^2 \rangle \hookrightarrow \Gamma \xrightarrow{\text{L}} \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

We denote the quotient $\Gamma \backslash \mathbb{E}^3$ by $\mathcal{S}_{\mathbb{Q}}^3$. It is a Euclidean manifold, which has a regular $\mathbb{Z}/2 \oplus \mathbb{Z}/2$ -covering space by a torus, and hence admits a complete affine structure.

EXERCISE 6.2.4. *Let $M^3 = \Gamma \backslash \mathbb{E}^3$ as above.*

- *Show that M^3 has the same rational homology as S^3 .*
- *Prove that every closed Euclidean 3-manifold is either a parallel suspension of an isometry of a closed Euclidean 2-manifold or is isometric to M^3 .*

Later we will see that every affine structure on $\mathcal{S}_{\mathbb{Q}}^3$ must be complete, and indeed a Euclidean structure as above.

6.3. Homogeneous subdomains

6.3.1. Projective structures from non-Euclidean geometry.

Using the Klein model of hyperbolic geometry

$$(\mathbb{H}^n, \text{PO}(n, 1)) \hookrightarrow (\mathbb{P}^n, \text{Proj}(\mathbb{P}^n))$$

every hyperbolic-geometry structure (that is, Riemannian metric of constant curvature -1) determines a projective structure. Using the inclusion of the projective orthogonal group $\text{PO}(n+1) \subset \text{PGL}(n+1; \mathbb{R})$ one sees that every elliptic-geometry structure (that is, Riemannian metric of constant curvature $+1$) determines a projective structure. Since every surface admits a metric of constant curvature, we obtain the following:

THEOREM 6.3.1. *Every surface admits an \mathbb{RP}^2 -structure.*

Similarly, the Poincaré model of 2-dimensional hyperbolic geometry

$$(\mathbb{H}^2, \mathrm{PGL}(2, \mathbb{R})) \hookrightarrow (\mathbb{CP}^1, \mathrm{PGL}(2, \mathbb{C}))$$

embeds the hyperbolic plane in complex-projective 1-dimensional geometry, and every hyperbolic structure on a surface determines a \mathbb{CP}^1 -structure.

THEOREM 6.3.2. *Every surface admits a \mathbb{CP}^1 -structure.*

6.4. Hopf manifolds

The basic example of an incomplete affine structure on a closed manifold is a *Hopf manifold*. (The one-dimensional cases were introduced in §5.4.2.) Consider the domain

$$\Omega := \mathbb{R}^n - \{0\}.$$

The group \mathbb{R}^+ of *positive homotheties* (that is, scalar multiplications) acts on Ω properly and freely. Indeed, there is an \mathbb{R}^+ -equivariant homeomorphism

$$(25) \quad \begin{aligned} \Omega &\xrightarrow{h} \mathbb{R} \times S^{n-1} \\ \mathbf{v} &\longmapsto (\log(\|\mathbf{v}\|), \mathbf{v}/\|\mathbf{v}\|) \end{aligned}$$

where \mathbb{R}^+ acts by translation on the first factor and identically on the second. Clearly the affine structure on Ω is incomplete. If $\lambda \in \mathbb{R}$ and $\lambda > 1$, then the cyclic group $\langle \lambda \rangle$ is a discrete subgroup of \mathbb{R}^+ and the quotient $\Omega/\langle \lambda \rangle$ is a compact incomplete affine manifold M . We shall denote this manifold by \mathbf{Hopf}_λ^n . (A geodesic whose tangent vector “points” at the origin will be incomplete; on the manifold M the affinely parametrized geodesic will circle around with shorter and shorter period until in a finite amount of time will “run off” the manifold.) If $n = 1$, then M consists of two disjoint copies of the *Hopf circle* $\mathbb{R}^+/\langle \lambda \rangle$ — this manifold is an incomplete closed geodesic (and every incomplete closed geodesic is isomorphic to a Hopf circle). For $n > 1$, then M is connected and is diffeomorphic to the product $S^1 \times S^{n-1}$. For $n > 2$ both the holonomy homomorphism and the developing map are injective.

If $n = 2$, then M is a torus whose holonomy homomorphism maps $\pi_1(M) \cong \mathbb{Z} \oplus \mathbb{Z}$ onto the cyclic group $\langle \lambda \rangle$. Note that $\widetilde{M} \xrightarrow{\mathrm{dev}} \mathbb{R}^2$ is neither injective nor surjective, although it is a covering map onto its image. For $k \geq 1$ let $\pi^{(k)} \subset \pi$ be the unique subgroup of index k which intersects $\mathrm{Ker}(h) \cong \mathbb{Z}$ in a subgroup of index k . Let $M^{(k)}$ denote the corresponding covering space of M . Then $M^{(k)}$ is another closed affine

manifold diffeomorphic to a torus whose holonomy homomorphism is a surjection of $\mathbb{Z} \oplus \mathbb{Z}$ onto $\langle \lambda \rangle$.

EXERCISE 6.4.1. *Show that for $k \neq l$, the two affine manifolds $M^{(k)}$ and $M^{(l)}$ are not isomorphic. (Hint: consider the invariant defined as the least number of breaks of a broken geodesic representing a simple closed curve on M whose holonomy is trivial.) Thus two different affine structures on the same manifold can have the same holonomy homomorphism.*

EXERCISE 6.4.2. *Suppose that $\lambda < -1$. Then $M = (\mathbb{R}^n - \{0\})/\langle \lambda \rangle$ is an incomplete compact affine manifold doubly covered by Hopf_λ^n . What is M topologically?*

EXERCISE 6.4.3. *Let $A \in \text{GL}(n, \mathbb{R})$ be a linear expansion, that is a linear map all of whose eigenvalues have modulus > 1 . Suppose that A preserves orientation, that is, $\det(A) > 0$. Then for every $\lambda > 1$, find a homeomorphism*

$$\mathbb{R}^n \xrightarrow{\phi} \mathbb{R}^n$$

such that $\phi(A(\mathbf{v})) = \lambda\phi(\mathbf{v})$. Show that $(\mathbb{R}^n \setminus \{\mathbf{0}\})/\langle A \rangle$ is a closed incomplete affine manifold homeomorphic to $S^{n-1} \times S^1$.

What can you say if $\det(A) < 0$?

Hopf manifolds play an important role in relating projective structures and affine structures on closed manifolds.

6.4.1. Geodesics on Hopf manifolds. These *geodesically incomplete* structures model *incomplete closed geodesics* on affine manifolds. Namely, the geodesic on \mathbf{A}^1 defined by

$$t \longmapsto 1 + t(\lambda^{-1} - 1)$$

begins at 1 and in time

$$t_\infty := 1 + \lambda^{-1} + \lambda^{-2} + \cdots = (1 - \lambda^{-1})^{-1} > 0$$

reaches 0. It defines a closed incomplete closed geodesic $p(t)$ on M starting at $p(0) = p_0$. The lift

$$(-\infty, t_\infty) \xrightarrow{\tilde{p}} \tilde{M}$$

satisfies

$$\text{dev}(\tilde{p}(t)) = 1 + t(\lambda^{-1} - 1),$$

which uniquely specifies the geodesic $p(t)$ on M . It is a geodesic since its velocity

$$p'(t) = (\lambda^{-1} - 1)\partial_x$$

is constant (parallel). However $p(t_n) = p_0$ for

$$t_n := \frac{1 - \lambda^{-n}}{1 - \lambda^{-1}} = 1 + \lambda^{-1} + \cdots + \lambda^{1-n}$$

and as viewed in M , seems to go “faster and faster” through each cycle. By time $t_\infty = \lim_{n \rightarrow \infty} t_n$, it seems to “run off the manifold:” the geodesic is only defined for $t < t_\infty$. The apparent paradox is that $p(t)$ has *zero acceleration*: it would have “constant speed” if “speed” were only defined.

6.4.2. The sphere of directions. An important example is the following, which in many contexts is a more useful model space than projective space.

DEFINITION 6.4.4. *Let V be an \mathbb{R} -vector space with origin $\mathbf{0}$. Define the sphere of directions in V as the quotient space of $V \setminus \{\mathbf{0}\}$ by the group \mathbb{R}^+ of positive scalar multiplications, and denote it by:*

$$\mathbb{S}(V) := V \setminus \{\mathbf{0}\} / \mathbb{R}^+$$

If $V = \mathbb{R}^{n+1}$, write $\mathbb{S}^n := \mathbb{S}(V)$.

EXERCISE 6.4.5. *Let $V = \mathbb{R}^{n+1}$ and $G = \mathrm{GL}(n+1; \mathbb{R})$ its group of linear automorphisms.*

- *Show that $\mathbb{S}(V) \approx S^n$ by explicitly constructing a section of the principal \mathbb{R}^+ -bundle defined by the quotient $V \setminus \{\mathbf{0}\} \rightarrow \mathbb{S}(V)$.*
- *Construct an explicit double covering $\mathbb{S}(V) \rightarrow \mathbb{P}(V)$, realizing the sphere of directions as the universal covering space of projective space.*
- *Show that the action of the collineation group $\mathrm{PGL}(n+1, \mathbb{R})$ lifts to the linear action of $\mathrm{GL}(n+1, \mathbb{R})$ on \mathbb{S}^n and compute its kernel of the action of $\mathrm{GL}(n+1, \mathbb{R})$ on \mathbb{S}^n .*

This construction relates to the Hopf manifolds Hopf_λ^n as follows. For each $\lambda > 1$, form the quotient by the cyclic subgroup $\langle \lambda \rangle < \mathbb{R}^+$ rather than all of \mathbb{R}^+ . The resulting quotient map is a principal $\mathbb{R}^+ / \langle \lambda \rangle$ -fibration

$$\mathrm{Hopf}_\lambda^{n+1} \rightarrow S^n$$

which is $\mathrm{GL}(n+1, \mathbb{R})$ -equivariant.

6.4.3. Hopf tori. There is another point of view concerning Hopf manifolds in dimension two. Let M be a two-torus; we may explicitly realize M as a quotient \mathbb{C} / Λ where $\Lambda \subset \mathbb{C}$ is a lattice. The complex exponential map $\mathbb{C} \xrightarrow{\exp} \mathbb{C}^\times$ is a universal covering space having the property that

$$\exp \circ \tau(z) = e^z \cdot \exp$$

where $\tau(z)$ denotes translation by $z \in \mathbb{C}$. For various choices of lattices Λ , the exponential map

$$\tilde{M} = \mathbb{C} \xrightarrow{\exp} \mathbb{C}^\times$$

is a developing map for a (complex) affine structure on M with holonomy homomorphism

$$\pi \cong \Lambda \xrightarrow{\exp} \exp(\Lambda) \hookrightarrow \mathbb{C}^\times \subset \text{Aff}(\mathbb{C})$$

We denote this affine manifold by $\exp(\mathbb{C}/\Lambda)$; it is an incomplete complex affine 1-manifold or equivalently an incomplete similarity 2-manifold. Every compact incomplete orientable similarity manifold is equivalent to an $\exp(\mathbb{C}/\Lambda)$ for a unique lattice $\Lambda \subset \mathbb{C}$. Taking $\Lambda \subset \mathbb{C}$ to be the lattice generated by $\log \lambda$ and $2\pi i$ we obtain the Hopf manifold Hopf_λ^2 . More generally the lattice generated by $\log \lambda$ and $2k\pi i$ corresponds to the k -fold covering space of Hopf_λ^2 described above. There are “fractional” covering spaces of the Hopf manifold obtained from the lattice generated by $\log \lambda$ and $2\pi/n$ for $n > 1$; these manifolds admit n -fold covering spaces by Hopf_λ^2 . The affine manifold M admits no closed geodesics if and only if $\Lambda \cap \mathbb{R} = \{0\}$. Note that the exponential map defines an isomorphism $\mathbb{C}/\Lambda \rightarrow M$ which is definitely not an isomorphism of affine manifolds.

Any $\lambda > 1$ generates a lattice inside the multiplicative group \mathbb{R}_+ , which acts affinely on \mathbb{A}^1 . The quotient $\mathbb{R}_+/\langle \lambda \rangle$ also defines an affine structure on M , which is not a Euclidean structure since dilation by λ is not an isometry. Explicitly, take f to be a diffeomorphism onto the interval $[1, \lambda] \subset \mathbb{R} \approx \mathbb{A}^1$, so that dev is a diffeomorphism of \tilde{M} onto $(0, \infty) = \mathbb{R}_+ \subset \mathbb{A}^1$.

Like the preceding example, this affine structure is also bi-invariant with respect to the natural Lie group structure on $\mathbb{R}_+/\langle \lambda \rangle$.

Observe that, since the exponential map

$$\begin{aligned} \mathbb{R} &\longrightarrow \mathbb{R}_+ \\ x &\longmapsto e^x \end{aligned}$$

converts addition (translation) to multiplication (dilation), it defines a diffeomorphism between two quotients

$$\mathbb{R}/l\mathbb{Z} \longrightarrow \mathbb{R}_+/\langle \lambda \rangle$$

where $l := \log(\lambda)$. This map also defines a (non-affine) analytic isomorphism between the corresponding Lie groups.

The preceding construction then applies and we obtain a radiant affine structure on the total space M' of a principal \mathbb{R}^+ -bundle over M with holonomy representation \tilde{h} . The radiant vector field $\mathbf{R}_{M'}$ generates

the (fiberwise) action of \mathbb{R}^+ ; this action of \mathbb{R}^\times on M' is affine, given locally in affine coordinates by homotheties. (This construction is due to Benzécri [35] where the affine manifolds are called *variétés coniques affines*. He observes there that this construction defines an embedding of the category of $\mathbb{R}P^n$ -manifolds into the category of $(n + 1)$ -dimensional affine manifolds.)

Since \mathbb{R}^+ is contractible, every principal \mathbb{R}^+ -bundle is trivial (although there is in general no preferred trivialization). Choose any $\lambda > 1$; then the cyclic group $\langle \lambda \rangle \subset \mathbb{R}^+$ acts properly and freely on M' by affine transformations. We denote the resulting affine manifold by M'_λ and observe that it is homeomorphic to $M \times S^1$. (Alternatively, one may work directly with the Hopf manifold $\text{Hopf}_\lambda^{n+1}$ and its \mathbb{R}^\times -fibration $\text{Hopf}_\lambda^{n+1} \rightarrow \mathbb{R}P^n$.) We thus obtain:

PROPOSITION 6.4.6 (Benzécri [35], §2.3.1). *Suppose that M is an $\mathbb{R}P^n$ -manifold. Let $\lambda > 1$. Then $M \times S^1$ admits a radiant affine structure for which the trajectories of the radiant vector field are all closed geodesics each affinely isomorphic to the Hopf circle $\mathbb{R}^+/\langle \lambda \rangle$.*

Since every (closed) surface admits an $\mathbb{R}P^2$ -structure, we obtain:

COROLLARY 6.4.7 (Benzécri [35]). *Let Σ be a closed surface. Then $\Sigma \times S^1$ admits an affine structure.*

If Σ is a closed hyperbolic surface, the affine structure on $M = \Sigma \times S^1$ can be described as follows. A developing map maps the universal covering of M onto the convex cone

$$\Omega = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 - z^2 < 0, z > 0\}$$

which is invariant under the identity component G of $\text{SO}(2, 1)$. The group $G \times \mathbb{R}^+$ acts transitively on Ω with isotropy group $\text{SO}(2)$. Choosing a hyperbolic structure on Σ determines an isomorphism of $\pi_1(\Sigma)$ onto a discrete subgroup Γ of G ; then for each $\lambda > 1$, the group $\Gamma \times \langle \lambda \rangle$ acts properly and freely on Ω with quotient the compact affine 3-manifold M .

EXERCISE 6.4.8. *A $\mathbb{C}P^n$ -structure is a geometric structure modeled on complex projective space $\mathbb{C}P^n$ with coordinate changes locally from the projective group $\text{PGL}(n + 1; \mathbb{C})$. Let M be a $\mathbb{C}P^n$ -manifold.*

- *Show that there is a T^2 -bundle over M which admits a complex affine structure and an S^1 -bundle over M which admits an $\mathbb{R}P^{2n+1}$ -structure.*
- *Show that this is a contact $\mathbb{R}P^{2n+1}$ -structure as defined in Exercise 6.1.2.*

Compare Guichard-Wienhard [141].

Suppose that \mathfrak{F} is a foliation of a manifold M ; then \mathfrak{F} is locally defined by an atlas of smooth submersions $U \rightarrow \mathbb{R}^q$ for coordinate patches U . An (G, X) -atlas transverse to \mathfrak{F} is defined to be a collection of coordinate patches U_α and coordinate charts

$$U_\alpha \xrightarrow{\psi_\alpha} X$$

such that for each pair (U_α, U_β) and each component $C \subset U_\alpha \cap U_\beta$ there exists an element $g_C \in G$ such that

$$g_C \circ \psi_\alpha = \psi_\beta$$

on C . An (G, X) -structure transverse to \mathfrak{F} is a maximal (G, X) -atlas transverse to \mathfrak{F} . Consider an (G, X) -structure transverse to \mathfrak{F} ; then an immersion $\Sigma \xrightarrow{f} M$ which is transverse to \mathfrak{F} induces an (G, X) -structure on Σ .

A foliation \mathfrak{F} of an affine manifold is said to be *affine* if its leaves are parallel affine subspaces (that is, totally geodesic subspaces). It is easy to see that transverse to an affine foliation of an affine manifold is a natural affine structure. In particular if M is an affine manifold and ζ is a parallel vector field on M , then ζ determines a one-dimensional affine foliation which thus has a transverse affine structure. Moreover if Σ is a cross-section to ζ , then Σ has a natural affine structure for which the Poincaré map $\Sigma \rightarrow \Sigma$ is affine.

EXERCISE 6.4.9. *Show that the Hopf manifold Hopf_λ^n has an affine foliation with one closed leaf if $n > 1$ (two if $n = 1$) and its complement consists of two Reeb components.*

6.5. Radiant manifolds

A Hopf manifold is the prototypical example of a *radiant affine manifold*. Many properties of Hopf manifolds are shared by radiant structures, and use the existence of a radiant vector field. For example, a closed radiant affine manifold M is always incomplete, and a radiant vector field is always nonsingular. Therefore $\chi(M) = 0$.

In this section we discuss general properties of radiant affine structures.

6.5.1. Radiant vector fields. Recall, from §1.5.2, that a vector field R on an affine manifold M is *radiant* if it is locally equivalent to the *Euler vector field*,

$$R_0 := \sum_{i=1}^n x^i \frac{\partial}{\partial x^i}$$

Exercise 1.5.6 gives alternate characterizations of radiance.

PROPOSITION 6.5.1. *Let M be an affine manifold with development pair $(\text{dev}, \mathfrak{h})$. The following conditions are equivalent:*

- *The affine holonomy group $\Gamma = \mathfrak{h}(\pi)$ fixes a point in \mathbf{A} (by conjugation we may assume this fixed point is the origin $\mathbf{0} \in \mathbf{V}$);*
- *M is isomorphic to a $(\mathbf{V}, \text{GL}(\mathbf{V}))$ -manifold;*
- *M possesses a radiant vector field \mathbf{R}_M .*

If \mathbf{R}_M is a radiant vector field on M , we shall often refer to the pair (M, \mathbf{R}_M) as well as a *radiant affine structure*. Then a Γ -invariant radiant vector field $\mathbf{R}_\mathbf{A}$ on \mathbf{A} exists, such that

$$\Pi^*\mathbf{R}_M = \text{dev}^*\mathbf{R}_\mathbf{A}.$$

THEOREM 6.5.2. *The developing image $\text{dev}(\widetilde{M})$ does not contain any stationary points of the affine holonomy.*

COROLLARY 6.5.3. *Let (M, \mathbf{R}_M) be a closed radiant affine manifold.*

- *M is incomplete.*
- *The radiant vector field \mathbf{R}_M is nonsingular.*
- *The Euler characteristic $\chi(M) = 0$.*

PROOF. Choosing affine coordinates (x^1, \dots, x^n) and a developing pair $(\text{dev}, \mathfrak{h})$, we may assume that $\mathbf{0}$ is fixed by the affine holonomy $\mathfrak{h}(\pi_1(M))$, so that

$$\mathbf{R}_\mathbf{A} = \sum_{i=1}^n x^i \frac{\partial}{\partial x^i}$$

with flow

$$x \xrightarrow{\Psi_t} e^t x k.$$

We prove that $0 \notin \text{dev}(\widetilde{M})$.

We find a vector field $\widetilde{\mathbf{R}} \in \text{Vec}(\widetilde{M})$ which is Π -related to $\mathbf{R}_\mathbf{A}$. Since dev is a local diffeomorphism, the pullback $\widetilde{\mathbf{R}} := \text{dev}^*(\mathbf{R}_\mathbf{A})$ is defined by (3). Let $\widetilde{\Phi}_t$ be the corresponding local flow. By the Naturality of Flows,

$$\text{dev}\widetilde{\Phi}_t(x) = e^t \text{dev}(x).$$

M is radiant, so \mathfrak{h} preserves $\mathbf{R}_\mathbf{A}$ and therefore $\widetilde{\mathbf{R}}$ is $\pi_1(M)$ -invariant. It follows that $\exists \mathbf{R}_M \in \text{Vec}(M)$ which is Π -related to $\widetilde{\mathbf{R}}$:

$$\Pi^*\mathbf{R}_M = \widetilde{\mathbf{R}}.$$

Since M is closed, \mathbf{R}_M integrates to a global flow $M \xrightarrow{\Pi_t}$, defined $\forall t \in \mathbb{R}$. Exercise 6.5.4 (below) implies that $\widetilde{\mathbf{R}}$ is complete and integrates to a flow

$$\widetilde{M} \xrightarrow{\widetilde{\Phi}_t} \widetilde{M}$$

such that

$$\Pi \circ \widetilde{\Phi}_t = \Phi_t$$

Let $M_{\mathbf{0}} := \Pi(\text{dev}^{-1}(0))$. Since Π and dev are local diffeomorphisms and $0 \in \mathbf{A}$ is a $\mathfrak{h}(\pi_1(M))$ -invariant discrete set, $M_{\mathbf{0}} \subset M$ is discrete set. Compactness of M implies that $M_{\mathbf{0}}$ is a finite set. We show that $M_{\mathbf{0}} = \emptyset$.

Since the only zero of $\mathbf{R}_{\mathbf{A}}$ is the origin $\mathbf{0}$, the vector field \mathbf{R}_M is nonsingular on the complement of $M_{\mathbf{0}}$. Choose a neighborhood U of $M_{\mathbf{0}}$, each component of which develops to a small ball B about 0 in \mathbf{A} . Let $K \subset \widetilde{M}$ be a compact set such that the saturation $\Pi(K) = M$; then $\exists N \gg 0$ such that

$$e^{-t}(\text{dev}(K)) \subset B$$

for $t \geq N$. Thus $\widetilde{\Phi}_t(K) \subset B$ for $t \leq -N$. It follows that U is an attractor for the flow of $-\mathbf{R}_M$, that is, $\Phi_{-t}(M) \subset U$ for $t \gg N$. Consequently $M \xrightarrow{\Phi_N} U$ deformation retracts the closed manifold M onto U . Since a closed manifold is not homotopy-equivalent to a finite set, this contradiction implies $M_{\mathbf{0}} = \emptyset$ and $\mathbf{0} \notin \text{dev}(\widetilde{M})$ as desired. \square

EXERCISE 6.5.4. *Let $M \xrightarrow{f} N$ be a local diffeomorphism between smooth manifolds, and $\xi \in \text{Vec}(M), \eta \in \text{Vec}(N)$ be f -related vector fields. Suppose that f is a covering space. Then ξ is complete if and only if η is complete.*

THEOREM 6.5.5. *Let M be a compact radiant manifold. Then M cannot have parallel volume. (In other words a compact manifold cannot support a $(\mathbb{R}^n, \text{SL}(n; \mathbb{R}))$ -structure.)*

PROOF. Let $\omega_{\mathbf{A}} = dx^1 \wedge \cdots \wedge dx^n$ be a parallel volume form on \mathbf{A} and let ω_M be the corresponding parallel volume form on M , that is, $\Pi^*\omega_M = \text{dev}^*\omega_{\mathbf{A}}$. The interior product

$$\eta_M := \frac{1}{n} \iota_{\mathbf{R}_M} \omega_M$$

is an $(n-1)$ -form on M . Since

$$d\iota_{\mathbf{R}_M} \omega_M = n\omega_M,$$

$d\eta_M = \omega_M$. However, ω_M is a volume form on M and

$$0 < \text{vol}(M) = \int_M \omega_M = \int_M d\eta_M = 0$$

a contradiction. \square

Intuitively, the main idea in the proof above is that the radiant flow on M expands the parallel volume uniformly. Thus by “conservation of volume” a compact manifold cannot support both a radiant vector field and a parallel volume form.

EXERCISE 6.5.6. *The first Betti number of a closed radiant affine manifold is always positive. (Hint: Compare Exercise 11.1.2.)*

6.5.2. Radiant suspensions. Let (M, R_M) be a radiant affine manifold of dimension $+1$. Transverse to R_M is an $\mathbb{R}P^n$ -structure, as follows. In local affine coordinates the trajectories of R_M are rays through the origin in \mathbb{R}^{n+1} and projectivization maps coordinate patches submersively into $\mathbb{R}P^n$. In particular, if Σ is an n -manifold and $\Sigma \xrightarrow{f} M$ is transverse to R_M , then f determines an $\mathbb{R}P^n$ -structure on Σ .

PROPOSITION 6.5.7. *Let Σ be a compact $\mathbb{R}P^n$ -manifold and $f \in \text{Aut}(\Sigma)$ a projective automorphism. Then there exists a radiant affine manifold (M, R_M) and a cross-section $\Sigma \xrightarrow{\iota} M$ to R_M such that the Poincaré map for ι equals $\iota^{-1} \circ f \circ \iota$. In other words, the mapping torus of a projective automorphism of an compact $\mathbb{R}P^n$ -manifold admits a radiant affine structure.*

PROOF. Let S^n be the double covering of $\mathbb{R}P^n$ (realized as the sphere of directions in \mathbb{R}^{n+1}) and let

$$\mathbb{R}^{n+1} \setminus \{0\} \xrightarrow{\Phi} S^n$$

be the corresponding principal \mathbb{R}^+ -fibration. Let N be the principal \mathbb{R}^+ -bundle over M constructed in §6.4.2 and choose a section $M \xrightarrow{\sigma} N$. Let $\{\xi_t\}_{t \in \mathbb{R}}$ be the radiant flow on N and denote by $\{\tilde{\xi}_t\}_{t \in \mathbb{R}}$ the radiant flow on \tilde{N} . Let (dev, h) be a development pair; then f lifts to an affine automorphism \tilde{f} of \tilde{M} . Furthermore there exists a projective automorphism $g \in \text{GL}(n+1; \mathbb{R})/\mathbb{R}^+$ of the sphere of directions S^n such that

$$\begin{array}{ccc} \tilde{N} & \xrightarrow{\text{dev}'} & \mathbb{R}^{n+1} \setminus \{0\} \\ \tilde{f} \downarrow & & \downarrow g \\ \tilde{N} & \xrightarrow{\text{dev}'} & \mathbb{R}^{n+1} \setminus \{0\} \end{array}$$

commutes. Choose a compact set $K \subset \widetilde{M}$ such that

$$\pi_1(M) \cdot K = \widetilde{M}.$$

Let $\tilde{K} \subset \tilde{N}$ be the image of K under a lift of σ to a section $\widetilde{M} \rightarrow \tilde{N}$. Then

$$\tilde{K} \cap \tilde{f}\tilde{\xi}_t(\tilde{K}) = \emptyset$$

whenever $t > t_0$, for some t_0 . It follows that the affine automorphism $\xi_t \tilde{f}$ generates a free and proper affine \mathbb{Z} -action on N for $t > t_0$. We denote the quotient by M . In terms of the trivialization of $N \rightarrow M$ arising from σ , it is clear that the quotient of N by this \mathbb{Z} -action is diffeomorphic to the mapping torus of f . Furthermore the section σ defines a cross-section $\Sigma \hookrightarrow M$ to \mathbf{R}_M whose Poincaré map corresponds to f . \square

We call the radiant affine manifold (M, \mathbf{R}_M) the *radiant suspension* of the pair (Σ, f) .

EXERCISE 6.5.8. *Express the Hopf manifolds of Exercise 6.4.3 as radiant suspensions of the automorphism of S^{n-1} given by the linear expansion A of \mathbb{R}^n .*

6.5.2.1. *Radiant similarity manifolds.* Hopf manifolds provide another example of a refined geometric structure, which arise in the classification of similarity structures on closed manifolds (§11.4).

EXERCISE 6.5.9. *Let $X = \mathbb{E}^n \setminus \{0\}$ and $G \subset \text{Sim}(\mathbb{E}^n)$ the stabilizer of 0. Let M be a compact (G, X) -manifold with holonomy group $\Gamma \subset G$.*

- *Prove that $G \cong \mathbb{R}^+ \times \mathbf{O}(n)$.*
- *Find a G -invariant Riemannian metric \mathfrak{g}_0 on X .*
- *Suppose that $n > 2$. Prove that $M \cong \Gamma \backslash X$, and that M admits a finite covering space isomorphic to a Hopf manifold.*
- *Suppose $n = 2$. Find an example where M is not isomorphic to $\Gamma \backslash X$.*

6.5.2.2. *Radiant affine surfaces.* A small modification of these constructions lead to the classification of radiant affine structures on closed 2-manifolds, and, with the classification of affine Lie group structures on the 2-torus, to the full list of closed affine 2-manifolds.

By Benzécri's theorem 9.1.1, every closed affine 2-manifold is homeomorphic to a torus or a Klein bottle. By passing to a covering space we can reduce to affine structures on a 2-torus T^2 .

These examples are obtained as follows. Begin with a linear expansion A of \mathbb{R}^2 . The cyclic group $\langle A \rangle$ acts properly on $\Omega := \mathbb{R}^2 - \{0\}$ with quotient M_A homeomorphic to a torus, as in Exercise 6.4.3.

Here is an explicit description of the development, which will be necessary to describe the modifications needed for all inhomogeneous affine tori. Choose a circle \mathcal{C} centered at $\mathbf{0} \in \mathbb{R}^2$. Then \mathcal{C} and its image $\mathcal{C}' := A(\mathcal{C})$ cobound an annulus $\mathcal{A} \subset \Omega$, which is a fundamental domain for the action of the holonomy group $\langle A \rangle$ on Ω .

Choose a point $\tilde{x}_0 \in \mathcal{C}$ and an arc

$$\tilde{x}_0 \xrightarrow{\tilde{a}} A(\tilde{x}_0)$$

in \mathcal{A} . Split \mathcal{A} along \tilde{a} obtaining a quadrilateral \square with four sides:

- S1:** \mathcal{C} split along \tilde{x}_0 ;
- S2:** The original arc \tilde{a} ;
- S3:** $A(\mathcal{C})$ split along $A(\tilde{x}_0)$;
- S4:** Another arc corresponding to \tilde{a} .

The annulus \mathcal{A} is the quotient of \square by an identification \lfloor which identifies sides S2 and S4. The torus M_A is the quotient of \mathcal{A} by A , which induces an identification of sides S1 and S3.

The image x_0 of the vertex \tilde{x}_0 of \square serves as a basepoint in M_A , and the fundamental group $\pi := \pi_1(M_A, x_0)$ is free abelian. Relative homotopy classes of the based loops corresponding to S1 and S2 define elements $a, b \in \pi$, respectively, which form a basis of π .

A model for the universal covering space \widetilde{M}_A of M_A is then the quotient of $\square \times \pi$ by identifications described above. The mapping of \square into Ω extending the embedding on the interior of \square generates a developing map $\widetilde{M}_A \rightarrow \Omega$. The corresponding holonomy homomorphism \mathfrak{h} maps a to A and b to the identity.

One can modify this construction in various ways. One modification involves passing to an n -fold covering space with the “same holonomy.” That is, one passes to the covering space $\widetilde{M}/\langle a, b^n \rangle$ which unwinds in the direction with trivial holonomy. These manifolds are all quotients of the n -fold covering space $\Omega^{(n)}$ of Ω .

All of these holonomy groups are cyclic, and the developing map factors through the covering space $\Omega^{(n)} \rightarrow \Omega$.

However we can modify these structures so that the holonomy of b is nontrivial, and find examples where the holonomy homomorphism is *injective*. Choose an affine transformation β which commutes with A ; since $\mathbf{0}$ is the unique point fixed by A , the affine transformation β is necessarily *linear*. We can replace Ω by the quotient Ω_β of $\square \times \pi$ by identifications generated by taking S1 to S2 by β . Equivalently, Ω_β is the quotient of the universal covering space $\widetilde{\Omega}$ by the cyclic group $\langle b \circ \widetilde{\beta} \rangle$ where $\widetilde{\beta}$ is the mapping on $\widetilde{\Omega}$ induced by β .

Since A commutes with β , it defines an affine automorphism A_β of Ω_β , and the quotient

$$M_{A,\beta} := \Omega_\beta / \langle A_\beta \rangle$$

is a radiant affine torus with holonomy homomorphism

$$\begin{aligned} \pi &\xrightarrow{h} \text{Aff}(\mathbb{A}^2) \\ a &\longmapsto A \\ b &\longmapsto \beta \end{aligned}$$

EXERCISE 6.5.10. *Express these manifolds as radiant suspensions of automorphisms of closed $\mathbb{R}\mathbb{P}^1$ -manifolds.*

Compare Exercise 6.4.3.

Every closed orientable affine 2-manifold which is *not* covered by an affine Lie group is one of these manifolds.

EXERCISE 6.5.11. *Find an example of an affine Lie group which is one of these manifolds.*

These turn out to be the only affine structures that are *not homogeneous*: indeed every other affine structure on a 2-torus is an affine commutative Lie group.

6.5.2.3. *Cross-sections to the radiant flow.* A natural question is whether every closed radiant affine manifold is a radiant suspension. A radiant affine manifold (M, \mathbb{R}) is a radiant suspension if and only if the flow of \mathbb{R} admits a cross-section. David Fried [105, 109] constructed a closed affine 6-manifold with diagonal holonomy whose radiant flow admits no cross-section. Choi [69] (using work of Barbot [19]) proves that every radiant affine 3-manifold is a radiant suspension, and therefore is either a Seifert 3-manifold covered by a product $F \times S^1$, where F is a closed surface, a nilmanifold or a hyperbolic torus bundle.

In dimensions 1 and 2 all closed radiant manifolds are radiant suspensions. When M is *hyperbolic*, that is, a quotient of a sharp convex cone (see Chapter 12), the existence of the *Koszul 1-form* implies that M is a radiant suspension.

In general affine automorphisms of affine manifolds can display quite complicated dynamics and thus the flows of parallel vector fields and radiant vector fields can be similarly complicated. For example, any element of $\text{GL}(2; \mathbb{Z})$ acts affinely on the flat torus $\mathbb{R}^2/\mathbb{Z}^2$; the most interesting of these are the hyperbolic elements of $\text{GL}(2; \mathbb{Z})$ which determine Anosov diffeomorphisms on the torus. Their suspensions thus determine Anosov flows on affine 3-manifolds which are generated by parallel or radiant vector fields. Indeed, it can be shown (Fried [108])

that every Anosov automorphism of a nilmanifold M can be made affine for some complete affine structure on M .

As a simple example of this we consider the linear diffeomorphism of the two-torus $T^2 = \mathbb{R}^2/\mathbb{Z}^2$ defined by a hyperbolic element $A \in \mathrm{GL}(2; \mathbb{Z})$. The parallel suspension of A is the complete affine 3-manifold \mathbb{R}^3/Γ where $\Gamma \subset \mathrm{Aff}(\mathbb{R}^3)$ consists of the affine transformations

$$\left[\begin{array}{cc|c} A^n & 0 & p \\ 0 & 1 & n \end{array} \right]$$

where $n \in \mathbb{Z}$ and $p \in \mathbb{Z}^2$. Since A is conjugate in $\mathrm{SL}(2; \mathbb{R})$ to a diagonal matrix with reciprocal eigenvalues, Γ is conjugate to a discrete cocompact subgroup of the subgroup of $\mathrm{Aff}(\mathbb{R}^3)$

$$G = \left\{ \left[\begin{array}{ccc|c} e^u & 0 & 0 & s \\ 0 & e^{-u} & 0 & t \\ 0 & 0 & 1 & u \end{array} \right] \mid s, t, u \in \mathbb{R} \right\}$$

which acts simply transitively. Since there are infinitely many conjugacy classes of hyperbolic elements in $\mathrm{SL}(2; \mathbb{Z})$ (for example the matrices

$$\left[\begin{array}{cc} n+1 & n \\ 1 & 1 \end{array} \right]$$

for $n > 1, n \in \mathbb{Z}$ are all non-conjugate), there are infinitely many isomorphism classes of discrete groups Γ . Louis Auslander observed that there are infinitely many homotopy classes of compact complete affine 3-manifolds — in contrast to the theorem of Bieberbach that in each dimension there are only finitely many homotopy classes of compact flat Riemannian manifolds. Notice that each of these affine manifolds possesses a parallel Lorentz metric and hence is a flat Lorentz manifold. (Auslander-Markus [9]).

EXERCISE 6.5.12. *Express the complete affine structures on the 2-torus as mapping tori of affine automorphisms of the complete affine manifold \mathbb{R}/\mathbb{Z} .*

CHAPTER 7

Classification

Given a topology Σ and a geometry (G, X) , how does one determine the various ways of putting (G, X) -structures on Σ ? This chapter discusses how to organize the geometric structures on a fixed topology. This is the general *classification problem* for (G, X) -structures.

7.1. Marking geometric structures

We begin with two more familiar and classical cases:

- The moduli space of flat tori;
- The classification of marked Riemann surfaces by Teichmüller space.

The latter is only analogous to our classification problem, but plays an important role, both historically and technically, in the study of locally homogeneous structures.

7.1.1. Marked Riemann surfaces. The prototype of this classification problem is the classification of Riemann surfaces of genus g . The *Riemann moduli space* is a space \mathfrak{M}_g whose points correspond to the biholomorphism classes of genus g Riemann surfaces. It admits the structure of a quasiprojective complex algebraic variety. In particular it is a Hausdorff space, with a singular differentiable structure.

In general the set of (G, X) -structures on Σ will not have such a nice structure. The natural space will in general not be Hausdorff, so we must expand our point of view. To this end, we introduce additional structures, called *markings*, such that the marked (G, X) -structures admit a more tractable classification. As before, the prototype for this classification is the Riemann moduli space \mathfrak{M}_g , which can be understood as the quotient of the Teichmüller space \mathfrak{T}_g (comprising equivalence classes of marked Riemann surfaces of genus g) by the *mapping class group* Mod_g .

Here is the classical context for \mathfrak{T}_g and $\text{Mod}_g = \mathfrak{T}_g/\text{Mod}_g$: The fixed topology is a closed orientable surface Σ of genus g . A *marked Riemann surface of genus g* is a pair (M, f) where M is a Riemann surface and $\Sigma \xrightarrow{f} M$ is a diffeomorphism. The *Teichmüller space* is

defined as the set of equivalence classes of marked Riemann surfaces of genus g , where two such marked Riemann surfaces (M, f) , (M', f') are *equivalent* if and only if there is a biholomorphism $M \xrightarrow{\phi} M'$ such that $\phi \circ f$ is isotopic to f' .

EXERCISE 7.1.1. *Fix a Riemann surface M . The mapping class group*

$$\mathbf{Mod}_g := \pi_0(\mathrm{Diff}(\Sigma))$$

acts simply transitively on the set of equivalence classes of marked Riemann surfaces (M, f) . Thus the Riemann moduli space \mathbf{Mod}_g is the quotient of the Teichmüller space \mathfrak{T}_g by the mapping class group \mathbf{Mod}_g .

7.1.2. Moduli of flat tori. Another common classification problem concerns flat tori. Recall (§5.4.1 a *flat torus* is a Euclidean manifold of the form $M^n := \mathbb{R}^n/\Lambda$, where $\Lambda \subset \mathbb{R}^n$ is a lattice. A *marking* of M is just a basis of Λ . Clearly the set of marked flat n -tori is the set of bases of \mathbb{R}^n , which is a torsor for the group $\mathrm{GL}(n, \mathbb{R})$. (The columns (respectively rows) of invertible $n \times n$ matrices are precisely bases of \mathbb{R}^n .)

EXERCISE 7.1.2. *For $M = \mathbb{R}^n/\Lambda$ as above, compute the isometry group (respectively affine automorphism group) of M . Show that two invertible matrices $A, A' \in \mathrm{GL}(n, \mathbb{R})$ define isometric marked flat tori if and only if $A'A^{-1} \in \mathrm{O}(n)$. Show that all flat n -tori are affinely isomorphic.*

The deformation space of marked flat tori identifies with the homogeneous space $\mathrm{GL}(n, \mathbb{R})/\mathrm{O}(n)$. The *mapping class group* $\mathbf{Mod}(T^n)$ of the n -torus T^n identifies with $\mathrm{GL}(n, \mathbb{Z})$, which acts properly on the deformation space $\mathrm{GL}(n, \mathbb{R})/\mathrm{O}(n)$. The *moduli space of flat tori* in dimension n identifies with the biquotient $\mathrm{GL}(n, \mathbb{Z}) \backslash \mathrm{GL}(n, \mathbb{R})/\mathrm{O}(n)$.

7.1.3. Marked geometric manifolds. Now we define the analogous construction for Ehresmann structures. As usual, we choose to work in the smooth category since (G, X) -manifolds carry natural smooth (in fact real analytic) structures, and the tools of differential topology are convenient. However, in general, there are many options, it may be more natural to consider homeomorphisms, or even homotopy equivalences, depending on the context. Since our primary interest in dimension two, where these notions yield equivalent theories, we do not discuss the alternative context.

DEFINITION 7.1.3. *Let Σ be a smooth manifold. A marking of an (G, X) -manifold M (with respect to Σ) is a diffeomorphism $\Sigma \xrightarrow{f} M$.*

A marked (G, X) -manifold is a pair (M, f) where f is a marking of M . Say that two marked (G, X) -manifolds (f, M) and (f', M') are equivalent if and only if a (G, X) -isomorphism $M \xrightarrow{\phi} M'$ exists such that $\phi \circ f \simeq \phi'$.

7.1.4. The infinitesimal approach. More useful for computations is another approach, where geometric structures are defined infinitesimally as structures on vector bundles associated to the tangent bundle. For example, a Euclidean manifold M can be alternatively described as a *Riemannian metric* on M with vanishing curvature tensor. Another example is defining a Riemann surface as a 2-manifold together with an *almost complex structure*, that is, a complex structure on its tangent bundle. A third example is defining an affine structure as a connection on the tangent bundle with vanishing curvature tensor. Projective structures and conformal structures can be defined in terms of *projective connections* and *conformal connections*, respectively.

In all of these cases, the underlying smooth structure is fixed, and the geometric structure is replaced by an infinitesimal object as above. The diffeomorphism group acts on this space, and the quotient by the full diffeomorphism group would serve as the moduli space. However, to avoid pathological quotient spaces, we prefer to quotient by the identity component of $\text{Diff}(\Sigma)$. Alternatively define the deformation space of marked structures as the quotient of the space of the infinitesimal objects by the subgroup of $\text{Diff}(\Sigma)$ consisting of diffeomorphisms isotopic to the identity.

The “infinitesimal objects” above are *Cartan connections*, to which we refer to Sharpe [249].

7.2. Deformation spaces of geometric structures

Fundamental in the deformation theory of locally homogeneous (Ehresmann) structures is the following principle, first observed in this generality by Thurston [265]:

THEOREM 7.2.1. *Let X be a manifold upon which a Lie group G acts transitively. Let M be a compact (G, X) -manifold with holonomy representation $\pi_1(M) \xrightarrow{\rho} G$.*

- (1) *Suppose that ρ' is sufficiently near ρ in the representation variety $\text{Hom}(\pi_1(M), G)$. Then there exists a (nearby) (G, X) -structure on M with holonomy representation ρ' .*
- (2) *If M' is a (G, X) -manifold near M having the same holonomy ρ , then M' is isomorphic to M by an isomorphism isotopic to the identity.*

Here the topology on marked (G, X) -manifolds is defined in terms of the atlases of coordinate charts, or equivalently in terms of developing maps, or developing sections. In particular one can define a *deformation space* $\text{Def}_{(G,X)}(\Sigma)$ whose points correspond to equivalence classes of marked (G, X) -structures on Σ . One might *like* to say the holonomy map

$$\text{Def}_{(G,X)}(\Sigma) \xrightarrow{\text{hol}} \text{Hom}(\pi_1(\Sigma), G)/\text{Inn}(G)$$

is a local homeomorphism, with respect to the quotient topology on $\text{Hom}(\pi_1(\Sigma), G)/\text{Inn}(G)$ induced from the classical topology on the \mathbb{R} -analytic set $\text{Hom}(\pi_1(\Sigma), G)$. In many cases this is true (see below) but misstated in [127]. However, Kapovich [166] and Baues [21] observed that this is not quite true, because local isotropy groups acting on $\text{Hom}(\pi_1(\Sigma), G)$ may not fix marked structures in the corresponding fibers.

In any case, these ideas have an important consequence:

COROLLARY 7.2.2. *Let M be a closed manifold. The set of holonomy representations of (G, X) -structures on M is open in $\text{Hom}(\pi_1(M), G)$ (with respect to the classical topology).*

One can define a space of flat (G, X) -bundles (defined by a fiber bundle \mathcal{E}_M having X as fiber and G as structure group) and the foliation \mathcal{F} transverse to the fibration $\mathcal{E}_M \rightarrow M$. The foliation \mathcal{F} is equivalent to a reduction of the structure group of the bundle from G with the classical topology to G with the discrete topology. This set of flat (G, X) -bundles over Σ identifies with the quotient of the \mathbb{R} -analytic set $\text{Hom}(\pi_1(\Sigma), G)$ by the action of the group $\text{Inn}(G)$ of inner automorphisms action by left-composition on homomorphisms $\pi_1(\Sigma) \rightarrow G$.

Conversely, if two nearby structures on a compact manifold M have the same holonomy, they are equivalent. The (G, X) -structures are topologized as follows. Let $\Sigma \rightarrow M$ be a marked (G, X) -manifold, that is, a diffeomorphism from a fixed model manifold Σ to a (G, X) -manifold M . Fix a universal covering $\tilde{\Sigma} \rightarrow \Sigma$ and let $\pi = \pi_1(\Sigma)$ be its group of deck transformations. Choose a holonomy homomorphism $\pi \xrightarrow{\rho} G$ and a developing map $\tilde{\Sigma} \xrightarrow{\text{dev}} X$.

In the nicest cases, this means that under the natural topology on flat (G, X) -bundles $(X_\rho, \mathcal{F}_\rho)$ over M , the holonomy map hol is a local homeomorphism. Indeed, for many important cases such as hyperbolic geometry (or when the structures correspond to geodesically complete affine connections), hol is actually an embedding.

7.2.1. Historical remarks. Thurston’s holonomy principle has a long and interesting history.

The first application is the theorem of Weil [281] that the set of *discrete embeddings* of the fundamental group $\pi = \pi_1(\Sigma)$ of a closed surface Σ in $G = \mathrm{PSL}(2, \mathbb{R})$ is open in the quotient space $\mathrm{Hom}(\pi, G)/G$. Indeed, a discrete embedding $\pi \hookrightarrow G$ is exactly a holonomy representation of a *hyperbolic structure* on Σ . The corresponding subset of $\mathrm{Hom}(\pi, G)/G$ is called the *Fricke space* $\mathfrak{F}(\Sigma)$ of Σ . Weil’s results are clearly and carefully expounded in Raghunathan [238], (see Theorem 6.19), and extended in Bergeron-Gelander [41]. Fenchel and Nielsen proved that $\mathfrak{F}(\Sigma) \approx \mathbb{R}^{-\chi(\Sigma)}$; their approach is outlined in §7.4.

For \mathbb{CP}^1 -structures, Theorem 7.2.1 is due to Hejhal [146, 145]; see also Earle [95] and Hubbard [153]. This venerable subject originated with conformal mapping and the work of Schwarz, and closely relates to the theory of second order (Schwarzian) differential equations on Riemann surfaces. In this case, where $X = \mathbb{CP}^1$ and $G = \mathrm{PSL}(2, \mathbb{C})$, we denote the deformation space $\mathrm{Def}_{(G, X)}(\Sigma)$ simply by $\mathbb{CP}^1(\Sigma)$. See Dumas [91] and §14 below.

Thurston sketches the intuitive ideas for Theorem 7.2.1 in his unpublished notes [265], which contains the first explicit statement of this principle. The first detailed proofs of this fact are Lok [202], Canary-Epstein-Green [55], and Goldman [121] (the proof in [121] was worked out with M. Hirsch, and was independently found by A. Haefliger). The ideas in these proofs may be traced to Ehresmann [97], although he didn’t express them in terms of moduli of structures. Corollary 7.2.2 was noted by Koszul [185], Chapter IV, §3, Theorem 3; compare also the discussion in Kapovich [167], Theorem 7.2.

7.3. Representation varieties

As this theorem concerns the topology of the space of holonomy representations, we first discuss the space $\mathrm{Hom}(\pi, G)$ and its quotient $\mathrm{Rep}(\pi, G)$. Good general references for this theory are Kapovich [167], Lubotzky-Magid [204], Raghunathan [238] and Sikora [251].

We shall assume that G is a (real) Lie group and π is finitely generated. Let $\{\gamma_1, \dots, \gamma_N\}$ be a set of generators.

EXERCISE 7.3.1. *Consider the map*

$$\begin{aligned} \mathrm{Hom}(\pi, G) &\longrightarrow G^N \\ \rho &\longmapsto (\rho(\gamma_1), \dots, \rho(\gamma_N)) \end{aligned}$$

- *This map is injective.*

- Its image is an analytic subset of G^N defined by

$$R_\alpha(g_1, \dots, g_N) = 1,$$

where the R_α are the relations among the generators $\gamma_1, \dots, \gamma_N$ of π , regarded as an analytic map $G^N \xrightarrow{R_\alpha} G$.

- Furthermore the structure of this analytic variety is independent of the choice of generating set.
- The natural action of $\text{Aut}(\pi) \times \text{Aut}(G)$ on $\text{Hom}(\pi, G)$ preserves the analytic structure.

In many cases, G may be an *algebraic group*, that is a Zariski-closed subgroup of some $\text{GL}(m, \mathbb{R})$. In that case $\text{Hom}(\pi, G)$ has the structure of a *real algebraic subset* of $\text{GL}(m, \mathbb{R})^N$, and this algebraic structure is preserved by the natural $\text{Aut}(\pi) \times \text{Aut}(G)$ -action. Thus the map of Exercise 7.3.1 embeds $\text{Hom}(\pi, G)$ an analytic or algebraic set.¹ Unless otherwise stated, we give this set the *classical topology* inherited from the topology of G as a Lie group.

EXERCISE 7.3.2. *Suppose that π is ann-generator free group. Let G be a reductive Lie group.*

- Identify $\text{Hom}(\pi, G)$ with the Cartesian power G^n . How does $\text{Aut}(\pi)$ act on G^n ?
- Let $\text{Hom}(\pi, G)^-$ denote the subset comprising ρ such that the centralizer of $\rho(\pi)$ equals the center $\mathcal{Z}(G)$ of G . Show that $\text{Hom}(\pi, G)^-$ is $\text{Inn}(G)$ -invariant, open and dense in $\text{Hom}(\pi, G)$.
- Show that $\text{Inn}(G)$ acts freely and properly on $\text{Hom}(\pi, G)^-$ and the quotient map is a smooth principal $\text{Inn}(G)$ -fibration. Deduce that the quotient space is a real analytic manifold of dimension $(n-1)\dim(G) + \dim(\mathcal{Z}(G))$.

As the holonomy homomorphism $\pi_1(M) \xrightarrow{h} G$ is only defined up to conjugation, it is natural to form the quotient of $\text{Hom}(\pi, G)$ by the subgroup

$$\{1\} \times \text{Inn}(G) < \text{Aut}(\pi) \times \text{Aut}(G)$$

where $\text{Inn}(G) < \text{Aut}(G)$ is the normal subgroup consisting of *inner automorphisms* of G . With this quotient topology inherited from the classical topology on $\text{Hom}(\pi, G)$ as above, we denote this space by $\text{Hom}(\pi, G)/\text{Inn}(G)$ or simply $\text{Hom}(\pi, G)/G$. This quotient space is the one arising in differential geometry/topology as the space of equivalence

¹The Hilbert basis theorem implies that it is not necessary to assume that π has a finite presentation. An interesting question is how the defining ideal varies if π is finitely generated but not finitely presentable.

classes of flat connections, and is the quotient space upon which we concentrate.

Unfortunately this space is generally not well-behaved, and *Murphy's law* applies: *Everything that possibly could go wrong does go wrong*. In particular:

- Although the action of G by conjugation is algebraic/analytic, it is generally neither proper nor free. Thus $\mathbf{Hom}(\pi, G)/G$ is generally not a Hausdorff space.
 - Even if the $\mathbf{Inn}(G)$ -action is proper (for example if G is compact), then the action may not be free, and the quotient may not be a smooth manifold (although it underlies an *orbifold* structure).
- Furthermore the analytic set $\mathbf{Hom}(\pi, G)$ is generally not smooth, and forming the quotient by G can only make matters worse. Sometimes this can be repaired by forming the *algebraic-geometric quotient* (in the sense of *Geometric Invariant Theory*, although then points in this quotient generally do *not* correspond to G -orbits themselves).

The infinitesimal theory, and its relation to cohomology, can be found in Raghunathan [238]. Explicit formulas using the *free differential calculus* of Fox [103] are described in Goldman [130]. See Sikora [167] for a careful treatment of the infinitesimal theory as a scheme.

7.3.1. Example: $\mathbf{SL}(2)$ -characters of \mathbb{F}_2 . (This material is taken from [126], which includes proofs of the stated results.)

A classical theorem of Vogt [280] (and often attributed to Fricke-Klein [104]) asserts that, when $G = \mathbf{SL}(2, \mathbb{C})$ the algebraic-geometric (GIT) quotient of $\mathbf{Hom}(\mathbb{F}_2, G)$ by $\mathbf{Inn}(G)$ is \mathbb{C}^3 . This generalizes the elementary fact that the quotient $G//\mathbf{Inn}(G) \cong \mathbb{C}$, with coordinate given by the trace function

$$G \xrightarrow{\text{tr}} \mathbb{C}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \longmapsto a + d.$$

EXERCISE 7.3.3. *Compute the critical points and the critical values of tr .*

Recall that a function f on G is *regular* if $f(x)$ is a polynomial function of the matrix entries of $x \in G$.

EXERCISE 7.3.4. *Show that for any $\mathbf{Inn}(G)$ -invariant regular function f on G there exists a function $\mathbb{C} \xrightarrow{F} \mathbb{C}$ such that $f = \text{tr} \circ F$.*

Show that $\mathrm{tr}^{-1}(t)$ consists of a single $\mathrm{Inn}(G)$ -orbit when $t \neq \pm 2$. Describe $\mathrm{tr}^{-1}(t)$ when $t = \pm 2$.

It follows that tr is the GIT quotient map for the action of $\mathrm{Inn}(G)$ on G , and we write $G//\mathrm{Inn}(G) \cong \mathbb{C}$.

EXERCISE 7.3.5. Show that if $x \in G$, then $\mathrm{tr}(x) = \mathrm{tr}(x^{-1})$. Deduce that x and x^{-1} are conjugate in G . Is the same true when G is replaced by $\mathrm{SL}(2, \mathbb{R})$ or $\mathrm{GL}(2, \mathbb{R})$?

Writing $\mathbb{F}_2 = \langle X, Y \rangle$ for a pair of free generators X, Y , the identification

$$\begin{aligned} \mathrm{Hom}(\mathbb{F}_2, G) &\longleftrightarrow G \times G \\ \rho &\longleftrightarrow (\rho(X), \rho(Y)) \end{aligned}$$

is equivariant with respect to the action of $G \rightarrow \mathrm{Inn}(G)$ on $\mathrm{Hom}(\mathbb{F}_2, G)$ and the diagonal action of G on $G \times G$ given by:

$$(26) \quad g \cdot (x, y) := (gxg^{-1}, gyg^{-1}).$$

This action preserves the mapping

$$(27) \quad \begin{aligned} G \times G &\longrightarrow \mathbb{C}^3 \\ (x, y) &\longmapsto \begin{bmatrix} \xi := \mathrm{tr}(x) \\ \eta := \mathrm{tr}(y) \\ \zeta := \mathrm{tr}(xy) \end{bmatrix} \end{aligned}$$

which is a GIT-quotient map:

THEOREM 7.3.6 (Vogt [280], Fricke [104]). *Let*

$$\mathrm{SL}(2, \mathbb{C}) \times \mathrm{SL}(2, \mathbb{C}) \xrightarrow{f} \mathbb{C}$$

be a regular function which is invariant under the diagonal action (26) of $\mathrm{SL}(2, \mathbb{C})$ by conjugation. There exists a polynomial function $F(\xi, \eta, \zeta) \in \mathbb{C}[\xi, \eta, \zeta]$ such that

$$f(x, y) = F(\mathrm{tr}(x), \mathrm{tr}(y), \mathrm{tr}(\xi\eta)).$$

Furthermore, for all $(\xi, \eta, \zeta) \in \mathbb{C}^3$, there exists $x, y \in \mathrm{SL}(2, \mathbb{C})$ such that

$$\xi = \mathrm{tr}(x), \eta = \mathrm{tr}(y), \zeta = \mathrm{tr}(xy).$$

Conversely, suppose $x, y, x', y' \in \mathrm{SL}(2, \mathbb{C})$ satisfy

$$\begin{bmatrix} \mathrm{tr}(x) \\ \mathrm{tr}(y) \\ \mathrm{tr}(xy) \end{bmatrix} = \begin{bmatrix} \xi \\ \eta \\ \zeta \end{bmatrix} = \begin{bmatrix} \mathrm{tr}(x') \\ \mathrm{tr}(y') \\ \mathrm{tr}(x'y') \end{bmatrix}$$

where

$$(28) \quad \xi^2 + \eta^2 + \zeta^2 - \xi\eta\zeta \neq 4.$$

Then $(x', y') = g \cdot (x, y)$ for some $g \in G$.

Condition (28) means that the matrix group $\langle x, y \rangle$ acts *irreducibly* on \mathbb{C}^2 . That is, $\langle x, y \rangle$ preserves no proper nonzero linear subspace of \mathbb{C}^2 . The irreducibility condition is crucial in several alternate descriptions of $\mathrm{SL}(2, \mathbb{C})$ -representations of F_2 . In particular, it is equivalent to the condition that the $\mathrm{SL}(2, \mathbb{C})$ -orbit is closed in $\mathrm{Hom}(F_2, \mathrm{SL}(2, \mathbb{C}))$. This condition is in turn equivalent to the orbit being *stable* in the sense of Geometric Invariant Theory. In terms of hyperbolic geometry, it means that the representation fixes no point in \mathbb{CP}^1 .²

7.3.1.1. *Coxeter Extension.* A more geometric description involves the action of the representation ρ with $\rho(X) = x$ and $\rho(Y) = y$ on hyperbolic 3-space \mathbb{H}^3 . The group $\mathrm{PSL}(2, \mathbb{C})$ acts by orientation-preserving isometries of \mathbb{H}^3 . An *involution*, that is, an element $g \in \mathrm{PSL}(2, \mathbb{C})$ having order two, is reflection in a unique geodesic $\mathrm{Fix}(g) \subset \mathbb{H}^3$. Denote the space of such involutions by Inv .

THEOREM 7.3.7 (Coxeter extension). *Suppose that $x, y \in \mathrm{SL}(2, \mathbb{C})$ generate an irreducible representation and let $z = y^{-1}x^{-1}$ so that*

$$xyz = \mathbb{I}.$$

Then there exists a unique triple of involutions

$$\iota_{xy}, \iota_{yz}, \iota_{zx} \in \mathrm{Inv}$$

such that the corresponding elements $P(x), P(y), P(z) \in \mathrm{PSL}(2, \mathbb{C})$ satisfy:

$$\begin{aligned} P(x) &= \iota_{zx}\iota_{xy} \\ P(y) &= \iota_{xy}\iota_{yz} \\ P(z) &= \iota_{yz}\iota_{zx}. \end{aligned}$$

For the proof see [126]

EXERCISE 7.3.8. *Show that Inv identifies with the set of unoriented geodesics in \mathbb{H}^3 . Describe its topological type.*

EXERCISE 7.3.9. *Let $\widetilde{\mathrm{Inv}}$ denote the inverse image $P^{-1}(\mathrm{Inv})$. Show that $\widetilde{\mathrm{Inv}} = \mathfrak{sl}(2, \mathbb{C}) \cap \mathrm{SL}(2, \mathbb{C})$.*

²This situation is remarkably clean; for a description of $\mathrm{SL}(3, \mathbb{C})$, see Lawton [199].

We choose lifts $\tilde{\iota}_{xy}, \tilde{\iota}_{yz}, \tilde{\iota}_{zx} \in \widetilde{\text{Inv}}$ such that

$$\tilde{\iota}_{xy}\tilde{\iota}_{yz}\tilde{\iota}_{zx} = \mathbb{I}.$$

These lifts will be used to parametrize hyperbolic structures on surfaces in terms of traces in $\text{SL}(2, \mathbb{R})$.

7.3.1.2. *Hyperbolic three-holed spheres.* Theorem 7.3.7 implies the Fricke space of hyperbolic structures on the three-holed sphere Σ (sometimes called a “pair of pants” or a “trinion”) identifies with $(-\infty, -2]^3$ using trace coordinates. Namely, the three trace parameters correspond to the three boundary components of Σ . The Coxeter extension identifies a hyperbolic structure on Σ with (perhaps mildly degenerate) right-angled hexagon in the hyperbolic plane \mathbb{H}^2 . Right-angled hexagons are allowed to degenerate when some of the alternate edges covering boundary components degenerate to ideal points.

Suppose that $\xi, \eta, \zeta \leq -2$. Then the corresponding elements $x, y, z \in \text{SL}(2, \mathbb{C})$ have real representatives and are represented by hyperbolic or parabolic elements of $\text{SL}(2, \mathbb{R})$. Furthermore if $\xi, \eta, \zeta < -2$, they are represented by hyperbolic elements of $\text{SL}(2, \mathbb{R})$ whose axes do not intersect. The involutions $\iota_{xy}, \iota_{yz}, \iota_{zx}$ preserve $\mathbb{H}^2 \subset \mathbb{H}^3$ and their restrictions to \mathbb{H}^2 act by (orientation-reversing) reflections in geodesics which we denote by

$$\text{Fix}(\iota_{xy}), \text{Fix}(\iota_{yz}), \text{Fix}(\iota_{zx}) \subset \mathbb{H}^2$$

respectively. Theorem 7.3.7 implies that, for example, the invariant axis of x is the common orthogonal to the lines $\text{Fix}(\iota_{xy}), \text{Fix}(\iota_{zx})$. Their distance equals the distance between their closest points $\text{Fix}(\iota_{xy}) \cap \text{Axis}(x)$ and $\text{Fix}(\iota_{zx}) \cap \text{Axis}(x)$:

$$d(\text{Fix}(\iota_{xy}), \text{Fix}(\iota_{zx})) = d(\text{Fix}(\iota_{xy}) \cap \text{Axis}(x), \text{Fix}(\iota_{zx}) \cap \text{Axis}(x))$$

Since $x = \iota_{zx}\iota_{xy}$, the hyperbolic isometry x is a transvection of displacement

$$\ell_x := 2d(\text{Fix}(\iota_{xy}), \text{Fix}(\iota_{zx}))$$

and the trace of the matrix x equals

$$\xi = -2 \cosh(\ell_x/2).$$

For the detailed proof that $\xi, \eta, \zeta < 2$ implies that the six lines

$$\text{Axis}(x), \text{Fix}(\iota_{xy}), \text{Axis}(y), \text{Fix}(\iota_{yz}), \text{Axis}(z), \text{Fix}(\iota_{zx})$$

bound a convex right-angled hexagon, see §4.3 of [126]. This hexagon is a fundamental domain for the Coxeter group $\langle \iota_{xy}, \iota_{yz}, \iota_{zx} \rangle$. This Coxeter group contains $\langle x, y, z \rangle$ with index two. The union of two adjacent hexagons in the resulting tessellation is then a fundamental domain for $\langle x, y \rangle$. The quotient is a hyperbolic surface homeomorphic

to a three-holed sphere, with three boundary components of length ℓ_x, ℓ_y, ℓ_z .

7.3.2. Twist flows and Fenchel-Nielsen earthquakes. Given a surface Σ and a simple closed curve $\mathcal{C} \subset \Sigma$, we define deformations of representations of $\pi_1(\Sigma)$ which are “supported” on \mathcal{C} . To this end, bordify the complement $\Sigma \setminus \mathcal{C}$ as a surface-with-boundary $\Sigma|_{\mathcal{C}}$ with boundary components \mathcal{C}_i which are identified to form \mathcal{C} in the quotient (which is Σ).

Suppose first that \mathcal{C} separates Σ into two components Σ_1, Σ_2 so that Σ can be reconstructed from the disjoint union

$$\Sigma|_{\mathcal{C}} = \Sigma_1 \bigsqcup \Sigma_2$$

by a quotient map

$$\Sigma_1 \bigsqcup \Sigma_2 \xrightarrow{Q} \Sigma.$$

Write $Q^{-1}(\mathcal{C}) = \mathcal{C}_1 \sqcup \mathcal{C}_2$ where $\mathcal{C}_i \in \Sigma_i$, so that Q identifies \mathcal{C}_1 and \mathcal{C}_2 to form \mathcal{C} .

Choose a basepoint $x_0 \in \mathcal{C} \subset \Sigma$ and let $Q^{-1}(x_0) = \{x_1, x_2\}$ where $x_i \in \mathcal{C}_i \subset \Sigma_i$. Let $c \in \pi_1(\Sigma, x_0)$ be the element corresponding to \mathcal{C} . For $i = 1, 2$, let $c_i \in \pi_1(\Sigma_i, x_i)$, be the respective elements corresponding to c_i . By the Van Kampen theorems, $\pi_1(\Sigma, x_0)$ may be reconstructed from $\pi_1(\Sigma_i, x_i)$ as an *amalgamated free product*

$$\pi_1(\Sigma, x_0) \cong \pi_1(\Sigma_1, x_1) \amalg_{\langle c \rangle} \pi_1(\Sigma_2, x_2).$$

Suppose that $\pi_1(\Sigma) \xrightarrow{\rho} G$ is a representation and \mathfrak{z}_t is a parametrized family of elements of the centralizer of $\rho(c)$ in G . Then we can construct a parametrized family of representations ρ_t by the formula:

$$(29) \quad \rho_t(A) := \begin{cases} \rho(A) & \text{if } A \in \pi_1(\Sigma_1, x_1) \\ \mathfrak{z}_t \rho(A) \mathfrak{z}_t^{-1} & \text{if } A \in \pi_1(\Sigma_2, x_2) \end{cases}$$

Since $\pi_1(\Sigma_1, x_1)$ and $\pi_1(\Sigma_2, x_2)$ generate $\pi_1(\Sigma, x_0)$ and the only relations concern compatibility along c , that \mathfrak{z}_t centralizes $\rho(c)$ implies that (29) defines a family of representations.

EXERCISE 7.3.10. *Develop the analogous construction of a family of representations when \mathcal{C} does not separate Σ , that is, when $\Sigma \setminus \mathcal{C}$ is connected.*

When \mathcal{C} is a simple closed geodesic on a complete hyperbolic surface M , then $\text{dev}(\tilde{\mathcal{C}})$ is a geodesic in \mathbf{H}^2 and the holonomy $\rho(c)$ is a hyperbolic isometry stabilizing $\text{dev}(\tilde{\mathcal{C}})$. The stabilizer $\text{Stab}(\text{dev}(\tilde{\mathcal{C}}))$ is a

one-parameter subgroup of $\mathrm{PSL}(2, \mathbb{R})$ consisting of transvections. The family of representations ρ_t correspond to the following geometric operation: After cutting along \mathcal{C} , re-identify Σ from $\Sigma|_{\mathcal{C}}$ along the isometries corresponding to \mathfrak{z}_t . Thurston generalised this construction (originally due to Fenchel-Nielsen) to *earthquake flows* on Fricke space $\mathfrak{F}(\Sigma)$.

For a more general discussion of earthquakes and an important application, compare Kerckhoff [169]. Thurston's *bending deformations* of embeddings in $\mathrm{PSL}(2, \mathbb{C})$ and the higher-dimensional generalizations due to Johnson-Millson [164] are also special cases of this construction. McMullen [218] and Kamishima-Tan [165] consider a 2-parameter family of deformations in $\mathrm{PSL}(2, \mathbb{C})$ (*quakebend deformations*).

We describe a generalization to $\mathbb{R}\mathbb{P}^2$ -structures in §13.2.2.

EXERCISE 7.3.11. *If $\mathcal{C}_1, \dots, \mathcal{C}_N \subset \Sigma$ are disjoint, with respective centralizing one-parameter subgroups then the corresponding flows on $\mathrm{Hom}(\pi, G)$ commute.*

More generally, suppose $\mathcal{C} = \mathcal{C}_1 \sqcup \dots \sqcup \mathcal{C}_N$ is a *multicurve*, that is, a disjoint union of simple closed curves. Then Exercise 7.3.11 implies the above operation can be performed along each of the curves \mathcal{C}_i independently, obtaining an \mathbb{R}^N -action.

Furthermore every $\mathrm{Inn}(G)$ -invariant function $G \xrightarrow{f} \mathbb{R}$ defines an $\mathrm{Inn}(G)$ -invariant function

$$(30) \quad \begin{aligned} \mathrm{Hom}(\pi, G) &\xrightarrow{fc} \mathbb{R} \\ \rho &\longmapsto \sum_{i=1}^N f \circ \rho(c_i) \end{aligned}$$

where $c_i \in \pi$ corresponds to \mathcal{C}_i .

EXERCISE 7.3.12. *Show that the formula (30) is well-defined, that is, is independent of the elements c_i in the fundamental group.*

7.4. Fenchel-Nielsen coordinates on Fricke space

For a more detailed account, see Abikoff [2], Hubbard [154] or Farb-Margalit [100].

The Fenchel-Nielsen parametrization of $\mathfrak{F}(\Sigma)$ begins with the choice of a *pants decomposition*, that is, a decomposition into three-holed spheres along a multicurve \mathcal{C} as above.

EXERCISE 7.4.1. *Show that if Σ is a closed orientable surface of genus $g > 1$, then $N = 3g - 3$.*

In a sequence of papers, Wolpert [287, 288, 289, 290] initiated the study of the symplectic geometry of $\mathfrak{F}(\Sigma)$. In particular he studied the twist flows and showed they were Hamiltonian flows for geodesic length functions. These were put into the more general context in Goldman [118, 119]. When applied to the geodesic length function of a pants decomposition \mathcal{P} (as in (30)), one obtains a map

$$(31) \quad \mathfrak{F}(\Sigma) \xrightarrow{\ell_{\mathcal{P}}} \mathbb{R}^{+N}.$$

EXERCISE 7.4.2. *This map is a principal \mathbb{R}^N -fibration, where the fiber action is defined by the Fenchel-Nielsen earthquakes along the \mathcal{C}_i . Furthermore, giving $\mathfrak{F}(\Sigma)$ the Weil-Petersson symplectic structure, this \mathbb{R}^N -action is a Hamiltonian action and with momentum mapping (31).*

It follows that the Fenchel-Nielsen earthquake flow defines a *completely integrable Hamiltonian system* ([289]).

More generally Wolpert [290] showed that the length functions ℓ_1, \dots, ℓ_N are part of a *global Darboux coordinate system*

$$(\ell_1, \dots, \ell_N, \tau_1, \dots, \tau_N) \in (\mathbb{R}^+)^N \times \mathbb{R}^N$$

on the symplectic manifold $(\mathfrak{F}(\Sigma), \omega_{WP})$:

$$\omega_{WP} = \sum_{i=1}^N d\ell_i \wedge d\tau_i.$$

The choice of the *twist coordinates* τ_i is not as natural as the length coordinates $\ell_{\mathcal{P}}$: they involve a choice of section s of the mapping $\ell_{\mathcal{P}}$. This section corresponds to when all the $\tau_i = 0$.

The *Fenchel-Nielsen section* arises from the decomposition corresponding to \mathcal{P}] as follows. The hyperbolic surfaces M decomposes into $2g - 2$ pairs-of-pants P_j (where $j = 1, \dots, 2g - 2$). Furthermore ∂P_j consists of closed geodesics

$$\partial P_j = \partial^1 P_j \sqcup \partial^2 P_j \sqcup \partial^3 P_j$$

where each boundary component is one of the \mathcal{C}_i (for $i = 1, \dots, 3g - 3$):

$$\partial^k P_j = \mathcal{C}_{i(j,k)}$$

for $k = 1, 2, 3$.

Each P_j decomposes into two right-angled hexagons $\mathcal{O}_j^+ \cup \mathcal{O}_j^-$; indeed each pants P_j is the *double* of a right-angled hexagon \mathcal{O}_j .

Now fix a collection of lengths

$$\ell = (\ell_1, \dots, \ell_{3g-3}) \in (\mathbb{R}^+)^{3g-3}.$$

The Fenchel-Nielsen section is a *marked* hyperbolic surface with the given length parameters ℓ . Specifically, choose right-angled hexagons $\square_1, \dots, \square_{3g-3}$ with alternate triples of edge-lengths

$$\frac{\ell_i^{(1)}}{2}, \frac{\ell_i^{(2)}}{2}, \frac{\ell_i^{(3)}}{2}$$

for $i = 1, \dots, 3g - 3$.

EXERCISE 7.4.3. Find other sections to $\ell_{\mathcal{P}}$.

EXERCISE 7.4.4. A Dehn twist about a simple closed curve $c \subset \Sigma$ is a homeomorphism $\Sigma \rightarrow \Sigma$ supported on a tubular neighborhood of c . Define a group of homeomorphisms \mathbb{Z}^{3g-3} preserving \mathcal{P} generated by Dehn twists and describe its action on Fenchel-Nielsen coordinates. Describe the action of a Dehn twist about a curve not in \mathcal{P} in Fenchel-Nielsen coordinates.

The following exercises are taken from [119] and will be used in §13.2.2. Choose an Ad -invariant nondegenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ on the Lie algebra \mathfrak{g} of G . Choose an orientation on Σ as well.

EXERCISE 7.4.5. Suppose that $G \xrightarrow{f} \mathbb{R}$ is a smooth $\text{Inn}(G)$ -invariant function. Define a function $G \xrightarrow{F} \mathbb{R}$ by:

$$\langle F(x), Y \rangle = \left. \frac{d}{dt} \right|_{t=0} f((x \exp(tY)))$$

for all $Y \in \mathfrak{g}$.

- Show that F is G -equivariant with respect to the action Inn of inner automorphisms on G and the adjoint representation Ad of G on \mathfrak{g} .
- If $x \in G$, show that $F(x)$ lies in the infinitesimal centralizer of x . In particular the one-parameter subgroup $\exp(tF(x))$ lies in the centralizer $\mathcal{Z}_x < G$ of x .

We call F the *variation function* associated to f .

Now let $c \in \pi_1(M)$ and define a function f_c on $\text{Hom}(\pi, G)$ as in (30) with $N = 1$. Generalizing Wolpert's theorem [288] that the Fenchel-Nielsen earthquake flow is the Hamiltonian flow for the geodesic length function is the following description of the Hamiltonian flow of f_c :

EXERCISE 7.4.6. Suppose that \mathcal{C} is a simple closed curve on M and let c be an element of $\pi_1(M)$ corresponding to \mathcal{C} . Let

$$\mathfrak{z}_t := \exp(tF(\rho(c)))$$

be the corresponding path in the centralizer of $\rho(c)$.

- Suppose first that \mathcal{C} separates Σ into subsurfaces Σ_1, Σ_2 as in §7.3.2. Then (29) describes a flow on $\text{Hom}(\pi, G)$ which leaves the function $f_{\mathcal{C}}$ invariant.
- Suppose that $M|_{\mathcal{C}}$ is connected. Describe the corresponding flow.

In the case of the Fricke component $\mathfrak{F}(\Sigma) \subset \text{Hom}(\pi, G)/G$, every representation ρ with $[\rho] \in \mathfrak{F}(\Sigma)$ has the property that $\rho(c)$ is hyperbolic $\forall c \in \pi \setminus \{1\}$. Denote the open subset of hyperbolic elements of G by Hyp and use the invariant function:

$$(32) \quad \begin{array}{c} \text{Hyp} \xrightarrow{\ell} \mathbb{R}^+ \\ \pm \begin{bmatrix} e^{l/2} & 0 \\ 0 & e^{-l/2} \end{bmatrix} \mapsto l \end{array}$$

EXERCISE 7.4.7. Using the trace form on $\mathfrak{sl}(2, \mathbb{R})$ as the Ad -invariant inner product, show that the corresponding variation function for ℓ is the function

$$\pm \begin{bmatrix} e^{l/2} & 0 \\ 0 & e^{-l/2} \end{bmatrix} \xrightarrow{L} \begin{bmatrix} 1/2 & 0 \\ 0 & -l/2 \end{bmatrix}.$$

The corresponding one-parameter subgroups \mathfrak{z}_t consists of transvections along $\text{Axis}(A)$ displacing points on $\text{Axis}(A)$ by distance t . In particular A itself equals $\mathfrak{z}_{\ell(A)}$.

When c corresponds to a simple closed curve $\mathcal{C} \subset \Sigma$, then ℓ_c is the function on $\mathfrak{F}(\Sigma)$ mapping a marked hyperbolic structure on Σ to the length of the unique closed geodesic homotopic to \mathcal{C} . The corresponding flow is the Fenchel-Nielsen earthquake flow along \mathcal{C} .

7.5. Open manifolds

The classification of (G, X) -structures on open manifolds is quite different than on closed manifolds. Indeed the classification is a relatively elementary special case of Gromov's *h-principle* [138], which extends the Smale-Hirsch theory of immersions. In particular the existence reduces to homotopy theory, and the effective classification uses a weaker equivalence relation. For a simple example, the (G, X) -structures on a disc D^n (for $n > 1$) correspond to immersions $D^2 \looparrowright X$, and the quotient by isotopy is still an infinite-dimensional space. A more suggestive equivalence relation is modeled on *regular homotopy* whereby regular homotopy classes are classified by homotopy classes of sections of a natural fiber bundle. Without extra assumptions — the most notable being completeness — the developing maps are intractable and can be highly pathological.

Constructing incomplete geometric structures on noncompact manifolds M is *easy*. Take any immersion $M \xrightarrow{f} X$ which is not bijective; then f induces an (G, X) -structure on M . If M is parallelizable, then such an immersion always exists (Hirsch [149]).

More generally, let $\pi \xrightarrow{h} G$ be a representation. If the associated flat (G, X) -bundle $E \rightarrow X$ possesses a section $M \xrightarrow{s} E$ whose normal bundle is isomorphic to $\mathbb{T}M$, then an (G, X) -structure exists having holonomy h . This follows from the extremely general *h-principle* of Gromov [138] (see Haefliger [144] or Eliashberg-Mishachev [98], for example).

Here is how it plays out in dimension two. First of all, every orientable noncompact surface admits an immersion into \mathbb{R}^2 and such an




FIGURE 7.1. Immersions of a one-holed torus into the plane and the sphere.




FIGURE 7.2. Development of one-holed torus with holonomy generated by translation

immersion determines an affine structure with trivial holonomy. Immersions can be classified up to crude relation of regular homotopy, although the isotopy classification of immersions of noncompact surfaces seems forbiddingly complicated. Furthermore suppose $\pi \xrightarrow{\mathbf{h}} \text{Aff}(E)$ is a homomorphism such that the character

$$\pi \xrightarrow{\det \circ L \circ \mathbf{h}} \mathbb{Z}/2$$

equals the first Stiefel-Whitney class. That is, suppose its kernel is the subgroup of π corresponding to the orientable double covering of M . Then M admits an affine structure with holonomy \mathbf{h} . Classifying general geometric structures on noncompact manifolds without extra geometric hypotheses seems hopeless under anything but the crudest equivalence relations.

Constructing incomplete geometric structures on compact manifolds is much harder. Indeed for certain geometries (G, X) , there exist closed manifolds for which every (G, X) -structure on M is complete. As a trivial example, if X is compact and M is a closed manifold with finite fundamental group, then Theorem 5.2.2 implies every (G, X) -structure is complete. As a less trivial example, if M is a closed manifold whose fundamental group contains a nilpotent subgroup of finite index and whose first Betti number equals one, then every affine structure on M is complete (see Fried-Goldman-Hirsch [111]). Compare the discussion of Markus's question about the relation of parallel volume to completeness in §11.

CHAPTER 8

Completeness

In many important cases the developing map is a diffeomorphism $\widetilde{M} \rightarrow X$, or at least a covering map onto its image. In particular if $\pi_1(X) = \{e\}$, such structures are *quotient structures*:

$$M \cong \Gamma \backslash X$$

We also call such quotient structures *tame*. This chapter develops criteria for taming the developing map.

Many important geometric structures are modeled on *homogeneous Riemannian manifolds*. These structures determine Riemannian structures, which are locally homogeneous *metric spaces*. For these structures, completeness of the metric space will tame the developing map.

Although it is not completely necessary, this closely relates to *geodesic completeness* of the associated Levi-Civita connection. The key tool is the *Hopf-Rinow theorem*: Geodesic completeness (of the Levi-Civita connection) is equivalent to completeness of the associated metric space. In particular compact Riemannian manifolds are geodesically complete. Many Ehresmann structures have natural Riemannian structures whose completeness tames of the developing map. In particular such structures are quotient structures as above.

After giving some general remarks on the developing map, its relation to the exponential map (for affine connections), we describe all the complete affine structures on T^2 . The chapter ends with a discussion of incomplete affine structures on T^2 , and a general discussion of the most important incomplete examples — *Hopf manifolds*, which were introduced in §6.4 of Chapter 6.

8.1. Locally homogeneous Riemannian manifolds

Suppose (G, X) is a *Riemannian homogeneous space*, that is, X possesses a G -invariant Riemannian metric \mathbf{g}_X . Equivalently, $X = G/H$ where the isotropy group H is compact. Precisely, the image of the adjoint representation $\text{Ad}(H) \subset \text{GL}(\mathfrak{g})$ is compact.

EXERCISE 8.1.1. *Prove that these two conditions on the homogeneous space (G, X) are equivalent.*

If (G, X) is a Riemannian homogeneous space, then every (G, X) -manifold M inherits a Riemannian metric locally isometric to \mathfrak{g}_X . We say that M is *complete* if such a metric is geodesically complete.

EXERCISE 8.1.2. *Prove that this notion of completeness is independent of the G -invariant Riemannian structure \mathfrak{g}_X on X .*

8.1.1. Complete locally homogeneous Riemannian manifolds. We use the following consequence of the *Hopf-Rinow theorem* from Riemannian geometry: *Geodesic completeness* of a Riemannian structure (the complete extendability of geodesics) is equivalent to the *completeness* of the corresponding metric space (convergence of Cauchy sequences). (Compare do Carmo [87], Kobayashi-Nomizu [181], Lee [200], Milnor [222], O’Neill [230], or Papadopoulos [232].) Our application to geometric structures is that *a local isometry from a complete Riemannian manifold is a covering space*.

Recall our standard notation from Chapter 5: M is a (G, X) -manifold with universal covering space $\widetilde{M} \xrightarrow{\Pi} M$; denote by π the associated fundamental group, and (dev, hol) a development pair.

PROPOSITION 8.1.3. *Let (G, X) be a Riemannian homogeneous space. Suppose that X is simply connected. Let M be a complete (G, X) -manifold. Then:*

- $\widetilde{M} \xrightarrow{\text{dev}} X$ is a diffeomorphism;
- $\pi \xrightarrow{\text{hol}} G$ is an isomorphism of π onto a cocompact discrete subgroup $\Gamma \subset G$.

COROLLARY 8.1.4. *Let (G, X) be a Riemannian homogeneous space, where X is simply connected, and let M be a compact (G, X) -manifold. Then the holonomy group $\Gamma \subset G$ is a discrete subgroup which acts properly and freely on X and M is isomorphic to the quotient X/Γ .*

PROOF OF COROLLARY 8.1.4 ASSUMING PROPOSITION 8.1.3. Since (G, X) is a Riemannian homogeneous, M inherits a Riemannian structure locally isometric to X . Since M is compact, this Riemannian structure is complete. Now apply Proposition 8.1.3. \square

PROOF OF PROPOSITION 8.1.3. The Riemannian metric

$$\widetilde{\mathfrak{g}} = \text{dev}^* \mathfrak{g}_X$$

on \widetilde{M} is invariant under the group of deck transformations $\pi_1(M)$ of \widetilde{M} and hence there is a Riemannian metric \mathfrak{g}_M on M such that $\Pi^* \mathfrak{g}_M = \widetilde{\mathfrak{g}}$.

By assumption the metric \mathbf{g}_M on M is complete and so is the metric $\tilde{\mathbf{g}}$ on \tilde{M} . By construction,

$$(\tilde{M}, \tilde{\mathbf{g}}) \xrightarrow{\text{dev}} (X, \mathbf{g}_X)$$

is a local isometry. A local isometry from a complete Riemannian manifold into a Riemannian manifold is necessarily a covering map (Kobayashi-Nomizu [181]) so dev is a covering map of \tilde{M} onto X . Since X is simply connected, it follows that dev is a diffeomorphism. Let $\Gamma \subset G$ denote the image of h . Since dev is equivariant respecting h , the action of π on X given by h is equivalent to the action of π by deck transformations on \tilde{M} . Thus h is faithful and its image Γ is a discrete subgroup of G acting properly and freely on X . Furthermore dev defines a diffeomorphism

$$M = \tilde{M}/\pi \longrightarrow X/\Gamma.$$

□

When M is compact, more is true: X/Γ is compact (and Hausdorff). Since the fibration $G \longrightarrow G/H = X$ is proper, the homogeneous space $\Gamma \backslash G$ is compact, that is, Γ is cocompact in G .

One may paraphrase the above observation abstractly as follows. Let (G, X) be a Riemannian homogeneous space. Then there is an equivalence of categories:

$$\left\{ \text{Compact } (G, X)\text{-manifolds/maps} \right\} \iff \left\{ \text{Discrete cocompact subgroups of } G \text{ acting freely on } X \right\}$$

where the morphisms in the latter category are inclusions of subgroups composed with inner automorphisms of G . (Equivalences of categories are discussed in §A.3.)

8.1.2. Topological rigidity of complete structures. We say that a (G, X) -manifold M is *complete* if $\tilde{M} \xrightarrow{\text{dev}} X$ is a diffeomorphism.¹ A (G, X) -manifold M is complete if and only if its universal covering \tilde{M} is (G, X) -isomorphic to X , that is, if M is isomorphic to the quotient X/Γ (at least if X is simply connected). Note that if (G, X) is contained in (G', X') in the sense of §5.2.3 and $X \neq X'$, then a complete (G, X) -manifold is *never* complete as an (G', X') -manifold.

Here is an interesting characterization of completeness using elementary properties of developing maps.

¹or a covering map if we don't insist that X be simply connected

EXERCISE 8.1.5. *Let (G, X) be a (not necessarily Riemannian) homogeneous space and X be simply connected. Let M be a closed (G, X) -manifold with developing pair (dev, hol) . Show that M is complete if and only if the holonomy representation $\pi \xrightarrow{\text{hol}} G$ is an isomorphism of π onto a discrete subgroup of G which acts properly and freely on X . Find a counterexample when M is not assumed to be closed.*

8.1.3. Euclidean manifolds. Euclidean structures on closed manifolds provide an important example of this. Namely, \mathbf{E}^n is a Riemannian homogeneous space whose isometry group $\text{Isom}(\mathbf{E}^n)$ acts properly with isotropy group the orthogonal group $\mathbf{O}(n)$. As above, Euclidean structures on closed manifolds identify with lattices $\Gamma \subset \text{Isom}(\mathbf{E}^n)$. This class of geometric structures forms the intersection of flat affine structures and locally homogeneous Riemannian structures.

EXERCISE 8.1.6. *Let \mathbf{E} be a Euclidean space with underlying vector space $\mathbf{V} = \text{Trans}(\mathbf{E})$. Then every closed $(\text{Trans}(\mathbf{E}), \mathbf{E})$ -manifold M is a quotient $\Lambda \backslash \mathbf{V}$, where $\Lambda < \mathbf{V}$ is a lattice, that is, M is a flat torus in the sense of §5.4.1.*

Since $\text{Trans}(\mathbf{E}) < \text{Isom}(\mathbf{E})$, every such structure is a Euclidean structure. Remarkably, every closed Euclidean manifold is finitely covered by a flat torus:

THEOREM 8.1.7 ((Bieberbach)). *Let M^n be a closed Euclidean manifold with affine holonomy group $\Gamma < \text{Isom}(\mathbf{E}^n)$. Then $M^n \cong \Gamma \backslash \mathbf{E}^n$ is complete. Furthermore the translation subgroup $\Gamma \cap \mathbb{R}^n$ is a lattice in \mathbb{R}^n and the quotient projection $\Lambda \backslash \mathbf{E}^n \rightarrow M$ is a finite covering space.*

Euclidean structures identify with the more traditional notion of flat Riemannian structures.

8.2. Affine structures and connections

We have seen that a G -invariant metric on X is a powerful tool in classifying (G, X) -structures. However, without this extra structure, many pathological developing maps may arise, even on closed manifolds. In this section we discuss the notion of completeness for affine structures, for which the lack of an invariant metric leads to fascinating phenomena. The simplest example of a compact incomplete affine structure is a *Hopf manifold*, for which the 1-dimensional case was discussed in §5.4.2 and the general case in §6.4.

Just as Euclidean structures are flat Riemannian structures, general Ehresmann structures can be characterized in terms of more general

differential-geometric objects. Affine structures are then *affine connections* ∇ which are locally equivalent to the affine connection ∇_A on a model affine space A . This is equivalent to the vanishing of both the curvature tensor and the torsion tensor of ∇ . Thus affine structures are flat torsionfree affine connections. Such a connection is the Levi-Civita connection for a Euclidean structure, and the Euclidean structure can be recast as an affine structure with parallel Riemannian structure, as described in §1.4.1.

8.3. Completeness and convexity of affine connections

A more traditional proof of Proposition 8.1.3 uses the theory of *geodesics*. Geodesics are curves with zero acceleration, where *acceleration* of a smooth curve is defined in terms of an *affine connection*, which is just a connection on the tangent bundle of a smooth manifold. Connections appear twice in our applications: first, as Levi-Civita connections for Riemannian homogeneous spaces, and second, for flat affine structures. These contexts meet in the setting of Euclidean manifolds.

After we briefly review the standard theory of affine connections and the geodesic flow, we discuss the theorem of Auslander-Markus characterizing complete affine structures. Then we discuss the closely related notion of *geodesic convexity* and prove Koszul's theorem relating convexity to the developing map.

8.3.1. Review of affine connections. Suppose that M is a smooth manifold with an affine connection ∇ . Let $p \in M$ be a point and $\mathbf{v} \in T_p M$ a tangent vector. Then

$$\exists a, b \in \mathbb{R} \cup \{\pm\infty\}$$

such that

$$-\infty \leq a < 0 < b \leq \infty$$

and a geodesic $\gamma(t)$, defined for $a < t < b$, with $\gamma(0) = p$ and $\gamma'(0) = \mathbf{v}$. We call (p, \mathbf{v}) the *initial conditions*. Furthermore γ is unique in the sense that two such γ agree on their common interval of intersection. We may choose the interval (a, b) to be maximal. When $b = \infty$ (respectively $a = -\infty$), the geodesic is *forwards complete* (respectively *backwards complete*). A geodesic is *complete* if and only if it is both forwards and backwards complete. In that case γ is defined on all of \mathbb{R} . We say (M, ∇) is *geodesically complete* if and only if every geodesic extends to a complete geodesic.

If γ is a geodesic with initial condition $(p, \mathbf{v}) \in TM$, then we write

$$\gamma(t) = \text{Exp}(t\mathbf{v})$$

in light of the uniqueness remarks above. For further clarification, we make the following definition:

DEFINITION 8.3.1. *The exponential domain $\mathcal{E} \subset \mathbb{T}M$ is the largest open subset of $\mathbb{T}M$ upon which Exp is defined. For $p \in M$, write $\mathcal{E}_p := \mathcal{E} \cap \mathbb{T}_p M$ and $\text{Exp}_p := \text{Exp}|_{\mathcal{E}_p}$.*

- \mathcal{E} contains the zero-section $\mathbf{0}_M$ of $\mathbb{T}M$.
- \mathcal{E}_p is *star-shaped* about $\mathbf{0}_p$, that is, if $\mathbf{v} \in \mathcal{E}_p$ and $0 \leq t < 1$, then $t\mathbf{v} \in \mathcal{E}_p$.
- The set of all $t \in \mathbb{R}$ such that $t\mathbf{v} \in \mathcal{E}_p$ is an open interval

$$(a_{\mathbf{v}}, b_{\mathbf{v}}) \subset \mathbb{R} \bigsqcup \{-\infty, +\infty\}$$

containing 0, and

$$\begin{aligned} (a_{\mathbf{v}}, b_{\mathbf{v}}) &\longrightarrow M \\ t &\longmapsto \text{Exp}_p(t\mathbf{v}) \end{aligned}$$

is a maximal geodesic.

- This maximal geodesic is complete if and only if $(a_{\mathbf{v}}, b_{\mathbf{v}}) = (-\infty, +\infty)$.

(M, ∇) is geodesically complete if and only if $\mathcal{E} = \mathbb{T}M$. Then

$$(p, \mathbf{v}) \xrightarrow{\Phi_t} \left(\text{Exp}_p(t\mathbf{v}), \frac{d}{dt} \text{Exp}_p(t\mathbf{v}) \right)$$

defines a *flow* (that is, an additive \mathbb{R} -action) on $\mathbb{T}M$, called the *geodesic flow* of (M, ∇) . The velocity vector

$$\frac{d}{dt} \text{Exp}_p(t\mathbf{v})$$

is the image of \mathbf{v} under parallel translation along the geodesic $\text{Exp}_p|_{[0,t]}$.

EXERCISE 8.3.2. *Suppose that M is connected, and ∇ is an affine connection on M . Let $p \in M$. Then (M, ∇) is complete if and only if $\mathcal{E}_p = \mathbb{T}_p(M)$.*

DEFINITION 8.3.3. *Let (M, ∇) be a manifold with an affine connection, and let $x, y \in M$. Then y is visible from x if and only if a geodesic joins x to y . Equivalently, y lies in the image $\text{Exp}_x(\mathcal{E}_x)$. Evidently y is visible from x if and only if x is visible from y . We say that y is invisible from x if and only if y is not visible from x .*

The following idea will be used later in §11.4 and §12.2.

EXERCISE 8.3.4. Let M be an affine manifold and $p \in M$. Show that the set $M(p)$ of points in M visible from p is open in M . More generally, let M be a projective manifold and $p \in M$. Show that the union $M(p)$ of geodesic segments beginning at p is open in M .

EXERCISE 8.3.5. Let M be a manifold with an affine connection. For each $p \in M$, show that the function

$$\begin{aligned} \mathbb{T}_p(M) &\longrightarrow \mathbb{R}^+ \cup \{\infty\} \\ X_p &\longmapsto \sup \{t \in \mathbb{R} \mid tX_p \in \mathcal{E}_p\} \end{aligned}$$

is lower semicontinuous.

(For a discussion of semicontinuous functions, see §D.)

When the affine connection is *flat*, that is, arises from an affine structure, the exponential map relates to the developing map as follows.

PROPOSITION 8.3.6. Let M be an affine manifold with developing map $M \xrightarrow{\text{dev}} \mathbf{A}$. Let $p \in M$. Then the composition

$$\begin{array}{ccc} & \text{dev} \circ \text{Exp}_p & \\ & \curvearrowright & \\ \mathcal{E}_p & \xrightarrow{\text{Exp}_p} M \xrightarrow{\text{dev}} & \mathbf{A} \end{array}$$

extends (uniquely) to an affine isomorphism $\mathbb{T}_p M \xrightarrow{\mathcal{A}_p} \mathbf{A}$: that is, the following diagram

$$(33) \quad \begin{array}{ccc} \mathcal{E}_p & \hookrightarrow & \mathbb{T}_x M \\ \text{Exp}_p \downarrow & & \downarrow \mathcal{A}_p \\ M & \xrightarrow{\text{dev}} & \mathbf{A} \end{array}$$

commutes.

EXERCISE 8.3.7. Prove Proposition 8.3.6.

EXERCISE 8.3.8. Relate the parallel transport along a path $x \overset{\gamma}{\rightsquigarrow} y$ to the composition

$$\mathbb{T}_x M \xrightarrow{\mathcal{A}_y^{-1} \circ \mathcal{A}_x} \mathbb{T}_y M.$$

8.3.2. Geodesic completeness and the developing map. Recall from Chapter 1 that geodesics — curves in A with zero acceleration — are curves in Euclidean space travelling along straight lines at constant speed. Of course, in affine geometry, the speed doesn't make sense, which is why we prefer to characterize geodesics by acceleration. A fundamental result of Auslander-Markus [8] is that geodesic completeness of affine manifolds is equivalent to the bijectivity of the developing map.

THEOREM 8.3.9 (Auslander-Markus [8]). *Let M be an affine manifold, with a developing map $\widetilde{M} \xrightarrow{\text{dev}} A$. Then dev is an isomorphism if and only if M is geodesically complete. That is, the following two conditions are equivalent:*

- M is a quotient of affine space by a discrete subgroup $\Gamma \subset \text{Aff}(A)$ acting properly on A ;
- A particle on M moving at constant speed in a straight line will continue indefinitely.

Clearly if M is geodesically complete, so is its universal covering \widetilde{M} . Hence we may assume M is simply connected. Let $p \in M$. If M is complete, then Exp_p is defined on *all* of T_pM and

$$\begin{array}{ccc} T_pM & \xrightarrow{(D\text{dev})_p} & T_{\text{dev}(p)}A \\ \text{Exp} \downarrow & & \downarrow \text{Exp} \\ M & \xrightarrow{\text{dev}} & A \end{array}$$

commutes. Since the vertical arrows and the top horizontal arrows are bijective, $M \xrightarrow{\text{dev}} A$ is bijective.

The other direction is a corollary of the following basic result (see also Kobayashi [179], Proposition 4.9, Shima [250], Theorem 8.1):

THEOREM 8.3.10 (Koszul [186]). *Let M be an affine manifold and $p \in M$. Suppose that the domain $\mathcal{E}_p \subset T_pM$ of the exponential map Exp_p is convex. Then $\widetilde{M} \xrightarrow{\text{dev}} A$ is a diffeomorphism of \widetilde{M} onto the open subset*

$$\Omega_p := \text{Exp}_p(\mathcal{E}_p) \subset A$$

PROOF OF THEOREM 8.3.10. Clearly we may assume that M is simply connected, so that $\widetilde{M} \xrightarrow{\text{dev}} A$ is defined.

LEMMA 8.3.11. *The image $\text{Exp}_p(\mathcal{E}_p) = M$.*

PROOF OF LEMMA 8.3.11. $\mathcal{E}_p \subset T_p M$ is open and Exp_p is an open map, so the image $\text{Exp}_p(\mathcal{E}_p)$ is open. Since M is assumed to be connected, we show that $\text{Exp}_p(\mathcal{E}_p)$ is closed.

Let $q \in \overline{\text{Exp}_p(\mathcal{E}_p)} \subset M$. Since M is simply connected, Exp_p maps \mathcal{E}_p bijectively onto $\text{Exp}_p(\mathcal{E}_p)$. Since $q \in \overline{\text{Exp}_p(\mathcal{E}_p)}$, there exists $\mathbf{v} \in T_p M$ such that

$$\lim_{t \rightarrow 1} \text{Exp}_p(t\mathbf{v}) = q.$$

Since the star-shaped open subset $\mathcal{E}_p \subset T_p M$ is convex, $t\mathbf{v} \in \mathcal{E}_p$ for $0 \leq t < 1$. We want to show that $\mathbf{v} \in \mathcal{E}_p$ and $\text{Exp}_p(\mathbf{v}) = q$.

Let $W_{\mathbf{v}} \ni \mathbf{v}$ be a convex open neighborhood of \mathbf{v} in $T_p M$, such that its parallel translate

$$W' := \mathbb{P}_{p,q}(W) \subset T_q M$$

lies in \mathcal{E}_q . Then $\mathcal{E}_q \cap W'$ is nonempty. Furthermore, $\exists t_1 > 0$ so that $\text{Exp}_q(\mathcal{E}_q \cap W')$ contains $\text{Exp}_p(t\mathbf{v})$ for $t_1 \leq t < 1$. Let $p_1 := \text{Exp}_p(t_1\mathbf{v})$ and $\mathbf{v}_1 := \mathbb{P}_{p,p_1}(\mathbf{v})$. Then

$$\text{Exp}_p(t\mathbf{v}) = \text{Exp}_{p_1}((t - t_1)\mathbf{v}_1)$$

for $t_1 \leq t < 1$ extends to $t = 1$, proving that $\mathbf{v} \in \mathcal{E}_p$ and $\text{Exp}_p(\mathbf{v}) = q$ as desired. \square

Now we conclude the proof of Theorem 8.3.10: By the commutativity of (33),

$$\begin{array}{ccc} T_p M & \xrightarrow[\cong]{(D\text{dev})_p} & T_{\text{dev}(p)} A \\ \uparrow \wr & & \uparrow \wr \\ \mathcal{E}_p & \xrightarrow[\cong]{(D\text{dev})_p} & (D\text{dev})_p(\mathcal{E}_p) \\ \text{Exp}_p \downarrow & & \downarrow \mathcal{A}_p \\ M & \xrightarrow[\text{dev}]{} & A \end{array}$$

commutes, where the first vertical arrows are inclusions. By the previous argument (now applied to the subset $\mathcal{E}_p \subset T_p M$) the developing map dev is injective. However, Lemma 8.3.11 implies that $\text{Exp}_p(\mathcal{E}_p) = M$ and thus $\text{dev}(M) = \Omega$. \square

EXERCISE 8.3.12. Find an example of a closed affine manifold M such that for every point $x \in M$, the restriction of dev to the closure $\overline{\text{Exp}_x(\mathcal{E}_x)}$ is not injective.

More properties of the exponential map, including criteria for incompleteness, are discussed in Chapter 12, § 12.3.

8.4. Unipotent holonomy

THEOREM 8.4.1 (Fried-Goldman-Hirsch [111], Theorem 8.4(a)).
Let M be a closed affine manifold whose affine holonomy is unipotent. Then M is complete.

Conversely, under the assumption that the affine holonomy group is a *nilpotent* group, completeness is equivalent to unipotent holonomy. See [111].

PROOF. Choose a basepoint $p_0 \in M$ and let $\widetilde{M} \xrightarrow{\Pi} M$ be the corresponding universal covering space; let \widetilde{p}_0 be a basepoint in \widetilde{M} with $\Pi(\widetilde{p}_0) = p_0$. Choose an origin $\mathbf{0}$ to identify \mathbb{A}^n with the vector space \mathbb{R}^n , and choose a developing map $\widetilde{M} \xrightarrow{\text{dev}} \mathbb{R}^n$ such that $\text{dev}(\widetilde{p}_0) = \mathbf{0}$. Let \mathfrak{h} be the corresponding affine holonomy representation and let $\Gamma := \mathfrak{h}(\pi_1(M))$ the affine holonomy group.

Suppose the linear holonomy group $L(\Gamma) < \text{GL}(n, \mathbb{R})$ is unipotent. Then $L(\Gamma)$ is upper-triangular with respect to some basis of \mathbb{R}^n . That is, $L(\Gamma)$ preserves a complete linear flag

$$0 = F^0 \subset F^1 \subset \dots \subset F^n = \mathbb{R}^n$$

where $\dim(F^k) = k$. Furthermore the induced action on F^k/F^{k-1} is trivial. Thus the restriction $L(\Gamma)|_{F^k}$ preserves a nonzero linear functional $F^k \xrightarrow{l_k} \mathbb{R}$ with kernel F^{k-1} . (See, for example, Humphreys [155].)

This invariant flag determines a family of parallel fields $\mathfrak{F}^k \subset \text{TM}$ of k -planes on M , for each $0 \leq k \leq n$. Evidently each \mathfrak{F}^k is integrable, and the leaves are totally geodesic affine submanifolds of M . Furthermore l_k determines parallel 1-forms ω_k on the leaves of \mathfrak{F}^k vanishing on \mathfrak{F}^{k-1} . A partition of unity on M enables the construction of vector fields $\phi_k \in \text{Vec}(M)$ such that:

- $(\phi_k)_p \in \mathfrak{F}^k(p)$ for each $p \in M$,
- $\omega_k(\phi_k) = 1$ on each leaf of \mathfrak{F}^k .

Since M is closed, each ϕ_k integrates to a smooth flow on M . Since $(\phi_k)_p \in \mathfrak{F}^k(p)$, the flow preserves each leaf of \mathfrak{F}^k .

Lift each vector field ϕ_k to a vector field $\widetilde{\phi} \in \text{Vec}(\widetilde{M})$. Then for each $\widetilde{p} \in \widetilde{M}$,

$$(\text{Ddev})_{\widetilde{p}}(\phi_k(\widetilde{p})) \in F^k.$$

Since ϕ_k integrates to a smooth flow on M , its lift $\tilde{\phi} \in \text{Vec}(\widetilde{M})$ integrates to a smooth flow $\tilde{\Phi}_k$ on \widetilde{M} :

$$\begin{aligned} \mathbb{R} \times \widetilde{M} &\longrightarrow \widetilde{M} \\ (t, \tilde{p}) &\longmapsto \tilde{\Phi}_k(t)(\tilde{p}) \end{aligned}$$

Furthermore, $\forall t \in \mathbb{R}, x \in \widetilde{M}$,

$$l_k(\text{dev} \circ \tilde{\Phi}_k(t))(x) = l_k(\text{dev}(x)) + t.$$

We first show that dev is onto; for any $\mathbf{v} \in \mathbb{R}^n \longleftrightarrow A^n$, we find $\tilde{p}_n \in \widetilde{M}$ with $\text{dev}(\tilde{p}_n) = \mathbf{v}$. We proceed, inductively by finding a sequence $\tilde{p}_0, \dots, \tilde{p}_k$ (for $k \leq n$), beginning at the basepoint \tilde{p}_0 , and by flowing \tilde{p}_k along $\tilde{\Phi}_{n-k}$ for time t_{n-k+1} to \tilde{p}_{k+1} , eventually ending at \tilde{p}_n .

To begin the induction (at $k = 0$), define $t_n := l_n(\mathbf{v})$. Then the restriction $\tilde{\Phi}_n|_{[0, t_n]}(\tilde{p}_0)$ is a geodesic path

$$\tilde{p}_0 \rightsquigarrow \tilde{p}_1 := \tilde{\Phi}_n(t_n)(\tilde{p}_0)$$

in \widetilde{M} . The vector $\text{dev}(\tilde{p}_1) \in \mathbb{R}^n$ satisfies $l_n(\text{dev}(\tilde{p}_1)) = t_n = l_n(\mathbf{v})$, and thus lies in $\mathbf{v} + \mathbf{F}^{n-1}$.

Inductively suppose:

- $\tilde{p}_k \in \widetilde{M}$;
- The vector $\text{dev}(\tilde{p}_k)$ lies in $\mathbf{v} + \mathbf{F}^{n-k}$.

Join \tilde{p}_k to $\tilde{p}_{k+1} \in \widetilde{M}$ with $\text{dev}(\tilde{p}_{k+1}) \in \mathbf{v} + \mathbf{F}^{n-k}$. To this end, define

$$t_{n-k} := l_{n-k}(\mathbf{v} - \text{dev}(\tilde{p}_k)).$$

Then the restriction $\tilde{\Phi}_{n-k}|_{[0, t_{n-k}]}(\tilde{p}_k)$ is a geodesic path

$$\tilde{p}_k \rightsquigarrow \tilde{p}_{k+1} := \tilde{\Phi}_{n-k}(t_{n-k})(\tilde{p}_k)$$

Since $l_{n-k}(\text{dev}(\tilde{p}_{k+1}) - \text{dev}(\tilde{p}_k)) = t_{n-k}$,

$$\text{dev}(\tilde{p}_{k+1}) \in \mathbf{v} + \mathbf{F}^{n-k-1}.$$

Continue this induction until $k = n - 1$, when

$$\text{dev}(\tilde{p}_n) \in \mathbf{v} + \mathbf{F}^0 = \{\mathbf{v}\}$$

so $\text{dev}(\tilde{p}_n) = \mathbf{v}$, as claimed.

Next we prove injectivity.

Suppose $[a, b] \xrightarrow{\tilde{\gamma}} \widetilde{M}$ is a path such that $\text{dev}(\tilde{\gamma}(a)) = \text{dev}(\tilde{\gamma}(b))$. We may assume that $\tilde{\gamma}$ is a loop based at \tilde{p}_0 , that is, $\tilde{\gamma}(a) = \tilde{p}_0$, and we continue to assume $\text{dev}(\tilde{p}_0) = \mathbf{0}$. We shall prove that $\tilde{\gamma}(b) = \tilde{p}_0$ by induction on $k = 0, \dots, n - 1$. We assume that $\tilde{\gamma}_k$ is a path

$$\tilde{\gamma}_k(a) = \tilde{p}_0 \rightsquigarrow \tilde{\gamma}_k(b) = \tilde{\gamma}(b)$$

with velocity $(\tilde{\gamma}_k)'(t) \in \mathbb{F}^{n-k}$.

Begin the induction ($k = 0$) with $\tilde{\gamma}_0 = \tilde{\gamma}$. Suppose, inductively, for $k = 0, \dots, n-1$ that $\tilde{\gamma}_k$ is a path as above. Applying the flow $\tilde{\Phi}_k(s)$ to the path $\tilde{\gamma}_k(t)$,

$$l_k \circ \mathbf{dev}(\tilde{\Phi}_k(s)(\tilde{\gamma}_k(t))) = l_k \circ \mathbf{dev}(\tilde{\gamma}_k(t)) + s$$

implies

$$\tilde{\gamma}_{k+1}(t) := \tilde{\Phi}_k(\sigma(t))(\tilde{\gamma}_k(t)),$$

where

$$\sigma(t) := - \int_0^t l_k((\mathbf{dev} \circ \tilde{\gamma}_k)'(s)) ds,$$

is a path satisfying the desired properties. Thus $\tilde{\gamma}$ is relatively homotopic to $\tilde{\gamma}_k$ for $k = 1, \dots, n$. At the final stage ($k = n$),

$$(\tilde{\gamma}_n)'(t) \in \mathbb{F}^0 = 0,$$

$\tilde{\gamma}_n$ is constant, proving that $\tilde{\gamma}(b) = \tilde{p}_0$ as desired. The proof of Theorem 8.4.1 is now complete. \square

Fried [108] proves the following generalization of Theorem 8.4.1:

THEOREM 8.4.2. *Suppose that M is a closed affine manifold whose linear holonomy preserves a flag*

$$0 = \mathbb{F}^0 \subset \mathbb{F}^1 \subset \dots \subset \mathbb{F}^r = \mathbb{R}^n$$

and acts orthogonally on each quotient $\mathbb{F}^s/\mathbb{F}^{s-1}$. Then M is complete.

Indeed, M is finitely covered by a complete affine nilmanifold. The hypothesis is equivalent to the *distality* of the affine holonomy; see [108] for further details.

8.5. Complete affine structures on the 2-torus

The compact complete affine 1-manifold \mathbb{R}/\mathbb{Z} is unique up to affine isomorphism. Its Cartesian square $\mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$ is a Euclidean structure on the two-torus, unique up to affine isomorphism. In this section we shall describe all other complete affine structures on the two-torus and show that they are parametrized by the plane \mathbb{R}^2 .

These structures were first discussed by Kuiper [191]; compare also Baues [23, 22] and Baues-Goldman [191].

THEOREM 8.5.1 (Baues [22]). *The deformation space of marked complete affine structures on T^2 is homeomorphic to \mathbb{R}^2 .*

Indeed, changing the marking corresponds to the action of the *mapping class group* of T^2 , which is naturally isomorphic to $\mathrm{GL}(2, \mathbb{Z})$, on the deformation space. This action identifies with the usual *linear action* of $\mathrm{GL}(2, \mathbb{Z})$ on the vector space \mathbb{R}^2 . The dynamics of this action is very complicated — it is ergodic with respect to Lebesgue measure (which is invariant but infinite) — but the union of its discrete orbits is dense. Its quotient $\mathrm{GL}(2, \mathbb{Z}) \backslash \mathbb{R}^2$ is an intractable non-Hausdorff space. In contrast, $\mathrm{GL}(2, \mathbb{Z})$ acts *properly* on the deformation space of marked Euclidean structures on the torus, which identifies with the homogeneous space $\mathrm{GL}(2, \mathbb{R})/\mathrm{O}(2)$. (Compare §7.1.2.)

We begin by considering the one-parameter family of (quadratic) diffeomorphisms of the affine plane \mathbf{A}^2 defined by

$$\phi_r(x, y) = (x + ry^2, y)$$

Since $\phi_r \circ \phi_s = \phi_{r+s}$, the maps ϕ_r and ϕ_{-r} are mutually inverse. If $\mathbf{u} = (s, t) \in \mathbb{R}^2$ we denote translation by \mathbf{v} as $\mathbf{A} \xrightarrow{\tau_{\mathbf{v}}} \mathbf{A}$. Conjugation of the translation $\tau_{\mathbf{u}}$ by ϕ_r yields the affine transformation

$$\alpha_r(\mathbf{u}) = \phi_r \circ \tau_{\mathbf{u}} \circ \phi_{-r} = \left[\begin{array}{cc|c} 1 & 2rt & s + rt^2 \\ 0 & 1 & t \end{array} \right]$$

and

$$\mathbb{R}^2 \xrightarrow{\alpha_r} \mathrm{Aff}(\mathbf{A})$$

defines a simply transitive affine action. (Compare [110], §1.19.) If $\Lambda \subset \mathbb{R}^2$ is a lattice, then $\mathbf{A}/\alpha_r(\Lambda)$ is a compact complete affine 2-manifold $M = M(r; \Lambda)$ diffeomorphic to a 2-torus.

The parallel 1-form dy defines a parallel 1-form η on M and its cohomology class

$$[\eta] \in H^1(M; \mathbb{R})$$

is a well-defined invariant of the affine structure up to scalar multiplication. In general, M will have no closed geodesics. If $\gamma \subset M$ is a closed geodesic, then it must be a trajectory of the vector field on M arising from the parallel vector field $\partial/\partial x$ on \mathbf{A} ; then γ is closed if and only if the intersection of the lattice $\Lambda \subset \mathbb{R}^2$ with the line $\mathbb{R} \oplus \{0\} \subset \mathbb{R}^2$ is nonzero.

To classify these manifolds, note that the normalizer of $G_r = \alpha_r(\mathbb{R}^2)$ equals

$$\left\{ \left[\begin{array}{cc|c} \mu^2 & a \\ 0 & \mu \end{array} \right] \mid \mu \in \mathbb{R}^\times, a \in \mathbb{R} \right\} \cdot G_r$$

which acts on G_r conjugating

$$\alpha_r(s, t) \mapsto \alpha_r(\mu^2 s + at, \mu t)$$

Let

$$N = \left\{ \begin{bmatrix} \mu^2 & a \\ 0 & \mu \end{bmatrix} \mid \mu \in \mathbb{R}^\times, a \in \mathbb{R} \right\};$$

then the space of affine isomorphism classes of these tori may be identified with the homogeneous space $\mathrm{GL}(2, \mathbb{R})/N$ which is topologically $\mathbb{R}^2 - \{0\}$. The groups G_r are all conjugate and as $r \rightarrow 0$, each representation $\alpha_r|_\pi$ converges to an embedding of π as a lattice of translations $\mathbb{R}^2 \rightarrow \mathbb{R}^2$. It follows that the deformation space of complete affine structures on T^2 form a space which is the union of $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ with a point O (representing the Euclidean structure) which is in the closure of every other structure.

These structures generalize to *left-invariant affine structures* on Lie groups, which form a rich and interesting algebraic theory, which will be discussed in §10. Many (but not all) closed affine 2-manifolds arise from invariant affine structures on T^2 just as many (but not all) projective 1-manifolds arise from invariant projective structures on T^1 (see §5.4).

We briefly summarize this more general point of view, referring to §10.2 for further details.

EXERCISE 8.5.2. *Let \mathfrak{a} be a 2-dimensional commutative associative \mathbb{R} -algebra and let $\Lambda < \mathfrak{a}$ be a lattice.*

- *Adjoin a (two-sided) identity element $\mathbf{1}$ to \mathfrak{a} to define a 3-dimensional commutative associative \mathbb{R} -algebra with unit:*

$$\mathfrak{a}' := \mathfrak{a} \oplus \mathbb{R}\mathbf{1}$$

Let G be the (commutative) group of invertible elements in the multiplicatively closed affine plane $\mathbf{A} := \mathfrak{a} \oplus \mathbf{1}$. Then G acts locally simply transitively (or étale) on \mathbf{A} , and G inherits an invariant affine structure.

- *If \mathfrak{a} is nilpotent, then $\mathfrak{a}^3 = 0$. Then every element of the affine plane \mathbf{A} is invertible and $G = \mathbf{A}$.*
- *Continue to assume that \mathfrak{a} is nilpotent. The quotient Lie group $M := \Lambda \backslash G$ inherits a complete affine structure, and every orientable complete affine 2-manifold arises in this way.*
- *There are two isomorphism classes of nilpotent algebras \mathfrak{a} , depending on whether $\mathfrak{a}^2 = 0$ or $\mathfrak{a}^2 \neq 0$.*

We call complete affine 2-tori M arising from a pair (\mathfrak{a}, Λ) *Euclidean* if $\mathfrak{a}^2 = 0$ (in which case M is a flat (Euclidean) torus; otherwise we call M *non-Riemannian*).




FIGURE 8.1. Tilings corresponding to some complete affine structures on the 2-torus. The second picture depicts a complete non-Riemannian deformation where the affine holonomy contains no nontrivial horizontal translation. The corresponding torus contains no closed geodesics.

EXERCISE 8.5.3. Let $\mathcal{S}_{\mathbb{Q}}^3$ be the rational homology 3-sphere constructed in Exercise 6.2.3.1. Prove that every affine structure on $\mathcal{S}_{\mathbb{Q}}^3$ is complete.

8.6. Complete affine manifolds

This section describes the general theory of complete affine structures; compare §5.4.1 of Chapter 5 and § 8.5 of Chapter 8 for the specific cases of the circle and the two-torus, respectively. The model for the classification is Bieberbach's theorem that every closed *Euclidean* manifold M is finitely covered by a flat torus: that is, M is a quotient of \mathbb{A}^n by a lattice of translations. (An excellent general reference for this classification of Euclidean manifolds and the algebraic theory of their fundamental groups is Charlap [63].)

For complete affine structures on closed manifolds, the conjectural picture replaces the simply transitive group of translations by a more general simply transitive group G of affine transformations, such as the group

$$G = \alpha_r(\mathbb{R}^2) \subset \text{Aff}(\mathbb{A}^2)$$

of § 8.5. This statement had been claimed by Auslander [10] but his proof was flawed. The ideas were clarified in Milnor's wonderful paper [225] and Fried-Goldman [110], which classifies complete affine structures on closed 3-manifolds. (Compare §15.2 of Chapter 15.) Milnor observed that Auslander's claim was equivalent to the *amenability* of the fundamental group. He asked whether the fundamental group of any complete manifold (possibly noncompact) must be amenable; this is equivalent by Tits's theorem [268] to whether a two-generator free group can act properly and affinely.

In the late 1970's, Margulis proved that such group actions do exist; compare §15.3 of Chapter 15.

8.6.1. The Bieberbach theorems. In 1911-1912 Bieberbach found a general group-theoretic criterion for such groups in arbitrary dimension. In modern parlance, Γ is a *lattice* in $\text{Isom}(\mathbb{E}^n)$, that is, a discrete cocompact subgroup. Furthermore $\text{Isom}(\mathbb{E}^n)$ decomposes as a semidirect product $\mathbb{R}^n \rtimes \text{O}(n)$ where \mathbb{R}^n is the vector space of *translations*. In particular every isometry g is a composition of a translation $x \mapsto x + \mathbf{b}$ by a vector $\mathbf{b} \in \mathbb{R}^n$, with an orthogonal linear map $A \in \text{O}(n)$:

$$(34) \quad x \xrightarrow{g} Ax + \mathbf{b}$$

We call A the *linear part* of g and denote it $L(g)$ and \mathbf{b} the *translational part* of g , and denote it $u(g)$. When A is only required to be linear,

then g is *affine*. An affine automorphism is a Euclidean isometry if and only if its linear part lies in $O(n)$.

Bieberbach showed:

- $\Gamma \cap \mathbb{R}^n$ is a lattice $\Lambda \subset \mathbb{R}^n$;
- The quotient Γ/Λ is a finite group, mapped isomorphically into $O(n)$.
- Any isomorphism $\Gamma_1 \rightarrow \Gamma_2$ between Euclidean crystallographic groups $\Gamma_1, \Gamma_2 \subset \text{Isom}(\mathbb{E}^n)$ is induced by an *affine automorphism* $\mathbb{E}^n \rightarrow \mathbb{E}^n$.
- There are only finitely many isomorphism classes of crystallographic subgroups of $\text{Isom}(\mathbb{E}^n)$.

A *Euclidean manifold* is a flat Riemannian manifold, that is, a Riemannian manifold of zero curvature. A Euclidean manifold is *complete* if the underlying metric space is complete, which by the Hopf-Rinow theorem, is equivalent to the condition of geodesic completeness.

A torsionfree Euclidean crystallographic group $\Gamma \subset \text{Isom}(\mathbb{E}^n)$ acts freely on \mathbb{E}^n and the quotient \mathbb{E}^n/Γ is a complete Euclidean manifold. Conversely every complete Euclidean manifold is a quotient of \mathbb{E}^n by a crystallographic group. The geometric version of Bieberbach's theorems is:

- Every compact complete Euclidean manifold is a quotient of a flat torus \mathbb{E}^n/Λ (where $\Lambda \subset \mathbb{R}^n$ is a lattice of translations by a finite group of isometries acting freely on \mathbb{E}^n/Λ).
- Any homotopy equivalence $M_1 \rightarrow M_2$ of compact complete Euclidean manifolds is homotopic to an affine diffeomorphism.
- There are only finitely many affine isomorphism classes of compact complete Euclidean manifolds in each dimension n .

8.6.2. Complete affine solvmanifolds. This gives a very satisfactory qualitative picture of compact Euclidean manifolds, or, (essentially) equivalently Euclidean crystallographic groups. Does a similar picture hold for *affine crystallographic groups*, that is, for discrete subgroups $\Gamma \subset \text{Aff}(\mathbb{A}^n)$ which act properly on \mathbb{A}^n ?

Auslander and Markus [9] constructed examples of *flat Lorentzian crystallographic groups* Γ in dimension 3, for which all three Bieberbach theorems directly fail. In their examples, the quotient $M^3 = \mathbb{A}^3/\Gamma$ is a flat Lorentzian manifold. Topologically these are all 2-torus bundles over S^1 ; conversely every torus bundle over the circle admits such a structure.

These examples arise from a more general construction: namely, Γ embeds as a lattice in a closed Lie subgroup $G \subset \text{Aff}(\mathbf{A})$ whose identity component G^0 acts *simply transitively* on \mathbf{A} .

Furthermore $\Gamma^0 := \Gamma \cap G^0$ has finite index in Γ , so the flat Lorentz manifold M^3 is finitely covered by the homogeneous space G^0/Γ^0 . Necessarily G^0 is simply connected solvable. The group G^0 plays the role of the translation group \mathbb{R}^n and G is called the *crystallographic hull* in Fried-Goldman [110].

A weaker version of this construction is the *syndetic hull*, defined in [110], but known to H. Zassenhaus, H. C. Wang and L. Auslander. (A good general reference for this theory is Raghunathan [238]).

If $\Gamma \subset \text{GL}(n)$ is a solvable group, then a *syndetic hull* for Γ is a subgroup G such that:

- $\Gamma \subset G \subset A(\Gamma)$, where $A(\Gamma) \subset \text{GL}(n)$ is the Zariski closure (algebraic hull) of Γ in $\text{GL}(n)$;
- G is a closed subgroup having finitely many connected components;
- G/Γ is compact (although not necessarily Hausdorff).

The last condition is somewhat called *syndetic*, since “cocompact” sometimes refers to a subgroup whose coset space is compact *and Hausdorff*. (This terminology is due to Gottschalk and Hedlund [135].) Equivalently, $\Gamma \subset G$ is syndetic if and only if $\exists K \subset G$ which is compact and meets every left coset $g\Gamma$, for $g \in G$. (Compare §A.2.)

EXERCISE 8.6.1. *Find an example of an affine crystallographic group with infinitely many syndetic hulls.*

If $M = \Gamma \backslash \mathbf{A}$ is a complete affine manifold, then $\Gamma \subset \text{Aff}(\mathbf{A})$ is a discrete subgroup acting properly and freely on \mathbf{A} . However, in the example above, $\langle A \rangle$ is a discrete subgroup which doesn’t act properly. A proper action of a discrete group is the usual notion of a *properly discontinuous action*. If the action is also free (that is, no fixed points), then the quotient is a (Hausdorff) smooth manifold, and the quotient map $\mathbf{A} \rightarrow \Gamma \backslash \mathbf{A}$ is a covering space. A properly discontinuous action whose quotient is compact as well as Hausdorff is said to be *crystallographic*, in analogy with the classical notion of a *crystallographic group*: A *Euclidean crystallographic group* is a discrete cocompact group of Euclidean isometries. Its quotient space is a Euclidean orbifold. Since such groups act isometrically on metric spaces, discreteness here does imply properness; this dramatically fails for more general discrete groups of *affine transformations*.

L. Auslander [10] claimed to prove that the Euler characteristic vanishes for a compact complete affine manifold, but his proof was flawed.

It rested upon the following question, which in [110], was demoted to a “conjecture,” and is now known as the “Auslander Conjecture”:

CONJECTURE 8.6.2. *Let M be a compact complete affine manifold. Then $\pi_1(M)$ is virtually polycyclic.*

In that case the affine holonomy group $\Gamma \cong \pi_1(M)$ embeds in a closed Lie subgroup $G \subset \text{Aff}(\mathbf{A})$ satisfying:

- G has finitely many connected components;
- The identity component G^0 acts simply transitively on \mathbf{A} .

Then $M = \Gamma \backslash \mathbf{A}$ admits a finite covering space $M^0 := \Gamma^0 \backslash \mathbf{A}$ where

$$\Gamma^0 := \Gamma \cap G^0.$$

The simply transitive action of G^0 define a complete *left-invariant affine structure* on G^0 and the developing map is just the evaluation map of this action. Necessarily G^0 is a 1-connected solvable Lie group and M^0 is affinely isomorphic to the *complete affine solvmanifold* $\Gamma^0 \backslash G^0$. In particular $\chi(M^0) = 0$ and thus $\chi(M) = 0$.

This theorem is the natural extension of Bieberbach’s theorems describing the structure of flat Riemannian (or Euclidean) manifolds; see Milnor [224] for an exposition of this theory and its historical importance. Every flat Riemannian manifold is finitely covered by a *flat torus*, the quotient of \mathbf{A} by a lattice of translations. In the more general case, G^0 plays the role of the group of translations of an affine space and the solvmanifold M^0 plays the role of the flat torus. The importance of Conjecture 8.6.2 is that it would provide a detailed and computable structure theory for compact complete affine manifolds.

Fried-Goldman [110] established Conjecture 8.6.2 in dimension 3. The proof involves classifying the possible Zariski closures $A(\mathbf{L}(\Gamma))$ of the linear holonomy group inside $\text{GL}(\mathbf{A})$. Goldman-Kamishima prove Conjecture 8.6.2 for flat Lorentz manifolds. Grunewald-Margulis Conjecture 8.6.2 when the Levi component of $\mathbf{L}(\Gamma)$ lies in a real rank-one subgroup of $\text{GL}(\mathbf{A})$. See Tomanov Abels-Margulis-Soifer for further results. The conjecture is now known in all dimensions ≤ 6 (Abels-Margulis-Soifer

Part 3

Affine and projective structures

CHAPTER 9

Affine structures on surfaces and the Euler characteristic

One of our first goals is to classify affine structures on closed 2-manifolds. As noted in §7.5, classification of structures on noncompact manifolds is much different, and reduces to a homotopy-theoretic problem since the equivalence relation is much bigger.

The classification of closed affine 2-manifolds splits into two steps: first is the basic result of Benzécri that a closed surface admits an affine structure if and only if its Euler characteristic vanishes. From this it follows that the affine holonomy group of a closed affine 2-manifold is abelian and the second step uses simple algebraic methods to classify affine structures. The first step, and its generalizations, is the subject of this chapter.

9.1. Benzécri's theorem on affine 2-manifolds

The following result was first proved in [34]. Shortly afterwards, Milnor [221] gave a more general proof, clarifying its homotopic-theoretic nature. For generalizations of Milnor's result, see Benzécri [36], Gromov [137], Sullivan [261] and Smillie [252]. For an interpretation of this inequality in terms of hyperbolic geometry, see [115]. More recent developments are surveyed in [128].

THEOREM 9.1.1 (Benzécri 1955). *Let M be a closed 2-dimensional affine manifold. Then $\chi(M) = 0$.*

PROOF. Replace M by its orientable double covering to assume that M is orientable. By the classification of surfaces, M is diffeomorphic to a closed surface of genus $g \geq 0$. Since a simply connected closed manifold admits no affine structure, (§5.2.4), M cannot be a 2-sphere and hence $g \neq 0$. We assume that $g > 1$ and obtain a contradiction.

9.1.1. The surface as an identification space. Begin with the topological model for M . There exists a decomposition of M along $2g$ simple closed curves $a_1, b_1, \dots, a_g, b_g$ which intersect in a single point

$x_0 \in M$. (Compare Fig. 9.1.) The complement

$$M \setminus \bigcup_{i=1}^g (a_i \cup b_i)$$

is the interior of a $4g$ -gon F with edges

$$a_1^+, a_1^-, b_1^+, b_1^-, \dots, a_g^+, a_g^-, b_g^+, b_g^-.$$

(Compare Fig. 9.2.) There exist maps

$$A_1, B_1, \dots, A_g, B_g \in \pi$$

defining indentifications:

$$A_i(b_i^+) = b_i^-,$$

$$B_i(a_i^+) = a_i^-$$

for a quotient map $F \rightarrow M$. A universal covering space is the quotient space of the product $\pi \times F$ by identifications defined by the generators $A_1, B_1, \dots, A_g, B_g$.

Fix a development pair (dev, h) .




FIGURE 9.1. Decomposing a genus $g = 2$ surface along $2g$ curves into a $4g$ -gon. The single common intersection of the curves is a single point which decomposes into the $4g$ vertices of the polygon.




FIGURE 9.2. Identifying the edges of a $4g$ -gon into a closed surface of genus g . The sides are paired into $2g$ curves, which meet at the single vertex.

9.1.2. The turning number. Let $[t_0, t_1]$ be a closed interval. If $[t_0, t_1] \xrightarrow{f} \mathbb{R}^2$ is a smooth immersion, then its *turning number* $\tau(f)$ is defined as the total angular displacement of its tangent vector (normalized by dividing by 2π). Explicitly, if $f(t) = (x(t), y(t))$, then

$$\tau(f) = \frac{1}{2\pi} \int_{t_1}^{t_2} d \tan^{-1}(y'(t)/x'(t)) = \frac{1}{2\pi} \int_{t_1}^{t_2} \frac{x'(t)y''(t) - x''(t)y'(t)}{x'(t)^2 + y'(t)^2} dt.$$

Extend τ to piecewise smooth immersions as follows. Suppose that $[t_0, t_N] \xrightarrow{f} \mathbb{R}^2$ is an immersion which is smooth on subintervals $[a_i, a_{i+1}]$ where

$$t_0 < t_1 < \cdots < t_{N-1} < t_N.$$

Let $f'_+(t_i) = \lim_{t \rightarrow t_i^+} f'(t)$ and $f'_-(t_i) = \lim_{t \rightarrow t_i^-} f'(t)$ be the two tangent vectors to f at t_i ; define the total turning number of f by:

$$\tau(f) := \tau^{cont}(f) + \tau^{disc}(f)$$

where the *continuous contribution* is:

$$\tau^{cont}(f) := \sum_{i=0}^{N-1} (\tau(f|_{[t_i, t_{i+1}]})$$

and the *discrete contribution* is:

$$\tau^{disc}(f) := \frac{1}{2\pi} \sum_{i=0}^{N-1} \angle(f'_-(t_{i+1}), f'_+(t_{i+1}))$$

where $\angle(v_1, v_2)$ represents the positively measured angle between the vectors v_1, v_2 .

Here are some other elementary properties of τ :

- Denote $-f$ the immersion obtained by reversing the orientation on t :

$$(-f)(t) := f(t_0 + t_N - t)$$

Then $\tau(-f) = -\tau(f)$.

- If $g \in \text{Isom}(\mathbb{E}^2)$ is an orientation-preserving Euclidean isometry, then $\tau(f) = \tau(g \circ f)$.
- If f is an immersion of S^1 , then $\tau(f) \in \mathbb{Z}$.
- Furthermore, if an immersion $\partial D^2 \xrightarrow{f} \mathbb{E}^2$ extends to an orientation-preserving immersion $D^2 \rightarrow \mathbb{E}^2$, then $\tau(f) = 1$.

The *Whitney-Graustein theorem* asserts that immersions $S^1 \xrightarrow{f_i} \mathbb{R}^2$ ($i = 1, 2$) are regularly homotopic if and only if $\tau(f_1) = \tau(f_2)$, which implies the last remark.

EXERCISE 9.1.2. *Suppose that S is a compact oriented surface with boundary components $\partial_1 S, \dots, \partial_k S$. Suppose that $S \xrightarrow{f} \mathbb{E}^2$ is an orientation-preserving immersion. Then*

$$\sum_{i=1}^k \tau(f|_{\partial_i S}) = \chi(S).$$

An elementary property relating turning number to affine transformations is the following:

LEMMA 9.1.3. *Suppose that $[a, b] \xrightarrow{f} \mathbb{R}^2$ is a smooth immersion and $\phi \in \mathbf{Aff}^+(\mathbb{R}^2)$ is an orientation-preserving affine automorphism. Then*

$$|\tau(f) - \tau(\phi \circ f)| < \frac{1}{2}$$

PROOF. If ψ is an orientation-preserving Euclidean isometry, then $\tau(f) = \tau(\psi \circ f)$; by composing ϕ with an isometry we may assume that

$$\begin{aligned} f(a) &= (\phi \circ f)(a) \\ f'(a) &= \lambda(\phi \circ f)'(a) \end{aligned}$$

for $\lambda > 0$. That is,

$$(35) \quad \mathbf{L}(\phi)(f'(a)) = \lambda f'(a).$$

Suppose that $|\tau(f) - \tau(\phi \circ f)| \geq 1/2$. Since for $a \leq t \leq b$, the function

$$|\tau(f|_{[a,t]}) - \tau(\phi \circ f|_{[a,t]})|$$

is a continuous function of t and equals 0 for $t = a$ and is $\geq 1/2$ for $t = b$. The intermediate value theorem implies that there exists $0 < t_0 \leq b$ such that

$$|\tau(f|_{[a,t_0]}) - \tau(\phi \circ f|_{[a,t_0]})| = 1/2.$$

Then the tangent vectors $f'(t_0)$ and $(\phi \circ f)'(t_0)$ have opposite direction, that is, there exists $\mu < 0$ such that

$$(36) \quad \mathbf{L}(\phi)(f'(t_0)) = (\phi \circ f)'(t_0) = \mu f'(t_0).$$

Combining (35) with (36), the linear part $\mathbf{L}(\phi)$ has eigenvalues λ, μ with $\lambda > 0 > \mu$. However ϕ preserves orientation, contradicting $\mathbf{Det}(\mathbf{L}(\phi)) = \lambda\mu < 0$. \square

We apply these ideas to the restriction of the developing map \mathbf{dev} to ∂F . Since $f := \mathbf{dev}|_{\partial F}$ is the restriction of the immersion $\mathbf{dev}|_F$ of the 2-disc,

$$1 = \tau(f) = \tau^{disc}(f) + \tau^{cont}(f)$$

where

$$\begin{aligned} \tau^{cont}(f) &= + \sum_{i=1}^g \tau(\text{dev}|_{a_i^+}) + \tau(\text{dev}|_{a_i^-}) + \tau(\text{dev}|_{b_i^+}) + \tau(\text{dev}|_{b_i^-}) \\ &= \sum_{i=1}^g \tau(\text{dev}|_{a_i^+}) - \tau(h(B_i) \circ \text{dev}|_{a_i^+}) \\ &\quad + \tau(\text{dev}|_{b_i^+}) - \tau(h(A_i) \circ \text{dev}|_{b_i^+}) \end{aligned}$$

since $h(B_i)$ identifies $\text{dev}|_{a_i^+}$ with $-\text{dev}|_{a_i^-}$ and $h(A_i)$ identifies $\text{dev}|_{b_i^+}$ with $-\text{dev}|_{b_i^-}$. By Lemma 9.1.3, each

$$\begin{aligned} |\tau(\text{dev}|_{a_i^+}) - \tau(h(B_i) \circ \text{dev}|_{a_i^+})| &< \frac{1}{2} \\ |\tau(\text{dev}|_{b_i^+}) - \tau(h(A_i) \circ \text{dev}|_{b_i^+})| &< \frac{1}{2} \end{aligned}$$

and thus

$$(37) \quad |\tau^{cont}(f)| < \sum_{i=1}^g \frac{1}{2} + \frac{1}{2} = g$$

Now we estimate the discrete contribution. The j -th vertex of ∂F contributes $1/2\pi$ of the angle

$$\angle(f'_-(t_j), f'_+(t_j)).$$

which is supplementary to the i -th *interior angle* α_j of the polygon ∂F , as measured in the Euclidean metric $\text{dev}^* \mathbf{g}_{\mathbb{E}^2}$.

Let $m_0 \in M$ be the point corresponding to the $4g$ vertices of F . The total angle around m_0 (as measured in the metric $\text{dev}^* \mathbf{g}_{\mathbb{E}^2}$ restricted to the lift of a coordinate patch) equals 2π and we would like to identify this as the sum $\sum_{j=1}^{4g} \alpha_j = 2\pi$. However, the interior angle of the side of ∂F may not equal the corresponding angle in the tangent space of m_0 , since they are related by an element of the holonomy group, which is an affine transformation. Angles will generally *not* be preserved by affine transformations, unless they are multiples of a straight angle π radians. Thus, we assume that the edges emanating from each vertex meet at an angle α_j , which is a multiple of π . (Benzécri considers the case when all of the angles are 0 except one, which equals 2π , as in Figure 9.4.)




FIGURE 9.3. Cell-division of torus where all but one angle at the vertex is 0.




FIGURE 9.4. Doubly periodic tiling of the Euclidean plane with all but one angle at vertex 0.

Then the total cone angle at m_0 (as measured in this local Euclidean metric) equals 2π , that is,

$$\sum_{j=1}^{4g} \alpha_j = 2\pi$$

as desired, and

$$\begin{aligned} \tau^{disc}(f) &= \frac{1}{2\pi} \sum_{j=1}^{4g} \angle(f'_-(t_j), f'_+(t_j)) \\ &= \frac{1}{2\pi} \sum_{j=1}^{4g} (\pi - \alpha_j) = 2g - 1. \end{aligned}$$

Now

$$\tau^{cont}(f) = \tau(f) - \tau^{disc} = 1 - (2g - 1) = 2 - 2g$$

but (37) implies $2g - 2 < g$, that is, $g < 2$ as desired. \square

Benzécri's original proof uses a decomposition where all the sides of F have the same tangent direction at x_0 ; thus all the α_j equal 0 except for one which equals 2π (as in Figure 9.4).

9.1.3. The Milnor-Wood inequality.

Shortly after Benzécri proved the above theorem, Milnor observed that this result follows from a more general theorem on flat vector bundles. Let E be the 2-dimensional oriented vector bundle over M whose total space is the quotient of $\widetilde{M} \times \mathbb{R}^2$ by the diagonal action of π by deck transformations on \widetilde{M} and via Loh on \mathbb{R}^2 , (that is, the *flat vector bundle over M associated to the linear holonomy representation*.) This bundle has a natural flat structure, since the coordinate changes for this bundle are (locally) constant linear maps. Now an oriented \mathbb{R}^2 -bundle ξ over a space M is classified by its *Euler class*

$$\text{Euler}(\xi) \in H^2(M; \mathbb{Z}).$$

For M a closed oriented surface, the orientation defines an isomorphism $H^2(M; \mathbb{Z}) \cong \mathbb{Z}$, and we henceforth identify these groups when the context is clear. If ξ is an oriented \mathbb{R}^2 -bundle over M which admits a flat structure, Milnor [221] showed that

$$|\text{Euler}(\xi)| < g.$$

Furthermore every integer in this range is realized by a flat oriented 2-plane bundle. If M is an affine manifold, then the bundle E is isomorphic to the tangent bundle TM of M and hence has Euler number

$$\text{Euler}(\text{TM}) = 2 - 2g.$$

Thus the only closed orientable surface whose tangent bundle has a flat structure is a torus. Furthermore Milnor showed that any \mathbb{R}^2 -bundle whose Euler number satisfies the above inequality has a flat connection.

Extensions of the Milnor-Wood inequality to higher dimensions have been proved by Benzécri [36], Smillie [252], Sullivan [261], Burger-Iozzi-Wienhard [52] and Bucher-Gelander [48].

9.2. Higher dimensions

The Euler Characteristic in higher dimensions In the early 1950's Chern suggested that in general the Euler characteristic of a compact affine manifold must vanish. Based on the Chern-Weil theory of representing characteristic classes by curvature, several special cases of this conjecture can be solved: if M is a compact *complex* affine manifold, then the Euler characteristic is the top Chern number and hence can be expressed in terms of curvature of the complex linear connection (which is zero). However, in general, for a real vector bundle, only the Pontrjagin classes are polynomials in the curvature — indeed Milnor's examples show that the Euler class *cannot* be expressed as a polynomial in the curvature of a linear connection (although it can be expressed as a polynomial in the curvature of an *orthogonal* connection).

This has been an extremely important impetus for research in this subject.

Deligne-Sullivan [86] proved a strong vanishing theorem for flat *complex* vector bundles. Namely, every flat complex vector bundle ξ over a finite complex M is *virtually trivial*: that is, there exists a finite covering space $\hat{M} \xrightarrow{f} M$ such that $f^*\xi$ is trivial. This immediately implies that $\text{Euler}(\xi) = 0$. Hirsch and Thurston [150] gave a very general criterion for vanishing of the Euler class of flat bundle with amenable holonomy; compare Goldman-Hirsch [131] for an elementary proof in the case of flat vector bundles.

For an ingenious argument proving the vanishing of the Euler characteristic for integral holonomy, see Sullivan [260].

Recently the vanishing of the Euler characterisitic for closed affine manifolds with parallel volume has been proved by Bruno Klingler [175]. He uses the natural geometric structure on the total space of the tangent bundle TM of a compact affine manifold M which he calls a *parahypercomplex structure*. Such a structure is an integrable reduction of the structure group to the split quaternions.

9.2.1. The Chern-Gauss-Bonnet Theorem. Most of the known special cases of the Chern-Sullivan conjecture follow from the Chern's

intrinsic generalization of the Gauss-Bonnet theorem [65] and the Chern-Weil theory of characteristic classes. This includes flat pseudo-Riemannian manifolds, flat complex manifolds, and complete affine manifolds (Kostant-Sullivan). Notable exceptions are Benzécri's theorem for surfaces and the Kobayashi-Vey theorem for hyperbolic affine structures.

Chern's theorem concerns an oriented orthogonal rank n vector bundle $\xi \rightarrow M$ over an oriented closed n -dimensional manifold M . That is, ξ is a smooth \mathbb{R}^n -bundle over M with an orthogonal connection ∇ and an orientation on the fibers. Let

$$\text{Euler}(\xi) \in H^n(M, \mathbb{Z})$$

denote the Euler class of the oriented \mathbb{R}^n -bundle ξ . (Compare Milnor-Stasheff [226] and Steenrod [259].) The orthogonal connection ∇ determines an exterior n -form $\text{Euler}(\nabla)$, the *Euler form* of ∇ on M , such that

$$\int_M \text{Euler}(\nabla) = \text{Euler}(\xi) \cdot [M]$$

where $[M] \in H_n(M, \mathbb{Z})$ denotes the fundamental class of M arising from the orientation. The Euler form is a polynomial expression in the curvature of ∇ and vanishes if ∇ is flat. When ξ is the tangent bundle of M , then

$$\text{Euler}(\xi) \cdot [M] = \chi(M),$$

the Euler characteristic of M .

Milnor [221] showed that, over a closed oriented surface of genus $g > 1$, every oriented \mathbb{R}^2 -bundle E with $|\text{Euler}(\xi)| < g$ admits a flat structure. That is, ξ admits a flat *linear connection* ∇ , but if $\text{Euler}(\xi) \neq 0$, then ∇ cannot be orthogonal.

We summarize some of the ideas in Chern's theorem, referring to Poor [237] (§3.56–3.73, pp. 138–49) for detailed proofs and discussion. According to Poor, this geometric approach is due to Gromoll.

A key point in this proof is the use of the associated principal $\text{SO}(n)$ -bundle over M , which is the bundle of positively oriented orthonormal frames. When $n = 2$, this is the unit tangent bundle of M and is an S^1 -bundle over M . As discussed in Steenrod [259] and Milnor-Stasheff [226], the Euler class is really an invariant of the associated oriented S^{n-1} -bundle. The quotient of the total space by the antipodal map on the fiber is an $\mathbb{R}P^{n-1}$ -bundle, which when $n = 2$, identifies with a sphere bundle itself. (Compare Exercise 9.2.1.)

Let ξ be a smooth oriented real vector bundle of even rank $n = 2m$ over an oriented smooth n -manifold M with an orthogonal connection

∇ . Let $\mathfrak{so}(\xi)$ be the vector bundle to ξ associated to the adjoint representation

$$\mathrm{SO}(2m) \longrightarrow \mathrm{Aut}(\mathfrak{so}(2m)).$$

The *curvature tensor* $R(\nabla)$ is an $\mathfrak{so}(\xi)$ -valued exterior 2-form on M . The *Pfaffian* is an Ad-invariant polynomial mapping $\mathfrak{so}(2m, \mathbb{R}) \xrightarrow{\mathrm{Pfaff}} \mathbb{R}$ such that

$$\mathrm{Det}(A) = \mathrm{Pfaff}(A)^2$$

and Pfaff is a polynomial of degree m . For example, for $m = 1$, and

$$A = \begin{bmatrix} 0 & -a \\ a & 0 \end{bmatrix},$$

$\mathrm{Det}(A) = a^2$ and $\mathrm{Pfaff}(A) = a$. Applying the Pfaffian to $R(\nabla)$ yields an exterior $2m$ -form $\mathrm{Pfaff}(R(\nabla))$ on M^{2m} .

The Euler number of ξ (relative to the orientations of ξ and A can be computed by the *Poincaré-Hopf theorem*: Namely, let η be a section of ξ with isolated zeroes p_1, \dots, p_k . Find an open ball B_i containing p_i and a trivialization

$$E_{B_i} \xrightarrow{\approx} B_i \times \mathbb{R}^{2m}.$$

With respect to this trivialization, the restriction of η to B_i is the graph of a map $B_i \xrightarrow{f_i} \mathbb{R}^{2m}$ where $f_i(x) \neq 0$ if $x \neq p_i$. The degree of the smooth map

$$\begin{aligned} S^{2m-1} &\approx \partial B_i \longrightarrow S^{2m-1} \\ x &\longmapsto \frac{f_i(x)}{\|f_i(x)\|} \end{aligned}$$

is independent of the trivialization, and is called the *Poincaré-Hopf index* $\mathrm{Ind}(\xi, p_i)$ of η at p_i . The *Euler number* of E , defined as

$$\mathrm{Euler}(\xi, \eta) := \sum_{i=1}^k \mathrm{Ind}(\eta, p_i) \in \mathbb{Z}$$

is independent of ξ and the local trivializations of ξ . Compare Bott-Tu [46], Theorem 11.17, p.125.)

The *intrinsic Gauss-Bonnet theorem*, due to Chern [65], states that there is a constant $c_m \in \mathbb{R}$ such that

$$\mathrm{Euler}(\xi) = c_m \int_M \mathrm{Pfaff}(R(\nabla)).$$

In particular if $\xi = \mathrm{T}M$, and $R(\nabla) = 0$, then

$$\chi(M) = \mathrm{Euler}(\mathrm{T}M) = 0.$$

Gromoll's proof (Poor [237], §3.56–3.73, pp. 138–49) begins with a smooth vector field η on M which is nonzero in the complement of a finite subset $Z \subset M$. From an orthogonal connection and η determine a $2m - 1$ -form whose exterior derivative equals $\text{Pfaff}(R(\nabla))$ on M^{2m} . Applying Stokes's theorem on the complement of a small neighborhood of $Z \subset M$ implies Chern's theorem.

9.2.2. Smillie's examples of flat tangent bundles. Two oriented 2-plane bundles over a space M are isomorphic if their Euler classes in $H^2(M; \mathbb{Z})$ are equal. (Compare Milnor-Stasheff [226].) Milnor [221] showed that an oriented 2-plane bundle ξ over an oriented surface of genus $g \geq 0$ admits a flat structure if and only if

$$|\text{Euler}(\xi)| < g.$$

Suppose that ξ is such a bundle which is nontrivial, that is, $\text{Euler}(\xi) \neq 0$. Then ξ admits a connection ∇ with curvature zero.

EXERCISE 9.2.1. *Show that the complexification of such a bundle is trivial.*

EXERCISE 9.2.2. *Suppose that $F \rightarrow \Sigma$ is an oriented S^1 -bundle which admits a free proper action of the cyclic group $\mathbb{Z}/m\mathbb{Z}$ on the fibers so that $F' := F/(\mathbb{Z}/m\mathbb{Z})$ is an oriented S^1 -bundle over Σ . Show that*

$$m|\text{Euler}(F)$$

and

$$\text{Euler}(F') = \text{Euler}(F)/m.$$

Deduce that the Euler number of a flat \mathbb{R}^2 -bundle over Σ is always even.

EXERCISE 9.2.3. *Show that the 3-sphere S^3 admits a flat affine connection (that is, a connection on its tangent bundle $\text{T}M$ with vanishing curvature), but no flat affine connection with vanishing torsion.*

Thus, in general, a manifold can have a flat tangent bundle even if it fails to have a flat affine structure. In this direction, Smillie [254] showed that Chern's conjecture is false if one only requires that the curvature vanishes. We outline his (elementary) argument below.

First, he considers the class of *stably parallelizable manifolds* manifolds, that is, manifolds with stably trivial tangent bundles. Recall that a vector bundle $E \rightarrow M$ is *stably trivial* if the Whitney sum $E \oplus \mathbb{R}_M$ is trivial, where $\mathbb{R}_M \rightarrow M$ denotes the trivial line bundle over M . (Smillie uses the terminology “almost” instead of “stably” although we think that “stable” is more standard.)

EXERCISE 9.2.4. If $\xi \rightarrow M$ is a stably trivial vector bundle, then $\xi = f^*TS^n$, for some map $M \xrightarrow{f} S^n$. Furthermore two stably trivial vector bundles ξ, ξ' are isomorphic if and only if

$$\text{Euler}(\xi) = \text{Euler}(\xi') \in H^n(M; \mathbb{Z}).$$

An oriented 2-plane bundle ξ over a closed oriented surface is stably trivial if and only if its Euler number $\text{Euler}(\xi)$ is even (equivalently, if its second Stiefel-Whitney class $w_2(\xi) = 0$).

EXERCISE 9.2.5. Let M be an orientable n -manifold. Show that the following conditions are equivalent:

- M is stably parallelizable;
- M immerses in \mathbb{R}^{n+1} ;
- For any point, the complement $M \setminus \{x\}$ is parallelizable.

Deduce that the connected sum of two stably parallelizable manifolds is parallelizable.

Smillie constructs a 4-manifold N^4 with $\chi(N) = 4$, and a 6-manifold Q^6 with $\chi(Q) = 8$ such that both TN and TQ have flat structures. He begins with a closed orientable surface Σ_3 of genus 3 and the flat $\text{SL}(2, \mathbb{R})$ -bundle ξ over Σ_3 with $\text{Euler}(\xi) = 2$. (This bundle arises by lifting a Fuchsian representation

$$\pi_1(\Sigma_3) \longrightarrow \text{PSL}(2, \mathbb{R})$$

from $\text{PSL}(2, \mathbb{R})$ to $\text{SL}(2, \mathbb{R})$.) Then ξ is stably trivial and admits a flat structure.

The product 4-plane bundle $\xi \times \xi$ over $\Sigma_3 \times \Sigma_3$ is also stably trivial and admits a flat structure. Furthermore its Euler number

$$\text{Euler}(\xi \times \xi) = 2 + 2 = 4.$$

Let P^4 be a parallelizable 4-manifold and let N be the connected sum of six copies of P with $\Sigma_3 \times \Sigma_3$.

EXERCISE 9.2.6. Prove that $TN \cong f^*(\xi \times \xi)$ for some degree one map

$$N \xrightarrow{f} \Sigma_3 \times \Sigma_3.$$

Deduce that TN admits a flat structure and that $\chi(N) = 4$. Find a similar construction for a 6-manifold Q^6 with flat tangent bundle but $\chi(Q) = 8$. Find, for any even $n \geq 8$, an n -dimensional manifold with flat tangent bundle and positive Euler characteristic.

9.2.3. The Kostant-Sullivan Theorem. In 1960, L. Auslander published a false proof that the Euler characteristic of a closed *complete* affine manifold is zero [10]. Of course, the difficulty is that the Euler characteristic can only be computed as a curvature integral for an *orthogonal connection*.

This difficulty was overcome by a clever trick by Kostant and Sullivan [182] who showed that the Euler characteristic of a compact *complete* affine manifold vanishes.

PROPOSITION 9.2.7 (Kostant-Sullivan [182]). *Let M^{2n} be a compact affine manifold whose affine holonomy group acts freely on A^{2n} . Then $\chi(M) = 0$.*

COROLLARY 9.2.8 (Kostant-Sullivan [182]). *The Euler characteristic of a compact complete affine manifold vanishes.*

The main lemma is the following elementary fact, which the authors attribute to Hirsch:

LEMMA 9.2.9. *Let $\Gamma \subset \text{Aff}(A)$ be a group of affine transformations which acts freely on A . Let G denote the Zariski closure of the linear part $L(\Gamma)$ in $GL(V)$. Then every element $g \in G$ has 1 as an eigenvalue.*

PROOF. First we show that the linear part $L(\gamma)$ has 1 as an eigenvalue for every $\gamma \in \Gamma$. This condition is equivalent to the non-invertibility of $L(\gamma) - \mathbb{I}$. Suppose otherwise; then $L(\gamma) - \mathbb{I}$ is invertible. Writing

$$x \xrightarrow{g} L(g)x + u(g),$$

where the vector $u(g)$ is the translational part ($u(g) = g(0)$) of g , the point

$$p := -(L(g) - \mathbb{I})^{-1}u(g)$$

is fixed by γ , contradicting our hypothesis that g acts freely.

Non-invertibility of $L(\gamma) - \mathbb{I}$ is equivalent to

$$\text{Det}(L(\gamma) - \mathbb{I}) = 0,$$

evidently a polynomial condition on γ . Thus $L(g) - \mathbb{I}$ is non-invertible for every $g \in G$, as desired. \square

PROOF OF PROPOSITION 9.2.7. To show that $\chi(M) = 0$, we find an orthogonal connection ∇ for which the Gauss-Bonnet integrand $\text{Pfaff}(R(\nabla)) = 0$. To this end, observe first that the tangent bundle TM is associated to the linear holonomy $L(\Gamma)$, representation of M , and hence its structure group reduces from $\text{Aff}(A)$ to the algebraic hull G of $L(\Gamma)$. Since M is complete, its affine holonomy group Γ acts

freely and Lemma 9.2.9 implies that every element of G has 1 as an eigenvalue.

Let $K \subset G$ be a maximal compact subgroup of G . (One can take K to be the intersection of G with a suitable conjugate of the orthogonal group $\mathbf{O}(2m) \subset \mathbf{GL}(2m, \mathbb{R})$.) A section of the G/K -bundle associated to the G -bundle corresponding to the tangent bundle always exists (since G/K is contractible), and corresponds to a Riemannian metric on M . Let ∇ be the corresponding Levi-Civita connection. Its curvature $R(\nabla)$ lies in the Lie algebra $\mathfrak{k} \subset \mathfrak{so}(2m)$.

Since every element of G has 1 as an eigenvalue, every element of K has 1 as an eigenvalue, and every element of \mathfrak{k} has 0 as eigenvalue. That is, $\text{Det}(k) = 0$ for every $k \in \mathfrak{k}$. Since

$$\text{Pfaff}(k)^2 = \text{Det}(k) = 0,$$

the Gauss-Bonnet integrand $\text{Pfaff}(R(\nabla)) = 0$. Applying Chern's intrinsic Gauss-Bonnet theorem (§9.2.1), $\chi(M) = 0$. \square

Affine structures on Lie groups and algebras

Many geometric structures on closed manifolds arise from *homogeneous structures* — structures invariant under a transitive Lie group action. In particular, left-invariant structures on Lie groups themselves furnish many examples, as we have already seen in dimensions 1 and 2. A large class of affine structures on the 2-torus T^2 arise from invariant affine structures on the Lie group T^2 — see Baues’s survey [23] for an account of this. For example, the content of Auslander’s approach to classifying affine crystallographic groups is that every compact complete affine manifold arises from a left-invariant complete affine structure on a solvable Lie group G with finitely many connected components and a lattice $\Gamma < G$. Similarly, Dupont’s classification [94] of affine structures on 3-dimensional hyperbolic torus bundles reduces to left-invariant affine structures on the solvable unimodular exponential non-nilpotent Lie group $\text{Isom}^0(\mathbb{E}^{1,1})$.

We emphasize the interplay between the coordinates on a Lie group and the affine coordinates. In particular we find many important additional geometric structures in these affine structures.

Left-invariant objects on a Lie group G of course reduce to algebraic constructions on its Lie algebra \mathfrak{g} . Left-invariant affine structures on G correspond to *left-invariant structures* on \mathfrak{g} , where the bi-invariant structures correspond to compatible *associative* structures on \mathfrak{g} . Specifically, covariant differentiation of vector fields preserves the Lie subalgebra of left-invariant vector fields, leading to a bilinear operation

$$(38) \quad \begin{aligned} \mathfrak{g} \times \mathfrak{g} &\longrightarrow \mathfrak{g} \\ (X, Y) &\longmapsto XY := \nabla_X(Y) \end{aligned}$$

Conversely, any bilinear mapping $\mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g}$ defines a left-invariant connection on G . The connection is flat and torsionfree if and only if the corresponding algebra is *left-symmetric* (In this chapter an *algebra* is a finite-dimensional \mathbb{R} -vector space \mathfrak{a} together with bilinear map $\mathfrak{a} \times \mathfrak{a} \longrightarrow \mathfrak{a}$.) Denote the commutator operation by:

$$(39) \quad [x, y] := xy - yx$$

In general, however a left-invariant structure will *not* be right-invariant, but instead will satisfy the *left-symmetry* condition, that the *associator operation*

$$(40) \quad [x, y, z] := (xy)z - x(yz)$$

is symmetric in its first two arguments:

$$(41) \quad [x, y, z] = [y, x, z]$$

Such algebras arise in many mathematical contexts, and we call them *left-symmetric algebras*. (They also go by other names : *pre-Lie algebras*, *Koszul algebras*, *Vinberg algebras*, or *Koszul-Vinberg algebras*; see Rothaus [241], Dorfmeister [88], Vinberg [278], Koszul [183], and Matsushima [216].)

Their literature is vast. We particularly recommend Burde's survey article [50], as well as works by Vinberg [278], Helmstetter [147] and Vey [275], Medina [219], Segal [247], Kim [171]. and the references cited therein for more information.

We do not attempt to be comprehensive, but mainly give a glimpse into this fascinating algebraic theory, which provides a rich class of interesting geometric examples.

10.1. Étale representations and the developing map

If G is a Lie group and $a \in G$, then define the *left-* and *right-multiplication operations* $\mathfrak{L}_a, \mathfrak{R}_a$, respectively:

$$\mathfrak{L}_a(b) := ab$$

$$\mathfrak{R}_a(b) := ba$$

Suppose that G is a Lie group with an affine structure. The affine structure is *left-invariant* (respectively *right-invariant*) if and only if the operations $G \xrightarrow{\mathfrak{R}_a} G$ (respectively $G \xrightarrow{\mathfrak{L}_a} G$) are affine. An affine structure is *bi-invariant* if and only if it is both left-invariant and right-invariant. In this section we describe the relationship between left-invariant affine structures and étale (locally simply transitive) affine representations.

10.1.1. Reduction to simply-connected groups. Suppose that G is a Lie group with a left-invariant (respectively right-invariant, bi-invariant) affine structure. Let \tilde{G} be its universal covering group and

$$\pi_1(G) \hookrightarrow \tilde{G} \longrightarrow G$$

the corresponding central extension. Then the induced affine structure on \tilde{G} is also left-invariant (respectively right-invariant, bi-invariant).

Conversely, since $\pi_1(G)$ is central in \tilde{G} , every left-invariant (respectively right-invariant, bi-invariant) affine structure on \tilde{G} determines a left-invariant (respectively right-invariant, bi-invariant) affine structure on G . Thus there is a bijection between left-invariant (respectively right-invariant, bi-invariant) affine structures on a Lie group and left-invariant (respectively right-invariant, bi-invariant) affine structures on any covering group. For this reason we shall mainly only consider simply connected Lie groups.

10.1.2. The translational part of the étale representation.

Suppose that G is a simply connected Lie group with a left-invariant affine structure. Let $G \xrightarrow{\text{dev}} \mathbf{A}$ be a developing map. Corresponding to the affine action of G on itself by left-multiplication is a homomorphism $G \xrightarrow{\alpha} \text{Aff}(\mathbf{A})$ such that the diagram

$$(42) \quad \begin{array}{ccc} G & \xrightarrow{\text{dev}} & \mathbf{A} \\ \mathfrak{L}_g \downarrow & & \downarrow \alpha(g) \\ G & \xrightarrow{\text{dev}} & \mathbf{A} \end{array}$$

commutes for each $g \in G$. We may assume that dev maps the identity element $e \in G$ to an origin $p_0 \in \mathbf{A}$. Then (42) implies that the developing map is the translational part of the affine representation:

$$\text{dev}(g) = (\text{dev} \circ \mathfrak{L}_g)(e) = (\alpha(g) \circ \text{dev})(e) = \alpha(g)(p_0)$$

Furthermore since dev is open, it follows that the orbit $\alpha(G)(p_0)$ equals the developing image and is open. Indeed the translational part, which is the differential of the evaluation map

$$\mathbb{T}_e G = \mathfrak{g} \longrightarrow \mathbb{V} = \mathbb{T}_{p_0} \mathbf{A}$$

is a linear isomorphism. Such an action will be called *locally simply transitive* or simply an *étale* representation.

Conversely suppose that $G \xrightarrow{\alpha} \text{Aff}(\mathbf{A})$ is an étale affine representation with an open orbit $\mathfrak{D} \subset \mathbf{A}$. Then for any point $x_0 \in \mathfrak{D}$, the evaluation map

$$g \mapsto \alpha(g)(x_0)$$

defines a developing map for an affine structure on G . Since

$$\begin{aligned} \text{dev}(\mathfrak{L}_g h) &= \alpha(gh)(x_0) \\ &= \alpha(g)\alpha(h)(x_0) \\ &= \alpha(g) \text{dev}(h) \end{aligned}$$

for $g, h \in G$, this affine structure is left-invariant. Thus a left-invariant affine structure on a Lie group G corresponds to correspond precisely to an étale affine representation $G \rightarrow \text{Aff}(A)$ and a choice of open orbit.

Pull back the connection on $\mathfrak{D} \subset A$ by dev to obtain a flat torsion-free affine connection ∇ on G .

Since the affine representation $G \xrightarrow{\alpha} \text{Aff}(A)$ corresponds to left-multiplication, the associated Lie algebra representation, also denoted $\mathfrak{g} \xrightarrow{\alpha} \mathfrak{aff}(A)$, maps \mathfrak{g} into affine vector fields which correspond to the infinitesimal generators of left-multiplications, that is, to *right-invariant vector fields*. Thus with respect to a left-invariant affine structure on a Lie group G , the right-invariant vector fields are affine.

10.2. Two-dimensional commutative associative algebras

Commutative associative algebras provide many examples of affine structures on closed 2-manifolds as follows.

Let \mathfrak{a} be such an algebra and adjoin a two-sided identity element (denoted “ $\mathbf{1}$ ”) to form a new commutative associative algebra $\mathfrak{a} \oplus \mathbb{R}\mathbf{1}$ with identity. The invertible elements in $\mathfrak{a} \oplus \mathbb{R}\mathbf{1}$ form an open subset closed under multiplication. The element $e := 0 \oplus \mathbf{1}$ is two-sided identity element. The universal covering group G of the group of invertible elements of the form

$$a \oplus \mathbf{1} \in \mathfrak{a} \oplus \mathbb{R}\mathbf{1}$$

acts locally simply transitively and affinely on the affine space

$$A = \mathfrak{a} \oplus \{\mathbf{1}\}.$$

The Lie algebra of G naturally identifies with the algebra \mathfrak{a} and there is an exponential map $\mathfrak{a} \xrightarrow{\text{exp}} G$ defined by the usual power series (in \mathfrak{a}). The corresponding evaluation map at e defines a developing map for an invariant affine structure on the vector group \mathfrak{a}

Now let $\Lambda \subset \mathfrak{a}$ be a lattice. The quotient \mathfrak{a}/Λ is a torus with an invariant affine structure. The complete structures were discussed in §8.5. Now we discuss all the structures on \mathbb{R}^2 , in terms of commutative associative algebras. (Recall that for commutative algebras, associativity and left-symmetry are equivalent, see §10.3.1.)

The classification of 2-dimensional commutative associative algebras is not difficult; here we summarize the classification, in terms of a basis $X, Y \in \mathfrak{a}$:

- $\mathfrak{a}^2 = 0$ (all products are zero). The corresponding affine representation is the action of \mathbb{R}^2 on the plane by translation

and the corresponding affine structures on the torus are the Euclidean structures.

- $\dim(\mathfrak{a}^2) = 1$ and \mathfrak{a} is nilpotent. We may take X to be a generator of \mathfrak{a}^2 and $Y \in \mathfrak{a}$ to be an element with $Y^2 = X$. The corresponding affine representation is the simply transitive action discussed in §8.5, which we call the *non-Riemannian* (complete) structures. These are the two isomorphism classes described in §8.5. The corresponding affine structures are complete but non-Riemannian. These structures deform to the first one, where $\mathfrak{a} = \mathbb{R}[x, y]$ where $x^2 = ky$. These are all equivalent when $k \neq 0$, and deforms to the Euclidean structure as $k \rightarrow 0$.
- \mathfrak{a} is a direct sum of 1-dimensional algebras, one with zero multiplication (corresponding to the complete structure), and one with nonzero multiplication (corresponding to the radiant structure). We can choose $X^2 = X$ for the radiant summand and this is the only nonzero basic product.

For various choices of Λ one obtains parallel suspensions of Hopf circles. In these cases the developing image is a halfplane, and we call these structures *nonradiant halfplane structures*.

- The next structure is a radiant suspension of the Euclidean 1-dimensional structure. Take X to be the radiant vector field, so that it is an identity element in \mathfrak{a} . For various choices of Λ one obtains radiant suspensions of the complete affine 1-manifold \mathbb{R}/\mathbb{Z} . The developing image is a halfplane.
- \mathfrak{a} is a direct sum of nonzero 1-dimensional algebras. Taking X, Y to be the generators of these summands, we can assume they are *idempotent*, that is, $X^2 = X, Y^2 = Y$. This structure is radiant since $X + Y$ is an identity element, that is, a radiant vector field. Products of Hopf circles, and, more generally, radiant suspensions of Hopf circles are examples of these affine manifolds.

The developing image is a quadrant in \mathbb{R}^2 .

- Finally $\mathfrak{a} \cong \mathbb{C}$ is the field of complex structures, regarded as an \mathbb{R} -algebra. In this case we obtain the complex affine 1-manifolds, in particular the (usual) 2-dimensional Hopf manifolds are all obtained from this algebra. Clearly $X \leftrightarrow 1 \in \mathbb{C}$ is the identity and these structures are all radiant. The developing image is the complement of a point in the plane.

Baues [23], surveys the theory of affine structures on the 2-torus. In particular he describes how the homogeneous structures (which he calls

“affine Lie groups”) deform one into another. Kuiper [191] classified convex affine structures on T^2 , including the complete case (see Chapter 8, §8.5). Nagano-Yagi [228] and Arrowsmith-Furness [6],[7] completed the classification. The complex-affine structures can be understood easily in terms of nonzero abelian differentials on the underlying elliptic curve (see, for example, Gunning [142, 143] and the discussion in §14. Projective structures on T^2 were classified by Goldman [114] and in higher dimensions by Benoist [28].

Here are the multiplication tables for the 2-dimensional commutative associative algebras:

	X	Y
X	0	0
Y	0	0

	X	Y
X	0	0
Y	0	X

TABLE 1. Multiplication tables for complete (Euclidean and non-Riemannian) structures

	X	Y
X	X	Y
Y	Y	0

	X	Y
X	X	0
Y	0	X

	X	Y
X	X	0
Y	0	Y

TABLE 2. Multiplication tables for incomplete (radiant halfplane, nonradiant halfplane and hyperbolic) structures

10.3. Left-invariant connections and left-symmetric algebras

Let G be a Lie group with Lie algebra \mathfrak{g} , which we realize as *left-invariant vector fields*. Let ∇ be a left-invariant connection; that is, each \mathfrak{L}_g preserves ∇ . If $X, Y \in \mathfrak{g}$ are such left-invariant vector fields, then left-invariance of ∇ implies that the *covariant derivative* $\nabla_X Y$ is also a left-invariant vector field. Thus the operation defined in (38) turns \mathfrak{g} into a finite-dimensional algebra over \mathbb{R} .

The condition that the torsion Tor_∇ of ∇ vanishes is precisely the commutator property (39). Using (39), the condition that ∇ has zero curvature is:

$$(XY - YX)Z = [X, Y]Z = X(YZ) - Y(XZ).$$

This condition is equivalent to the left-symmetric property (41), that is, the *associator* defined in (40) is symmetric in its first two arguments:

$$[X, Y, Z] = [Y, X, Z].$$

An algebra satisfying this condition will be called *left-symmetric*.

Every left-symmetric algebra determines a Lie algebra. This generalizes the well-known fact that underlying every associative algebra is a Lie algebra. We shall sometimes call a left-symmetric algebra with underlying Lie algebra \mathfrak{g} an *affine structure* on the Lie algebra \mathfrak{g} .

EXERCISE 10.3.1. *Let \mathfrak{a} be an algebra with commutator operation $[X, Y] := XY - YX$, and define a trilinear alternating map*

$$\begin{aligned} \mathfrak{a} \times \mathfrak{a} \times \mathfrak{a} &\xrightarrow{\text{Jacobi}} \mathfrak{a}. \\ (X, Y, Z) &\mapsto [[X, Y], Z] + [[Z, X], Y] + [[Y, Z], X] \end{aligned}$$

Show that

$$\begin{aligned} \text{Jacobi}(X, Y, Z) &= [X, Y, Z] + [Y, Z, X] + [Z, X, Y] \\ &\quad - [Z, Y, X] - [X, Z, Y] - [Y, X, Z] \end{aligned}$$

where $[X, Y, Z]$ denotes the associator defined in (40). Deduce that underlying every left-symmetric algebra is a Lie algebra.

EXERCISE 10.3.2. *Find an algebra \mathfrak{a} which is not left-symmetric but its commutator nonetheless satisfies the Jacobi identity.*

In terms of left-multiplication and the commutator operation defined in (39), a condition equivalent to (41) is:

$$(43) \quad \mathfrak{L}([X, Y]) = [\mathfrak{L}(X), \mathfrak{L}(Y)]$$

that is, that $\mathfrak{g} \xrightarrow{\mathfrak{L}} \text{End}(\mathfrak{a})$ is a Lie algebra homomorphism. We denote by $\mathfrak{a}_{\mathfrak{L}}$ the corresponding \mathfrak{g} -module. Furthermore the identity map $\mathfrak{g} \xrightarrow{I} \mathfrak{a}_{\mathfrak{L}}$ defines a cocycle of the Lie algebra \mathfrak{g} with coefficients in the \mathfrak{g} -module $\mathfrak{a}_{\mathfrak{L}}$:

$$(44) \quad \mathfrak{L}(X)Y - \mathfrak{L}(Y)X = [X, Y]$$

Let A denote an affine space with associated vector space \mathfrak{a} ; then it follows from (43) and (44) that the map $\mathfrak{g} \xrightarrow{\alpha} \text{aff}(A)$ defined by

$$(45) \quad Y \mapsto \xrightarrow{\alpha(X)} \mathfrak{L}(X)Y + X$$

is a Lie algebra homomorphism.

THEOREM 10.3.3. *There is an isomorphism of categories between left-symmetric algebras and simply connected Lie groups with left-invariant affine structure. Under this isomorphism the associative algebras correspond to bi-invariant affine structures.*

We have proved the first assertion, and now prove the second assertion.

10.3.1. Bi-invariance and associativity. Under the correspondence between left-invariant affine structures on G and left-symmetric algebras \mathfrak{a} , *bi-invariant* affine structures corresponds to *associative* algebras \mathfrak{a} .

PROPOSITION 10.3.4. *Let \mathfrak{a} be the left-symmetric algebra corresponding to a left-invariant affine structure on G . Then \mathfrak{a} is associative if and only if the left-invariant affine structure is bi-invariant.*

PROOF. Let ∇ be the affine connection corresponding to a left-invariant affine structure on G . Then ∇ is left-invariant and defines the structure of a left-symmetric algebra \mathfrak{a} on the Lie algebra of left-invariant vector fields on G .

Suppose that the affine structure is bi-invariant; then ∇ is also right-invariant. Therefore right-multiplications on G are affine maps with respect to the affine structure on G . It follows that the infinitesimal right-multiplications — the left-invariant vector fields — are affine vector fields. For a flat torsionfree affine connection a vector field Z is affine if and only if the second covariant differential $\nabla\nabla Z$ vanishes. Now $\nabla\nabla Z$ is the tensor field which associates to a pair of vector fields X, Y the vector field

$$\begin{aligned}\nabla\nabla Z(X, Y) &:= \nabla_X(\nabla Z(Y)) - \nabla Z(\nabla_X Y) \\ &= \nabla_X(\nabla_Y Z) - \nabla_{\nabla_X Y}(Z).\end{aligned}$$

If X, Y, Z are left-invariant vector fields, then

$$\nabla\nabla Z(X, Y) = X(YZ) - (XY)Z = [X, Y, Z]$$

in \mathfrak{a} , so \mathfrak{a} is an associative algebra, as desired.

Conversely, suppose \mathfrak{a} is an associative algebra. We construct from \mathfrak{a} a Lie group $G = G(\mathfrak{a})$ with a bi-invariant structure, such that the corresponding left-symmetric algebra equals \mathfrak{a} .

Denote by $\mathbb{R}\mathbf{1}$ a 1-dimensional algebra (isomorphic to \mathbb{R} generated by a two-sided identity element $\mathbf{1}$). The direct sum $\mathfrak{a} \oplus \mathbb{R}\mathbf{1}$ admits an associative algebra structure where $\mathbf{1}$ is a two-sided identity element:

$$(a \oplus a_0\mathbf{1})(b \oplus b_0\mathbf{1}) := (ab + a_0b + ab_0) \oplus a_0b_0\mathbf{1},$$

that is, “adjoint to \mathfrak{a} a two-sided identity element.” The affine hyperplane $\mathfrak{a} \oplus \mathbf{1}$ is multiplicatively closed, with the *Jacobson product*

$$(a \oplus \mathbf{1})(b \oplus \mathbf{1}) = (a + b + ab) \oplus \mathbf{1}.$$

In particular left-multiplications and right-multiplications are affine maps.

Let $G = G(\mathfrak{a})$ be the set of all $a \oplus \mathbf{1}$ which have left-inverses (necessarily also in $\mathfrak{a} \oplus \{\mathbf{1}\}$). Associativity implies $a \oplus \mathbf{1}$ is left-invertible if and only if it is right-invertible as well. Evidently G is an open subset of $\mathfrak{a} \oplus \{\mathbf{1}\}$ and forms a group. Furthermore, associativity property implies both the actions of G by left- and right- multiplication, respectively, on \mathfrak{A} are affine. obtaining a bi-invariant affine structure on G .

The proof concludes with the following exercise. □

EXERCISE 10.3.5. *Show that the corresponding left-symmetric algebra on the Lie algebra of G is \mathfrak{a} .*

When G is commutative, left-invariance and right-invariance coincide. Thus every left-invariant affine structure is bi-invariant. It follows that every commutative left-symmetric algebra is associative. This purely algebraic fact has a purely algebraic proof, using the following relationship between commutators and associators.

EXERCISE 10.3.6. *Suppose that \mathfrak{a} is an \mathbb{R} -algebra. If $X, Y, Z \in \mathfrak{a}$, show that*

$$[X, Y, Z] - [X, Z, Y] + [Z, X, Y] = [XY, Z] + X[Z, Y] + [Z, X]Y$$

However, even in dimension two, commutativity alone does not imply associativity (or, equivalently, left-symmetry):

EXERCISE 10.3.7. *Show that the following table describes a commutative algebra which is not associative:*

	X	Y
X	Y	0
X	0	Y

TABLE 3. A commutative non-associative 2-dimensional algebra

The literature on left-symmetric algebras (under various names) is vast. We recommend Burde’s survey article [50], as well as works by Vinberg [278], Helmstetter [147] and Vey [275], Medina [219], Segal [247], Kim [170], and the references cited therein for more information.

10.3.2. Completeness and right-nilpotence. One can translate geometric properties of a left-invariant affine structure on a Lie group G into algebraic properties of the corresponding left-symmetric algebra \mathfrak{a} . For example, the following theorem of Helmstetter [147] and Segal [247]. indicates a kind of infinitesimal version of Markus's conjecture relating geodesic completeness to parallel volume. (See also Goldman-Hirsch [133].)

THEOREM 10.3.8. *Let G be a simply connected Lie group with left-invariant affine structure. Let $G \xrightarrow{\alpha} \text{Aff}(A)$ be the corresponding locally simply transitive affine action and \mathfrak{a} the corresponding left-symmetric algebra. The following conditions are equivalent:*

- G is a complete affine manifold;
- α is simply transitive;
- A volume form on G is parallel if and only if it is right-invariant;
- For each $g \in G$,

$$\det L(\alpha(g)) = \det \text{Ad}(g)^{-1},$$

that is, the distortion of parallel volume by α equals the modular function of G ;

- \mathfrak{a} is right-nilpotent: The subalgebra of $\text{End}(\mathfrak{a})$ generated by right-multiplications $\mathfrak{R}_a : x \mapsto xa$ is nilpotent.

The original equivalence of completeness and right-nilpotence is due to Helmstetter [147] and was refined by Segal [247]. Goldman-Hirsch [133] explain the characterization of completeness by right-invariant parallel volume as an “infinitesimal version” of the Markus conjecture (using the radiance obstruction). However, left-nilpotence of a left-symmetric algebra is a much more restrictive condition; indeed it implies right-nilpotence:

THEOREM 10.3.9 (Kim [170]). *The following conditions are equivalent:*

- The left-multiplications $x \xrightarrow{\mathfrak{L}_a} xa$ generate a nilpotent subalgebra of $\text{End}(\mathfrak{a})$;
- G is nilpotent and the affine structure is complete;
- \mathfrak{g} is a nilpotent Lie algebra and \mathfrak{a} is right-nilpotent.

10.3.3. Radiant vector fields. In a different direction, we may say that a left-invariant affine structure is *radiant* if and only if the affine representation corresponding to left-multiplication has a fixed point, that is, is conjugate to a representation $G \rightarrow \text{GL}(V)$. Equivalently, $\alpha(G)$ preserves a radiant vector field R on A . A left-invariant

affine structure on G is radiant if and only if the corresponding left-symmetric algebra has a right-identity, that is, an element $e \in \mathfrak{a}$ satisfying $ae = a$ for all $a \in \mathfrak{a}$.

10.3.4. Volume forms and the characteristic polynomial.

Let X_1, \dots, X_n be a basis for the right-invariant vector fields; it follows that the exterior product

$$\alpha(X_1) \wedge \cdots \wedge \alpha(X_n) = f(x) dx^1 \wedge \cdots \wedge dx^n$$

for a polynomial $f \in \mathbb{R}[x^1, \dots, x^n]$, called the *characteristic polynomial* of the left-invariant affine structure. In terms of the algebra \mathfrak{a} , we have

$$f(X) = \det(\mathfrak{R}_{X \oplus 1})$$

where $\mathfrak{R}_{X \oplus 1}$ denotes right-multiplication by $X \oplus 1$. By Helmsstetter [147] and Goldman-Hirsch [133], the developing map is a covering space, mapping G onto a connected the component of complement of $f^{-1}(0)$. In particular the nonvanishing of f is equivalent to completeness of the affine structure.

The following is due to Goldman-Hirsch [133]:

EXERCISE 10.3.10 (Infinitesimal Markus conjecture). *A left-invariant affine structure on a Lie group is complete if and only if right-invariant volumes forms are parallel. (Hint: first reduce to working over \mathbb{C} . Then the characteristic function f is a nonzero polynomial $\mathbb{C}^n \rightarrow \mathbb{C}$. Unless it is nonconstant, it vanishes somewhere, and the étale representation is not transitive.)*

10.4. Two-dimensional noncommutative associative algebras

Two-dimensional Lie algebras \mathfrak{g} fall into two isomorphism types:

- $\mathfrak{g} \cong \mathbb{R}^2$ (abelian);
- $\mathfrak{g} \cong \mathfrak{aff}(1, \mathbb{R})$ the Lie algebra of affine vector fields on the affine line \mathbb{A}^1 .

As we have just treated the abelian case, we turn now to the non-abelian case. The classification of affine structures on $\mathfrak{aff}(1, \mathbb{R})$ is due to Burde [49], Proposition 4.1.

If \mathfrak{g} is nonabelian, the corresponding 1-connected Lie group is the group $G^0 = \mathfrak{Aff}_+(1, \mathbb{R})$ of affine transformations of the line \mathbb{A}^1 with positive linear part. Thus G^0 is the open subset of \mathbb{A}^2 with coordinates (x, y) where $x \in \mathbb{R}$ is the translational part and $y \in \mathbb{R}^+$ is the linear part. Under this identification $G^0 \leftrightarrow \mathbb{R} \times \mathbb{R}^+ \subset \mathbb{A}^2$, both left- and right-multiplications extend to affine transformations of \mathbb{A}^2 , and thus define a *bi-invariant* affine structure on G^0 .

An element of $\text{Aff}_+(1, \mathbb{R})$ is the transformation

$$\begin{aligned} \mathbf{A}^1 &\xrightarrow{[y \mid x]} \mathbf{A}^1 \\ \xi &\longmapsto y\xi + x. \end{aligned}$$

which is the restriction of the usual linear representation

$$\begin{bmatrix} \xi \\ 1 \end{bmatrix} \longmapsto \begin{bmatrix} y & x \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \xi \\ 1 \end{bmatrix}$$

on \mathbb{R}^2 to the affine line $\mathbf{A}^1 \leftrightarrow \mathbb{R} \oplus \{1\} \subset \mathbb{R}^2$.

The identity element is $[1 \mid 0]$, inversion is:

$$[y \mid x] \longmapsto [-y^{-1}x \mid y^{-1}],$$

and the group operation is:

$$(46) \quad [y_1 \mid x_1] [y_2 \mid x_2] = [y_1 y_2 \mid x_1 + y_1 x_2].$$

In particular left-multiplication by $[\eta \mid \xi]$ extends from G^0 to the affine transformation

$$\begin{aligned} \mathbf{A}^2 &\longrightarrow \mathbf{A}^2 \\ \begin{bmatrix} x \\ y \end{bmatrix} &\longmapsto \begin{bmatrix} \eta x + \xi \\ \eta y \end{bmatrix} = \begin{bmatrix} \eta & 0 & \xi \\ 0 & \eta & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \end{aligned}$$

(taking $y_1 = \eta$, $x_1 = \xi$, $y_2 = y$, $x_2 = x$ in (46))

We will describe other left-invariant affine structures on G^0 in terms of this one, using an étale representation corresponding to left-multiplication. Left-multiplication by one-parameter subgroups define flows whose infinitesimal generators are *right-invariant* vector fields. By describing the developing maps in terms of one-parameter subgroups, we find the left-invariant vector fields and compute the left-symmetric algebra.

Here is the procedure applied to this first example.

One-parameter groups of positive homotheties $[e^t \mid 0]$ and translations $[1 \mid s]$ generate G^0 :

$$[e^t \mid s] = [1 \mid s] [e^t \mid 0]$$

and we use $s, t \in \mathbb{R}^2$ as coordinates on the group. Left multiplication by $[e^t \mid s]$ then corresponds to the affine representation

$$\begin{bmatrix} e^t & 0 & s \\ 0 & e^t & 0 \end{bmatrix}.$$

The developing map requires a choice of basepoint which lies in an open orbit of this affine representation of G^0 on \mathbf{A}^2 , so choose the basepoint

to be

$$(47) \quad p_0 := \begin{bmatrix} 0 \\ 1 \end{bmatrix} \in \mathbf{A}^2.$$

Now the developing map is given by the orbit map

$$G^0 \xrightarrow{\text{dev}} \mathbf{A}^2$$

$$[e^t \mid s] \mapsto \begin{bmatrix} e^t & 0 & \mid & s \\ 0 & e^t & \mid & 0 \end{bmatrix} p_0 = \begin{bmatrix} s \\ e^t \end{bmatrix}$$

and gives the *affine coordinates* on G^0 , which we relate to the *group coordinates* defined above. Writing

$$p = \begin{bmatrix} x \\ y \end{bmatrix}$$

we solve $[e^t \mid s] p_0 = p$ to express $(s, t) \in \mathbb{R}$ in terms of $(x, y) \in \mathbb{R} \times \mathbb{R}^+$:

$$s = x$$

$$e^t = y$$

Thus the left-multiplication which maps p_0 to p is the affine transformation

$$\begin{bmatrix} e^t & 0 & \mid & s \\ 0 & e^t & \mid & e^t \end{bmatrix} = \begin{bmatrix} y & 0 & \mid & x \\ 0 & y & \mid & y \end{bmatrix}$$

The developing map takes the identity element $e \in G^0$ to the basepoint p_0 . Moreover dev maps G^0 diffeomorphically onto the halfplane $\mathbb{R} \times \mathbb{R}^+ \subset \mathbf{A}^2$. A left-invariant vector field on G^0 is determined by its value on any point, for example e . Let $g \in G^0$. For any tangent vector $\mathbf{v} \in T_e(G^0)$, the value at g of the left-invariant vector field on G^0 extending \mathbf{v} equals the image $(D\mathfrak{L}_g)_e(\mathbf{v})$ of \mathbf{v} under the differential of left-multiplication

$$G^0 \xrightarrow{\mathfrak{L}_g} G^0$$

$$e \mapsto g$$

Since the differential of an affine transformation (in affine coordinates) is its linear part, the columns of the linear part form a basis for left-invariant vector fields. In this example, the first column and second column, respectively determine left-invariant vector fields which we denote X, Y :

$$(48) \quad \begin{bmatrix} y \\ 0 \end{bmatrix} \leftrightarrow y\partial_x =: X, \quad \begin{bmatrix} 0 \\ y \end{bmatrix} \leftrightarrow y\partial_y =: Y$$

(That these left-invariant vector fields are affine indicates that this affine structure is bi-invariant.) For future reference, the corresponding flows (right-multiplications by one-parameter subgroups) are:

$$\begin{aligned}\exp(sX) &= \begin{bmatrix} 1 & s \\ 0 & 1 \end{bmatrix} : \begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x + sy \\ y \end{bmatrix} \\ \exp(tY) &= \begin{bmatrix} 1 & 0 \\ 0 & e^t \end{bmatrix} : \begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ e^t y \end{bmatrix}\end{aligned}$$

and

$$(49) \quad \exp(tY)\exp(sX) = \begin{bmatrix} 1 & 0 \\ 0 & e^t \end{bmatrix} \begin{bmatrix} 1 & s \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & s \\ 0 & e^t \end{bmatrix}$$

We return to the algebra of left-invariant vector fields defined by covariant differentiation. Computing covariant derivatives yields the multiplication table for the corresponding left-symmetric structure, which is evidently associative. In particular $[Y, X] = X$ which defines the Lie algebra \mathfrak{g} of G^0 up to isomorphism, and we shall describe the left-symmetric structures on \mathfrak{g} in terms of this basis and the defining relation $[Y, X] = X$ for the Lie algebra.

	X	Y
X	0	0
Y	X	Y

TABLE 4. Left-invariant vector fields on $\text{Aff}_+(1, \mathbb{R})$ form an associative algebra $\mathfrak{a}_{\mathfrak{L}}$

EXERCISE 10.4.1. *The rows of the matrix correspond to the dual basis of left-invariant 1-forms:*

$$y^{-1}dx, \quad y^{-1}dy.$$

The sum of their squares is the left-invariant Poincaré metric on G^0 , regarded as the upper half-plane:

$$y^{-2}(dx^2 + dy^2)$$

In terms of the framing (X, Y) by left-invariant vector fields, the Levi-Civita connection is given by the multiplication table, which does not define a left-symmetric algebra:

	X	Y
X	Y	-X
Y	0	0

TABLE 5. Covariant derivatives of left-invariant vector fields with respect to the Levi-Civita connection of the Poincaré metric.

10.4.0.1. *Haar measures and the characteristic polynomial.* Recall that a Lie group is *unimodular* if and only if its left and right Haar measures agree. The group $\text{Aff}_+(1, \mathbb{R})$ is not *not* unimodular, and among two-dimensional 1-connected Lie groups is characterized by this. Unimodularity is obstructed by the *modular character*, the homomorphism

$$G \xrightarrow{\Delta} \mathbb{R}^+$$

$$g \mapsto \text{Det Ad}(g)$$

which relate the left-invariant and right-invariant Haar measures:

$$\mu_{\text{Right}} = \Delta \cdot \mu_{\text{Left}},$$

that is, if $g \in G$, then

$$(g_* \mu_{\text{Right}}) : S \mapsto \mu_{\text{Right}}(g^{-1}S) = \Delta(g) \cdot \mu_{\text{Right}}(S)$$

For the structure $\mathfrak{a}_{\mathcal{L}}$ above,

$$\begin{aligned} \mu_{\text{Right}} &= y \, |\partial_x \wedge \partial_y| \\ \mu_{\text{Left}} &= y^2 \, |\partial_x \wedge \partial_y| \\ \Delta([x \mid y]) &= y^{-1} \end{aligned}$$

The *characteristic polynomial* (defined in §10.3.4) is thus y .

10.4.1. The opposite structure. Since this affine structure is also invariant under right-multiplications, the left-invariant vector fields generate flows of right-multiplication by one-parameter subgroups, as in (48).

Following (49), the corresponding étale affine representation is:

$$(50) \quad \begin{bmatrix} 1 & s \\ 0 & e^t \end{bmatrix}$$

Taking the basepoint p_0 as in (47), the developing map and étale representation are:

$$(51) \quad \left[\begin{array}{cc|c} 1 & s & s \\ 0 & e^t & e^t \end{array} \right] = \left[\begin{array}{cc|c} 1 & x & x \\ 0 & y & y \end{array} \right]$$

	X'	Y'
X'	0	$-X'$
Y'	0	$-Y'$

TABLE 6. Right-invariant vector fields on $\text{Aff}_+(1, \mathbb{R})$ form an associative algebra $\mathfrak{a}_{\mathfrak{R}}$ opposite to $\mathfrak{a}_{\mathfrak{L}}$

The vector fields (where R denotes the radiant vector field)

$$(52) \quad X' := \partial_x, \quad Y' = -R = -x\partial_x - y\partial_y$$

form a basis of right-invariant vector fields satisfying the commutation relation $[Y', X'] = X'$, with covariant derivatives tabulated in Table 6. (To preserve this commutation relation, we chose $Y' = -R$ rather than $Y' = R$ — which is actually the second column of the linear part above.) We denote this algebra by $\mathfrak{a}_{\mathfrak{R}}$.

Here is the multiplication table:

EXERCISE 10.4.2. Let \mathfrak{a} be an \mathbb{R} -algebra.

- Its opposite is the algebra with multiplication defined by

$$\begin{aligned} \mathfrak{a} \times \mathfrak{a} &\longrightarrow \mathfrak{a} \\ (A, B) &\longmapsto BA \end{aligned}$$

Show that the opposite of an associative algebra is an associative algebra.

- Show $\mathfrak{a}_{\mathfrak{L}}$ and $\mathfrak{a}_{\mathfrak{R}}$ are opposite algebras but are not isomorphic to each other.
- Show that an associative algebra whose underlying Lie algebra equals $\mathfrak{g} = \text{aff}(1, \mathbb{R})$ is isomorphic to either $\mathfrak{a}_{\mathfrak{L}}$ or $\mathfrak{a}_{\mathfrak{R}}$.

10.4.2. A complete structure. The affine representation

$$[e^t \mid s] \longmapsto \left[\begin{array}{cc|c} e^t & 0 & s \\ 0 & 1 & t \end{array} \right]$$

defines a simply transitive affine of $\text{Aff}_+(1, \mathbb{R})$ on A^2 , and hence a left-invariant *complete* affine structure. The right-invariant vector fields are:

$$S := \partial_x, \quad T := x\partial_x + \partial_y$$

and the left-invariant vector fields are:

$$X := e^y\partial_x, \quad Y = \partial_y$$

with multiplication table:

	X	Y
X	0	0
Y	X	0

TABLE 7. Left-invariant vector fields for the complete left-invariant structures on $\text{Aff}_+(1, \mathbb{R})$

The left-invariant area form is $e^{-y}dx \wedge dy$ and a right-invariant area form is the parallel form $dx \wedge dy$. Note that in the basis X, Y , right-multiplications are:

$$\mathfrak{R}_X \leftrightarrow \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \mathfrak{R}_Y \leftrightarrow \mathbf{0}$$

and generate an nilpotent algebra. However, left-multiplication

$$\mathfrak{L}_Y \leftrightarrow \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

is *not* nilpotent.

10.4.3. The first halfspace family. We embed the algebra $\mathfrak{a}_{\mathbb{C}}$ in a family $\mathfrak{a}_{\mathbb{C}}^{\beta}$ as follows, where $\beta \in \mathbb{R}$ is a real parameter. The developing images of the corresponding affine structures are again halfplanes when $\beta \neq 0$, but for $\beta = 0$ the affine structure is complete.

To begin, assume that $\beta \neq 0$. The affine vector fields

$$\partial_x, \quad x\partial_x + \beta y\partial_y$$

generate an affine action of G^0 which agrees with the first action when $\beta = 1$. The corresponding one-parameter subgroups are:

$$\begin{bmatrix} 1 & 0 & | & s \\ 0 & 1 & | & 0 \end{bmatrix} \begin{bmatrix} e^t & 0 & | & 0 \\ 0 & e^{\beta t} & | & 0 \end{bmatrix} = \begin{bmatrix} e^t & 0 & | & s \\ 0 & e^{\beta t} & | & 0 \end{bmatrix}$$

which map p_0 to $\begin{bmatrix} s \\ e^{\beta t} \end{bmatrix}$, where p_0 is the basepoint defined in (47). Writing

$$p = \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} s \\ e^{\beta t} \end{bmatrix},$$

we see that the action is:

$$\begin{bmatrix} y^{1/\beta} & 0 & | & x \\ 0 & y & | & y \end{bmatrix}$$

and the columns of the linear part give left-invariant vector fields

$$y^{1/\beta}\partial_x, \quad y\partial_y.$$

Although the second vector field is affine, the first vector field is affine if and only if $\beta = 1$, that is, for the bi-invariant structure.

Their covariant derivatives form the multiplication table:

	$y^{1/\beta} \partial_x$	$y \partial_y$
$y^{1/\beta} \partial_x$	0	0
$y \partial_y$	$1/\beta y^{1/\beta} \partial_x$	$y \partial_y$

TABLE 8. Left-invariant vector fields of deformation of bi-invariant structure

When $\beta = 0$, the original affine representation has no open orbits. However, a simple modification extends this structure to $\beta = 0$, by including a *complete affine structure*.

To this end, replace the second one-parameter subgroup by:

$$\left[\begin{array}{cc|c} e^t & 0 & 0 \\ 0 & e^{\beta t} & f_\beta(t) \end{array} \right] = \exp \left[\begin{array}{cc|c} t & 0 & 0 \\ 0 & \beta t & t \end{array} \right]$$

where f_β denotes the continuous function $\mathbb{R} \rightarrow \mathbb{R}$ defined by:

$$f_\beta(t) := \begin{cases} \frac{e^{\beta t} - 1}{\beta} & \text{if } \beta \neq 0 \\ t & \text{if } \beta = 0 \end{cases}$$

(Note that when $\beta \neq 0$, this new action is conjugate to the original action by the translation $\text{Trans}_{(0, -1/\beta)}$.) The affine action of G^0 is given by

$$\left[\begin{array}{cc|c} 1 & 0 & s \\ 0 & 1 & 0 \end{array} \right] \left[\begin{array}{cc|c} e^t & 0 & 0 \\ 0 & e^{\beta t} & f_\beta(t) \end{array} \right] = \left[\begin{array}{cc|c} e^t & 0 & s \\ 0 & e^{\beta t} & f_\beta(t) \end{array} \right]$$

The orbit of p_0 is the halfplane defined by $y > -1/\beta$ if $\beta \neq 0$ and all of \mathbb{A}^2 if $\beta = 0$. Evaluating at p_0 yields the affine representation (taking p_0 to p)

$$\left[\begin{array}{cc|c} (1 + \beta y)^{1/\beta} & 0 & x \\ 0 & (1 + \beta y) & y \end{array} \right]$$

and taking the columns of the linear part yields a basis of left-invariant vector fields:

$$X = \begin{cases} (1 + \beta y)^{1/\beta} \partial_x & \text{if } \beta \neq 0 \\ e^y \partial_x & \text{if } \beta = 0 \end{cases}$$

$$Y = (1 + \beta y) \partial_y$$

whose covariant derivatives are recorded in Table 9. Note that $\beta = 1$ corresponds to the bi-invariant (associative) structure and $\beta = 0$ corresponds to a complete structure discussed in §10.4.2:

	X	Y
X	0	0
Y	X	βY

TABLE 9. Left-invariant vector fields on the halfspace family structures define an algebra $\mathfrak{a}_{\mathbb{C}}^{\beta}$ depending on a parameter β

10.4.3.1. *Lorentzian structure.* The case $\beta = -1$ is also interesting. Then the affine structure arises from an invariant flat Lorentzian structure. Explicitly, if X, Y base the left-invariant vector fields:

$$X = y^{-1} \partial_x, \quad Y = y \partial_y$$

and X^*, Y^* is the dual basis of left-invariant 1-forms:

$$X^* = y dx, \quad Y^* = y^{-1} dy,$$

then the symmetric product $X^* \odot Y^* = dx \odot dy$ is a parallel Lorentzian structure, defining a flat Lorentzian structure on G invariant under left-multiplications.

Again the developing image of the corresponding flat structure is a halfplane so this structure is an *incomplete flat Lorentzian structure*, which is *homogeneous*. This contrasts the theorem of Marsden [213] that a *compact* homogeneous pseudo-Riemannian manifold is geodesically complete, indicating that the compactness hypothesis is necessary.

CONJECTURE 10.4.3. *A compact locally homogeneous pseudo-Riemannian manifold is geodesically complete.*

This has been proved by Klingler [173] for *constant curvature* Lorentzian manifolds.

10.4.4. Parabolic deformations. The case where $\beta = 2$ is also interesting, for several reasons. Here there exist nonradiant deformations whose developing images are *parabolic subdomains* of \mathbb{A}^2 , the components of the complements of a parabola in \mathbb{A}^2 . (Recall that the complement of a parabola has *two* connected components, one which is convex and the other concave.) To this end, consider a parameter $\alpha \in \mathbb{R}$; when $\alpha = 0$, these examples are just the $\mathfrak{a}_{\mathbb{C}}^2$ as before, but when $\alpha \neq 0$, these examples are all affinely conjugate, but the halfplane deforms to a parabolic subdoman.

After describing this structure as a deformation, we mention its surprising role as the first *simple* (in the sense of Burde [49]) left-symmetric algebra whose underlying Lie algebra is solvable. Then we

describe it as a *clan* in the sense of Vinberg [278], and briefly describe Vinberg's theory of convex homogeneous domains, which was one of the historical origins of the theory of left-symmetric algebras.

10.4.4.1. *Parabolic domains.* Let

$$(53) \quad f_\alpha(x, y) := y - \frac{\alpha}{2}x^2;$$

the parabolic subdomain

$$\Omega_\alpha := \{(x, y) \in \mathbb{A}^2 \mid f_\alpha(x, y) > 0\}$$

is a halfplane if $\alpha = 0$, convex if $\alpha \geq 0$, and concave if $\alpha \leq 0$. The one-parameter subgroups

$$\begin{aligned} \exp \left[\begin{array}{cc|c} 0 & 0 & s \\ \alpha s & 0 & 0 \end{array} \right] \exp \left[\begin{array}{cc} t & 0 \\ 0 & 2t \end{array} \right] &= \left[\begin{array}{cc|c} 1 & 0 & s \\ \alpha s & 1 & \frac{\alpha}{2} s^2 \end{array} \right] \left[\begin{array}{cc} e^t & 0 \\ 0 & e^{2t} \end{array} \right] \\ &= \left[\begin{array}{cc|c} e^t & 0 & s \\ \alpha e^t s & e^{2t} & \frac{\alpha}{2} s^2 \end{array} \right] \end{aligned}$$

generate its affine automorphism group. The image of the the basepoint

$$p_0 := \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

is

$$p := \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} s \\ e^{2t} + \frac{\alpha}{2} s^2 \end{bmatrix}$$

so the group coordinates (s, t) relate to the affine coordinates (x, y) by

$$\begin{aligned} s &= x \\ t &= \frac{1}{2} \log \left(y - \frac{\alpha}{2} x^2 \right) = \frac{1}{2} \log f_\alpha(x, y) \end{aligned}$$

where f_α is defined in (53). The left-invariant vector fields are based by the first two columns of the matrix

$$\left[\begin{array}{cc|c} (y - \frac{\alpha}{2}x^2)^{1/2} & 0 & x \\ \alpha(y - \frac{\alpha}{2}x^2)^{1/2}x & (y - \frac{\alpha}{2}x^2) & y \end{array} \right] = \left[\begin{array}{cc|c} f_\alpha(x, y)^{1/2} & 0 & x \\ \alpha f_\alpha(x, y)^{1/2}x & f_\alpha(x, y) & y \end{array} \right]$$

which are:

$$X := \sqrt{f_\alpha(x, y)} (\partial_x + \alpha x \partial_y), \quad Y := f_\alpha(x, y) \partial_y$$

Table 10 describes the multiplication in the corresponding left-symmetric algebra.

EXERCISE 10.4.4. *Show that, as α varies, these structures are related by the polynomial diffeomorphism:*

$$\begin{aligned} \mathbb{A}^2 &\longrightarrow \mathbb{A}^2 \\ (x, y) &\longmapsto (x, y - \alpha x^2/2) = (x, f_\alpha(x, y)) \end{aligned}$$

10.4.4.2. *Simplicity.* The algebra $\mathfrak{a}_\mathbb{C}^2(\alpha)$, where $\alpha \neq 0$, also has special *algebraic* significance. Define a left-symmetric algebra to be *simple* if and only if it contains no nonzero proper two-sided ideals. Somewhat surprisingly, the Lie algebra underlying a simple left-symmetric algebra can even be solvable. Indeed, Burde [49] proved that, over \mathbb{C} , the complexification of the above example is the only *simple* left-symmetric algebra of dimension two. Over \mathbb{R} , this example, and the field \mathbb{C} itself (regarded as an \mathbb{R} -algebra), are the only simple 2-dimensional left-symmetric algebras. The classification of simple left-symmetric algebras in general is a difficult unsolved algebraic problem; see [49, 50] for more details.

10.4.4.3. *Clans and homogeneous cones.* Vinberg [278] classifies convex homogeneous domains in terms of special left-symmetric algebras, which he calls *clans*. The example above is the first nontrivial example of such a clan, and can be approached in several different ways.

Let $\Omega \subset \text{Mat}_2(\mathbb{R})$ denote the convex cone comprising *positive definite symmetric* 2×2 real matrices. It is an open subset of the 3-dimensional linear subspace of $\text{Mat}_2(\mathbb{R})$ consisting of *symmetric* matrices. Thus $\Omega \subset \mathbb{W}$ is an open convex cone.

$\Omega \subset \mathbb{W}$ is *homogeneous*: Namely $\text{Aut}(\mathbb{R}^2) = \text{GL}(2, \mathbb{R})$ acts on \mathbb{W} by the induced action on symmetric bilinear forms:

$$\begin{aligned} \text{GL}(2, \mathbb{R}) \times \mathbb{W} &\longrightarrow \mathbb{W}. \\ (A, w) &\longmapsto A^\dagger w A \end{aligned}$$

	X	Y
X	$\alpha/2 Y$	0
Y	X	$2Y$

TABLE 10. Left-invariant vector fields on the parabolic deformation $\mathfrak{a}_\mathbb{C}^2(\alpha)$

where A^\dagger denotes the transpose of A . Under the linear correspondence

$$\mathbb{W} \longrightarrow \mathbb{R}^3$$

$$\begin{bmatrix} x & y \\ y & z \end{bmatrix} \longmapsto \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

defines an action of $\mathrm{GL}(2, \mathbb{R})$ on \mathbb{R}^3 preserving the open subset $\Omega \subset \mathbb{W}$ defined by the positivity of determinant of the matrix

$$\Delta(x, y, z) = xz - y^2,$$

which appears as the *characteristic polynomial* of the corresponding left-invariant structure (or left-symmetric algebra). Furthermore, by the Gram-Schmidt orthonormalization process, $\mathrm{GL}(2, \mathbb{R})$ acts transitively on Ω .

10.4.5. Deformations of the opposite structure. The other associative structure $\mathfrak{a}_{\mathfrak{g}}$ has two kinds of deformations. The first one we describe uses the left-invariant parallel vector field on $\mathfrak{a}_{\mathfrak{g}}$ to construct a deformation, depending a real parameter $\eta \in \mathbb{R}$ as follows. It is nonradiant. The second one is radiant, and replaces the diagonal one-parameter subgroup with one eigenvalue 1 by one with two eigenvalues of varying strength.

10.4.5.1. *Nonradiant deformation.* Consider the étale representation

$$\left[\begin{array}{cc|c} 1 & s & \eta t \\ 0 & e^t & 0 \end{array} \right]$$

which is a deformation of the representation (50). The action is:

$$p_0 := \begin{bmatrix} 0 \\ 1 \end{bmatrix} \longmapsto \begin{bmatrix} s + \eta t \\ e^t \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$$

so the group coordinates relate to the affine coordinates by:

$$s := x - \eta \log(y)$$

$$t := \log(y)$$

and the étale affine representation is:

$$\left[\begin{array}{cc|c} 1 & x - \eta \log(y) & x \\ 0 & y & y \end{array} \right]$$

The columns of the linear part (adjusted) determine left-invariant vector fields:

$$X' := \partial_x, \quad Y'_\eta = (\eta \log(y) - x) \partial_x - y \partial_y$$

which specialize to the left-invariant vector fields on $\mathfrak{a}_{\mathfrak{g}}$ as in (52) with multiplication table:

	X'	Y'_η
X'	0	$-X'$
Y'_η	0	$-\eta X' - Y'_\eta$

TABLE 11. Nonradiant deformation of $\mathfrak{a}_{\mathfrak{R}}$ depending on a parameter $\eta \in \mathbb{R}$

10.4.5.2. *Radiant deformation.* Consider the étale affine representation

$$\left[\begin{array}{cc|c} e^{\mu t} & e^{\mu t} s & 0 \\ 0 & e^t & 0 \end{array} \right]$$

which is a deformation of the representation (50) when $\mu \neq 0$. (When $\mu = 0$, the resulting group is abelian.) The developing map is:

$$p_0 := \begin{bmatrix} 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} e^{\mu t} s \\ e^t \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$$

so the group coordinates relate to the affine coordinates by:

$$\begin{aligned} s &:= y^{-1/\mu} x \\ t &:= \log(y) \end{aligned}$$

and the étale affine representation is:

$$\left[\begin{array}{cc|c} y^\mu & x & x \\ 0 & y & y \end{array} \right]$$

The columns of the linear part (adjusted) determine left-invariant vector fields:

$$X'_\mu := y^\mu \partial_x, \quad Y' = -x \partial_x - y \partial_y$$

which specialize to the left-invariant vector fields on $\mathfrak{a}_{\mathfrak{R}}$ as in (52) with multiplication table:

	X'_μ	Y'
X'_μ	0	$-X'_\mu$
Y'	$\mu X'_\mu$	$-Y'$

TABLE 12. Radiant deformation of $\mathfrak{a}_{\mathfrak{R}}$ depending on an eigenvalue parameter $\mu \in \mathbb{R}$

10.4.5.3. *Radiancy and parallel volume.* When $\mu = -1$, the structure has parallel volume — that is, area forms are parallel if and only if they are left-invariant. In that case the vector fields $x\partial_x + y\partial_y$, $y^{-1}\partial_x$ base the space of left-invariant vector fields and the 1-forms $y^{-1}dy$, $ydx - xdy$ base the space of left-invariant 1-forms. In contrast to affine structures on closed manifolds, this left-invariant affine structure is both radiant and has parallel volume.

10.4.5.4. *Deforming to the complete structure.* As above, we can reparametrize this family to include the complete structure when $\mu = 0$. Namely, choose a new real parameter δ (which will be $1/(\mu - 1)$).

For each $\delta \in \mathbb{R}$, the vector fields

$$(1 + \delta y)\partial_x, \quad (1 + \delta)x\partial_x + (1 + \delta y)\partial_y$$

generate étale affine representations (and thus left-invariant affine structures) on $\text{Aff}_+(1, \mathbb{R})$ such that the vector fields

$$\begin{aligned} X &\longleftrightarrow X_R^\delta := (1 + \delta y)^{\frac{1}{\delta}+1} \partial_x \\ Y &\longleftrightarrow Y_R^\delta := \delta x \partial_x + (1 + \delta y) \partial_y \end{aligned}$$

base the left-invariant vector fields, and $[Y, X] = X$. The multiplication table is:

	X	Y
X	0	δX
Y	$(\delta + 1)X$	δY

TABLE 13. Deformations of $\mathfrak{a}_{\mathfrak{R}}$ depending on an eigenvalue parameter $\delta \in \mathbb{R}$ containing the complete structure at $\delta = 0$

10.5. Complete affine structures on closed 3-manifolds

The closed complete affine 3-manifolds were classified in Fried-Goldman [110], based on the resolution of Auslander’s question in dimension 3. There are two isomorphism types of nilpotent 3-dimensional Lie groups, namely \mathbb{R}^3 and the 3-dimensional Heisenberg group. In general one must consider semidirect products of \mathbb{R}^2 by a one-parameter group, but for the purpose of structures on closed 3-manifolds, only one new isomorphism type is needed, that is, when the action of \mathbb{R} on \mathbb{R}^2 is:

$$(x, y) \mapsto (e^t x, e^{-t} y)$$

In that case the group is realized as $\text{Isom}^0(\mathbb{E}^{1,1})$.

The classification in the nilpotent case involves computing the subgroup of translations in the center. The solvable case reduces to the nilpotent case by the useful fact that the unipotent radical of the Zariski closure of a simply transitive subgroup itself acts simply transitively. Thus, underlying every simply transitive affine action is a *unipotent* simply transitive affine action. Every solvable group with complete left-invariant affine structure has an underlying structure as a *nilpotent* Lie group with complete left-invariant affine structure.

In the nilpotent case, simple transitivity is equivalent to unipotence of the linear part:

PROPOSITION 10.5.1 (Scheuneman [244]). *Let G be a nilpotent group with left-invariant affine structure. Let $G \xrightarrow{\rho} \text{Aff}(\mathbf{A})$ be the étale representation corresponding to left-multiplication. Then the affine structure is complete (that is, ρ is simply transitive) if and only if $L \circ \rho$ is unipotent.*

PROOF. Suppose first that $L \circ h$ is unipotent. Let O be an open orbit corresponding to the developing image. By Rosenlicht [240] (and independently, Kostant (unpublished)), every orbit of a connected unipotent group is Zariski-closed. Thus O is both open and closed (in the classical topology). Since \mathbf{A} is connected, $O = \mathbf{A}$, that is, G acts transitively. Since $\dim(\mathbf{A}) = \dim(G)$, every isotropy group is discrete. Since G is unipotent, every isotropy group is torsionfree and Zariski closed, which implies that G acts freely. Thus G acts simply transitively as desired.

Conversely, suppose that G acts simply transitively. By the structure theorem for representations of nilpotent groups, there exists a maximal invariant affine subspace \mathbf{A}_0 (the *Fitting subspace*) upon which ρ acts unipotently. Since G acts transitively, $\mathbf{A}_0 = \mathbf{A}$ and thus ρ is unipotent. \square

This includes a sort of an infinitesimal converse to Theorem 8.4.1, that a closed affine manifold with unipotent holonomy is geodesically complete.

10.5.1. Central translations. The key step of the classification in dimension 3 is the existence of *central translations* in a simply transitive group of unipotent affine transformations. This was conjectured by Auslander [11] and erroneously claimed by Scheuneman [244]. Fried [108] produced a 4-dimensional counterexample; see Kim [170] and Dekimpe [] for further developments in this basic question.

In terms of left-symmetric algebras, translations correspond to *right-invariant parallel vector fields*. A translation is central if and only

if the corresponding parallel vector field is also also *left-invariant*. Left-invariant parallel vector fields correspond to elements $P \in \mathfrak{a}$ of the left-symmetric algebra \mathfrak{a} such that $\mathfrak{a} \cdot P = 0$; being central means that $[\mathfrak{a}, P] = 0$. Thus the central translations correspond to elements $P \in \mathfrak{a}$ such that $P\mathfrak{a} = \mathfrak{a}P = 0$. Such elements P clearly form a two-sided ideal $\mathfrak{Z} = \mathfrak{Z}(\mathfrak{a})$ and the quotient $\mathfrak{a}/\mathfrak{Z}$ is a left-symmetric algebra.

EXERCISE 10.5.2. *If \mathfrak{a} corresponds to a complete affine structure, then $\mathfrak{a}/\mathfrak{Z}$ corresponds to a complete affine structure.*

Now we discuss the 3-dimensional nilpotent left-symmetric algebras, or equivalently unipotent simply transitive affine actions on \mathbb{A}^3 , which are in turn or equivalently, complete left-invariant affine structures on nilpotent Lie groups. If 1-connected 3-dimensional Lie group is not abelian (in which case it's isomorphic to \mathbb{R}^3), then it is isomorphic to the *Heisenberg group*, consisting of 3×3 upper-triangular unipotent matrices:

$$\begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}, a, b, c \in \mathbb{R}$$

EXERCISE 10.5.3. *The 3-dimensional nilpotent associative algebras are: (Compare [110])*

$\dim(\mathfrak{g}) \geq 2$:

	X	Y	Z
X	0	0	0
Y	0	0	0
Z	0	0	0

	X	Y	Z
X	0	0	0
Y	0	0	0
Z	0	0	Y

TABLE 14. Multiplication tables for $\dim(\mathfrak{g}) = 3$ and $\dim(\mathfrak{g}) = 2$, respectively

$\dim(\mathfrak{g}) = 1$ and G/\mathfrak{g} Euclidean :

	X	Y	Z
X	0	0	0
Y	0	$a_{11}X$	$a_{12}X$
Z	0	$a_{21}X$	$a_{22}X$

TABLE 15. When $\dim(\mathfrak{g}) = 1$ and G/\mathfrak{g} Euclidean, the structure is defined by bilinear form $G/\mathfrak{g} \times G/\mathfrak{g} \xrightarrow{\mathfrak{a}} \mathfrak{g}$

$\dim(\mathfrak{g}) = 1$ and G/\mathfrak{g} non-Riemannian:

	X	Y	Z
X	0	0	0
Y	0	0	bX
Z	0	cX	Y

TABLE 16. When $\dim(\mathfrak{g}) = 1$ and G/\mathfrak{g} non-Riemannian, the structure is defined by a pair $(b, c) \in \mathbb{R}^2$

G is abelian when $\dim(\mathfrak{g}) \geq 2$, and in these cases G is a product of a 1-dimensional structure and a 2-dimensional structure. These cases appear as limits in the generic situation when $\dim(\mathfrak{g}) = 1$.

When $\dim(\mathfrak{g}) = 1$ and the structure on G/\mathfrak{g} is Euclidean, then the induced product

$$\begin{aligned} \mathfrak{a}/\mathfrak{g} \times \mathfrak{a}/\mathfrak{g} &\longrightarrow \mathfrak{a} \\ (A + \mathfrak{g}, B + \mathfrak{g}) &\longmapsto AB \end{aligned}$$

defines a bilinear form $\mathbb{R}^2 \times \mathbb{R}^2 \xrightarrow{a} \mathfrak{Z} \cong \mathbb{R}$. Table 15 arises by taking X to be a generator of \mathfrak{Z} . The structure is abelian if and only if \mathbf{a} is symmetric. In this case \mathfrak{a} is Euclidean if $\mathbf{a} = 0$ and a product ($\dim(\mathfrak{Z}) = 2$) when \mathbf{a} is nonzero and degenerate.

When $\dim(\mathfrak{Z}) = 1$ and the structure on G/\mathfrak{Z} is non-Riemannian, let X generate \mathfrak{Z} , and extend to a basis $\{X, Y, Z\}$ so that

$$(Z + \mathfrak{Z})^2 = Y$$

in the non-Riemannian quotient. The results are tabulated in Table 16. The structure is abelian if and only if $b = c$, and corresponds to the product structure ($\dim(\mathfrak{Z}) = 2$) if and only if $b = c = 0$.

10.6. Fried's counterexample to Auslander's conjecture

Auslander's conjecture [11] that a nilpotent simply transitive affine group contains central translations was disproved by Fried [108] by the following 4-dimensional example.

A nilpotent Lie algebra is said to be *filiform* if it is "maximally non-abelian," in the sense that its degree of nilpotence is one less than its dimension. That is, a k -step nilpotent Lie algebra is filiform if its dimension equals $k + 1$. Fried's example lives on the unique 4-dimensional filiform Lie algebra. (Curiously, Benoist's example of an 11-dimensional nilpotent Lie algebra admitting no faithful 12-dimensional linear representation — and hence no simply transitive affine actions — is a filiform Lie algebra.)

Let $t, u, v, w \in \mathbb{R}$ be real parameters. We parametrize the 4-dimensional filiform algebra as a semidirect sum

$$\mathfrak{g} := \mathbf{V}_0 \rtimes_{J_3} \mathbb{R}T,$$

where \mathbf{V}_0 is a 3-dimensional abelian ideal with coordinates u, v, w and T acts by the 3-dimensional Jordan block

$$J_3 := \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0. \end{bmatrix}$$

That is, \mathfrak{g} admits a basis U, V, W, T subject to nonzero commutation relations

$$[T, U] = V, \quad [T, V] = W$$

and all other brackets between basic elements are zero.

We start with a simply transitive affine action, where U, V, W act by translations. The Lie algebra representation is:

$$\mathcal{A}_0(t, u, v, w) := \left[\begin{array}{cccc|c} 0 & 0 & 0 & 0 & t \\ & 0 & t & 0 & u \\ & & 0 & t & v \\ & & & 0 & w \end{array} \right],$$

the group representation is:

$$A_0(t, u, v, w) := \exp \mathcal{A}_0(t, u, v, w) := \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & t \\ & 1 & t & t^2/2 & u + tv/2 + t^2w/6 \\ & & 1 & t & v + tw/2 \\ & & & 1 & w \end{array} \right]$$

and a basis of left-invariant vector fields is:

$$X_0 := \partial_x, \quad Y_0 := \partial_y, \quad Z_0 := \partial_z + x\partial_y, \quad W_0 := \partial_w + x\partial_z + \frac{x^2}{2}\partial_y$$

with multiplication table:

	X_0	Y_0	Z_0	W_0
X_0	0	0	Y_0	Z_0
Y_0	0	0	0	0
Z_0	0	0	0	0
W_0	0	0	0	0

TABLE 17. Complete affine structure on 4-dimensional filiform algebra

Then X_0, Y_0 are parallel vector fields and $\mathfrak{Z}(\mathfrak{g})$ is one-dimensional, spanned by Y_0 .

Now we deform using a parameter $\lambda \in \mathbb{R}$, in the direction of the parallel vector field $X_0 = \partial_x$. Consider the affine representation $A = A_\lambda$,

$$\mathcal{A}(t, u, v) = \mathcal{A}_\lambda := \left[\begin{array}{cccc|c} 0 & -\lambda w & \lambda v & -\lambda u & t \\ & 0 & t & 0 & u \\ & & 0 & t & v \\ & & & 0 & w \end{array} \right]$$

Since

$$a^2 := \left[\begin{array}{ccc|c} 0 & -\lambda tw & \lambda tv & \lambda(-2uw + v^2) \\ & 0 & t^2 & tv \\ & & 0 & tw \\ & & & 0 \end{array} \right],$$

$$a^3 := \left[\begin{array}{ccc|c} 0 & -\lambda t^2 w & 0 & \\ & 0 & t^2 w & \\ & & 0 & \end{array} \right], \quad A^4 := \left[\begin{array}{c|c} 0 & -\lambda t^2 w^2 \\ & 0 \end{array} \right],$$

the general group element is:

$$\exp(A_\lambda) := \left[\begin{array}{cccc|c} 1 & -\lambda w & \lambda v & \lambda(-u + tv/2 - t^2 w/6) & t + \lambda(-uw + v^2/2 - t^2 w^2/24) \\ & 1 & t & t^2/2 & u + tv/2 + t^2 w/6 \\ & & 1 & t & v + tw/2 \\ & & & 1 & w \end{array} \right].$$

The last column of this matrix is the developing map, and we can relate the group coordinates to the affine coordinates, by:

$$\begin{aligned} x &= t + \lambda(-uw + v^2/2 - t^2 w^2/24) \\ y &= u + tv/2 + t^2 w/6 \\ z &= v + tw/2 \\ w &= w \end{aligned}$$

A basis for right-invariant vector fields is:

$$T := \partial_x + z\partial_y + w\partial_z, \quad U := \partial_y - \lambda w\partial_x, \quad V := \partial_z + \lambda z\partial_x, \quad \widetilde{W} := \partial_w - \lambda y\partial_x$$

with U central.

I think the multiplication table for Fried's example is:

	X	Y	Z	W
X	0	0	Y	Z
Y	0	0	0	λX
Z	0	0	λX	0
W	0	λX	0	0

TABLE 18. Fried's counterexample to Auslander's conjecture on central translations

10.7. Solvable 3-dimensional algebras

If M^3 is a closed manifold with complete affine structure, then M is affinely isomorphic to a finite quotient of a *complete affine solv-manifold*, that is, a homogeneous space $\Gamma \backslash G$, where G is a Lie group with a complete affine structure and $\Gamma < G$ is a lattice. Equivalently, G admits a simply transitive affine action (corresponding to left-multiplication). Necessarily G is solvable and we may assume that G is simply-connected. Since G admits a lattice, it is unimodular.

We have already discussed the cases when G is nilpotent. There are two isomorphism classes of simply connected solvable unimodular non-nilpotent Lie groups:

- The universal covering $\widetilde{\text{Isom}}^0(\mathbb{E}^2)$ of orientation-preserving isometries of the group $\text{Isom}^0(\mathbb{E}^2)$ of orientation-preserving isometries of the Euclidean plane \mathbb{E}^2 ;
- The identity component of the group $\text{Isom}^0(\mathbb{E}^{1,1})$ of orientation-preserving isometries of 2-dimensional Minkowski space.

EXERCISE 10.7.1. *Prove that every lattice $\Gamma < \widetilde{\text{Isom}}^0(\mathbb{E}^2)$ contains a free abelian subgroup of finite index. In particular, Γ is a finite extension of \mathbb{Z}^3 and is a 3-dimensional Bieberbach group.*

Thus the only interesting remaining case for classifying complete affine 3-manifolds occurs for the group $\text{Isom}^0(\mathbb{E}^{1,1})$.

There are two isomorphism classes of complete left-invariant affine structures on this group. One, which probably goes back to Auslander-Markus [9] is a group of isometries of a parallel Lorentzian metric; the other is due to Auslander [11] and corresponds to the simply transitive action, where $\lambda \in \mathbb{R}$ is a parameter (compare Fried-Goldman [110], Theorem 4.1):

$$(s, t, u) \xrightarrow{\rho_\lambda} \left[\begin{array}{ccc|c} 1 & \lambda e^s u & \lambda e^{-s} t & s + \lambda t u \\ 0 & e^s & 0 & t \\ 0 & 0 & e^{-s} & u \end{array} \right]$$

When $\lambda \neq 0$, these actions are all affinely conjugate. The case $\lambda = 0$ is the original flat Lorentzian structure.

Auslander notes that when $\lambda \neq 0$, the simply transitive action contains *no* translations.

The corresponding left-invariant vector fields are:

$$\begin{aligned} X &:= \partial_x \\ Y &:= e^{x-\lambda yz} (\lambda z \partial_x + \partial_y) \\ Z &:= e^{-x+\lambda yz} (\lambda y \partial_x + \partial_z) \end{aligned}$$

with multiplication table:

	X	Y	Z
X	0	Y	$-Z$
Y	0	0	λX
Z	0	λX	0

10.8. Incomplete affine 3-manifolds

10.8.1. Parabolic cylinders. We can extend this structure to structures on a 3-dimensional solvable Lie group G , which admits compact quotients. These provide examples of compact convex incomplete affine 3-manifolds which are not *properly convex*, and nonradiant. Therefore Vey's result that compact hyperbolic affine manifolds are radiant is sharp.

Further examples from the same group action give *concave* affine structures on these same 3-manifolds.

The function:

$$\begin{aligned} \mathbb{A}^3 &\xrightarrow{f} \mathbb{R} \\ (x, y, z) &\longmapsto x - y^2/2 \end{aligned}$$

is invariant under the affine \mathbb{R}^2 -action defined by:

$$\begin{aligned} \mathbb{R}^2 &\xrightarrow{u} \text{Aff}(\mathbb{A}^3) \\ (t, u) &\longmapsto \exp \left[\begin{array}{ccc|c} 0 & t & 0 & 0 \\ 0 & 0 & 0 & t \\ 0 & 0 & 0 & u \end{array} \right] = \left[\begin{array}{ccc|c} 1 & t & 0 & t^2/2 \\ 0 & 1 & 0 & t \\ 0 & 0 & 1 & u \end{array} \right] \end{aligned}$$

Under the 1-parameter group of dilations

$$\begin{aligned} \mathbb{R} &\xrightarrow{\delta} \text{Aff}(\mathbb{A}^3) \\ s &\longmapsto \exp \left[\begin{array}{ccc} 2s & 0 & 0 \\ 0 & s & 0 \\ 0 & 0 & -s \end{array} \right] = \left[\begin{array}{ccc} e^{2s} & 0 & 0 \\ 0 & e^s & 0 \\ 0 & 0 & e^{-s} \end{array} \right] \end{aligned}$$

the function f scales as:

$$f \circ \delta(s) = e^{2s} f.$$

The group $G \subset \text{Aff}(\mathbb{A}^3)$ generated by $\mathcal{U}(t, u)\delta(s)$ (for $s, t, u \in \mathbb{R}$) acts simply transitively on the open convex parabolic cylinder defined by $f(x, y, z) > 0$ as well as on the open concave parabolic cylinder defined by $f(x, y, z) < 0$. The corresponding left-invariant affine structure on the Lie group G has a basis of left-invariant vector fields

$$\begin{aligned} X &:= f(x, y, z)\partial_x \\ Y &:= f(x, y, z)^{1/2}(y\partial_x + \partial_y) \\ Z &:= f(x, y, z)^{-1/2}\partial_z \end{aligned}$$

with multiplication recorded in Table 19. The dual basis of left-invariant 1-forms is:

$$\begin{aligned} X^* &:= f(x, y, z)^{-1}(dx - ydy) = -d \log(f) \\ Y^* &= f(x, y, z)^{-1/2}dy \\ Z^* &= f(x, y, z)^{1/2}dz \end{aligned}$$

with *bi-invariant* volume form

$$X^* \wedge Y^* \wedge Z^* = f(x, y, z)^{-1}dx \wedge dy \wedge dz.$$

This example is due to Goldman [116] providing examples of non-conical convex domains covering compact affine manifolds.

	X	Y	Z
X	X	$Y/2$	$-Z/2$
Y	0	X	0
Z	0	0	0

TABLE 19. Algebra corresponding to parabolic 3-dimensional halfspaces

10.8.2. Nonradiant deformations of radiant halfspace quotients. Another example arises from radiant suspensions. Namely,

consider the radiant affine representation

$$\begin{aligned} \mathbb{R} \times \mathbb{R}^2 &\xrightarrow{\rho_\alpha} \text{Aff}(\mathbf{A}^3) \\ (s; t, u) &\longmapsto e^{\alpha s} \exp \begin{bmatrix} s & 0 & t \\ 0 & -s & u \\ 0 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} e^{(\alpha+1)s} & 0 & e^{\alpha s} t \\ 0 & e^{(\alpha-1)s} & e^{\alpha s} u \\ 0 & 0 & e^{\alpha s} \end{bmatrix} \end{aligned}$$

depending on a parameter $\alpha \in \mathbb{R}$. When $\alpha \neq 0$, the action is locally simply transitive; the open orbits are the two halfspaces defined by $z > 0$ and $z < 0$ respectively. The vector fields

$$\begin{aligned} X &:= z^{(\alpha+1)/\alpha} \partial_x \\ Y &:= z^{(\alpha-1)/\alpha} \partial_y \\ Z &:= R = x\partial_x + y\partial_y + z\partial_z \end{aligned}$$

correspond to a basis of left-invariant vector fields, with multiplication recorded in Table 20.

	X	Y	Z
X	0	0	X
Y	0	0	Y
Z	$((\alpha+1)/\alpha)X$	$((\alpha-1)/\alpha)Y$	Z

TABLE 20. A radiant suspension

When $\alpha = \pm 1$, then this action admits *nonradiant* deformations. Namely let $\beta \in \mathbb{R}$ be another parameter, and consider the case when $\alpha = 1$. The affine representation

$$\begin{aligned} \mathbb{R} \times \mathbb{R}^2 &\xrightarrow{\rho^\beta} \text{Aff}(\mathbf{A}^3) \\ (s; t, u) &\longmapsto \exp \left[\begin{array}{ccc|c} 2s & 0 & 0 & 0 \\ 0 & 0 & 0 & \beta s \\ 0 & 0 & s & 0 \end{array} \right] \cdot \exp \begin{bmatrix} 0 & 0 & t \\ 0 & 0 & u \\ 0 & 0 & 0 \end{bmatrix} \\ &= \left[\begin{array}{ccc|c} e^{2s} & 0 & e^{2s}t & 0 \\ 0 & 1 & u & \beta s \\ 0 & 0 & e^s & 0 \end{array} \right] \end{aligned}$$

maps

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \xrightarrow{\rho^\beta} \begin{bmatrix} e^{2st} \\ u + \beta s \\ e^s \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

so the group element with coordinates (s, t, u) corresponds to the point with coordinates

$$\begin{aligned} x &= e^{2st} \\ y &= u + \beta s \\ z &= e^s \end{aligned}$$

with inverse transformation:

$$\begin{aligned} s &= \log(z) \\ t &= z^{-2}x \\ u &= y - \beta \log(z) \end{aligned}$$

Then the linear part $L\rho^\beta(s, t, u)$ corresponds to the matrix

$$\begin{bmatrix} z^2 & 0 & x \\ 0 & 1 & y - \beta \log(z) \\ 0 & 0 & z \end{bmatrix}$$

whose columns base the left-invariant vector fields:

$$\begin{aligned} X &:= z^2 \partial_x \\ Y &:= \partial_y \\ Z &:= x \partial_x + (y - \beta \log(z)) \partial_y + z \partial_z \end{aligned}$$

Table 21 records their covariant derivatives.

	X	Y	Z
X	0	0	X
Y	0	0	Y
Z	$2X$	0	$Z - \beta Y$

TABLE 21. Nonradiant Deformation

CHAPTER 11

Parallel volume and completeness

A particularly tantalizing open problem about closed affine manifolds is whether geodesic completeness (a geometric one-dimensional property) is equivalent to parallel volume (an algebraic n -dimensional property). This question was raised in 1963 by L. Markus [209] as a “Research Problem” in unpublished mimeographed lecture notes from the University of Minnesota (Problem 8, §6, p.58):

QUESTION. *Let M be a closed affine manifold. Then M is geodesically complete if and only if M has parallel volume.*

An affine manifold M has *parallel volume* if and only if it satisfies any of the following equivalent conditions:

- The orientable double-covering of M admits a parallel volume form (in the sense of §1.4.2 of Chapter 1);
- M admits a coordinate atlas where the coordinate changes are volume-preserving;
- M admits a refined $(\mathbf{SAff}(A), A)$ -structure, where $\mathbf{SAff}(A)$ denotes the subgroup $L^{-1}(\mathbf{SL}_{\pm}(\mathbb{R}^n))$ of volume-preserving linear transformations;
- For each $\phi \in \pi_1(M)$, the linear holonomy $L \circ \mathbf{h}(\phi)$ has determinant ± 1 .

11.1. The volume obstruction

EXERCISE 11.1.1. *Prove the equivalence of the conditions stated in the introduction to Chapter 11.*

The last condition suggests a topological interpretation. The composition of the linear holonomy representation $L \circ \mathbf{h}$ with the logarithm of the absolute value of the determinant

$$\pi_1(M) \xrightarrow{L \circ \mathbf{h}} \mathbf{GL}(E) \xrightarrow{|\det|} \mathbb{R}^+ \xrightarrow{\log} \mathbb{R}$$

defines an additive homomorphism $\nu_M \in \mathbf{Hom}(\pi_1(M), \mathbb{R}) \cong H^1(M; \mathbb{R})$ which we call the *volume obstruction*. M has parallel volume if and only if $\nu_M = 0$.

EXERCISE 11.1.2. *Suppose that M is a manifold with zero first Betti number. Then every affine structure on M must have parallel volume.*

One amusing corollary of this is that the *only* projective structures on the \mathbb{Q} -homology 3-sphere $\mathcal{S}_{\mathbb{Q}}^3$ defined in §6.2.3.1 are *complete affine* structures. Since $\mathcal{S}_{\mathbb{Q}}^3$ is covered by a 3-torus, the Markus conjecture for abelian holonomy (Smillie [253], Fried-Goldman-Hirsch [111]) implies that $\mathcal{S}_{\mathbb{Q}}^3$ must be complete, and must be covered by a complete affine nilmanifold (compare §11.2).

EXERCISE 11.1.3. *Classify all the projective structures on $\mathcal{S}_{\mathbb{Q}}^3$.*

Helmstetter’s theorem [147] that a left-invariant affine structure on a Lie group is complete \iff right-invariant volume forms are parallel is an “infinitesimal version” of Markus’s conjecture. (Compare also Goldman-Hirsch [132].)

The plausibility of Markus’s question seems to be one of the main barriers in constructing examples of affine manifolds. A purely topological consequence of this conjecture is that a compact affine manifold M with zero first Betti number $\beta_1(M)$ is covered by Euclidean space: in particular all of its higher homotopy groups vanish. Thus, if $\beta_1(M) = 0$ there should be no such structure on a nontrivial connected sum in dimensions greater than two. (In fact no affine structure — or projective structure — is presently known on a nontrivial connected sum.) Jo and Kim [163] resolve this question for *convex* affine manifolds.

11.2. Nilpotent holonomy

One of the first results on Markus’s question is its resolution in the case the affine holonomy group is nilpotent.

The structure theory of affine structures on closed manifolds with nilpotent holonomy is relatively well understood, due to the work of Smillie [253], Fried-Goldman-Hirsch [111] and Benoist [25, 28]. Smillie’s thesis develops the basic theory for affine structures with abelian holonomy, which Fried-Goldman-Hirsch extended to nilpotent holonomy, and Benoist extended to projective structures with nilpotent holonomy. The transition from nilpotent to solvable is much larger than the transition from abelian to nilpotent, and the next section discusses the few results in the solvable non-nilpotent case, due to Serge Dupont. Compare also Fried’s classification of closed similarity manifolds in §.

The key technique in the discussion of nilpotent holonomy is the structure theory of linear representations of nilpotent groups. The

guiding principle is that nilpotence ensures a nontrivial center, producing lots of commuting transformations. Specifically, elements of a nilpotent linear group have strongly compatible Jordan decompositions, which leads to invariant geometric structures for geometric manifolds with nilpotent holonomy. In another context this was used by Goldman [117] to give the first examples of 3-manifolds *without* flat conformal structures — for a geometric approach to these algebraic facts, see Thurston [266] and Ratcliffe [239].

The key to understanding nilpotent holonomy is the following algebraic fact:

THEOREM 11.2.1. *Let V be a vector space over \mathbb{C} and $\Gamma < \mathrm{GL}(V)$ a nilpotent group. Then $\exists k \in \mathbb{N}$ and Γ -invariant subspaces $V_i < V$ for $i = 1, \dots, k$ such that*

$$V = \bigoplus_{i=1}^k V_i$$

and homomorphisms $\Gamma \xrightarrow{\lambda_i} \mathbb{C}^$ such that for each $\gamma \in \Gamma$ and $i = 1, \dots, k$, the restriction $\gamma - \lambda_i \mathbb{I}$ to V_i is nilpotent. Furthermore there exists a basis of V_i such that each restriction $g|_{V_i}$ is upper-triangular with diagonal entry $\lambda_i(g)$.*

COROLLARY 11.2.2. *Let A be an affine space over \mathbb{C} and $\Gamma < \mathrm{Aff}(A)$. Then there exists a unique maximal Γ -invariant affine subspace $A_1 < A$ such that the restriction of Γ to A_1 is unipotent.*

A_1 is called the *Fitting subspace* in [111].

Since Γ preserves the affine subspace A_1 , it induces an affine action on the quotient space A/A_1 . Denote by V the vector space underlying A . Since the affine action on A/A_1 is radiant (it preserves the coset $A_1 < A$), we may describe A as an *affine direct sum*:

$$A = A_1 \oplus V_1$$

where $V_1 \subset V$ is an $L(\Gamma)$ -invariant linear subspace.

THEOREM 11.2.3 (Smillie [253], Fried-Goldman-Hirsch [111]). *Let M be a closed manifold with an affine structure whose affine holonomy group Γ is nilpotent. Let A_1 be its Fitting subspace and V_1 its linear complement as above. Then $\exists \gamma \in \Gamma$ such that the restriction $\gamma|_{V_1}$ is a linear expansion.*

As in [253, 111], the Markus conjecture for nilpotent holonomy follows:

COROLLARY 11.2.4. *Let M be a closed affine manifold whose affine holonomy group is nilpotent. Then M is complete if and only if it has parallel volume.*

PROOF. Suppose M has parallel volume. Theorem 11.2.3 guarantees an element $\gamma \in \Gamma$ whose linear part $L(\gamma)$ is an expansion on V_1 . Thus $V_1 = 0$, and $A = A_1$, that is, Γ is unipotent. Apply Theorem 8.4.1 to deduce that M is complete.

Conversely suppose M is complete, that is, the developing map $\widetilde{M} \xrightarrow{\text{dev}} A$ is a diffeomorphism. Then

$$M_1 := (\text{dev}^{-1}(A_1))/\Gamma \subset A/\Gamma \cong M$$

is a closed affine submanifold and $M_1 \hookrightarrow M$ is a homotopy-equivalence. Since both M_1 and M are aspherical,

$$\dim(A_1) = \dim(M_1) = \text{cd}(\Gamma) = \dim(M) = \dim(A),$$

$A_1 = A$ and $L(\Gamma)$ is unipotent, and hence volume-preserving. Thus M has parallel volume. \square

The geometric consequence of Theorem 11.2.3 is that the Γ -invariant decomposition

$$A_1 \hookrightarrow A \twoheadrightarrow V_1$$

defines two transverse affine foliations of M . The affine subspaces parallel to A_1 define the leaves of a foliation \mathcal{F}^u of M . The leaves of \mathcal{F}^u are affine submanifolds of M with unipotent holonomy. The affine subspaces parallel to V_1 define the leaves of a foliation \mathcal{F}^R of M . The leaves of \mathcal{F}^R are affine submanifolds of M with radiant affine structure.

EXERCISE 11.2.5. *Suppose that M is closed. Show that each leaf of \mathcal{F}^u is complete.*

11.3. Smillie's nonexistence theorem

THEOREM 11.3.1 (Smillie [255]). *Let M be a closed affine manifold with parallel volume. Then the affine holonomy homomorphism cannot factor through a free group.*

This theorem can be generalized much further — see Smillie [255] and Goldman-Hirsch [132].

COROLLARY 11.3.2 (Smillie [255]). *Let M be a closed manifold whose fundamental group is a free product of finite groups (for example, a connected sum of manifolds with finite fundamental group). Then M admits no affine structure.*

PROOF OF COROLLARY 11.3.2 ASSUMING THEOREM 11.3.1.

Suppose M has an affine structure. Since $\pi_1(M)$ is a free product of finite groups, the first Betti number of M is zero. Thus M has parallel volume. Furthermore if $\pi_1(M)$ is a free product of finite groups, there

exists a free subgroup $\Gamma \subset \pi_1(M)$ of finite index. Let \hat{M} be the covering space with $\pi_1(\hat{M}) = \Gamma$. Then the induced affine structure on \hat{M} also has parallel volume contradicting Theorem. \square

PROOF OF THEOREM 11.3.1. Let M be a closed affine manifold modeled on an affine space E , $\widetilde{M} \xrightarrow{\Pi} M$ a universal covering, and

$$\left(\widetilde{M} \xrightarrow{\text{dev}} E, \pi \xrightarrow{\text{hol}} \text{Aff}(E) \right)$$

a development pair. Suppose that M has parallel volume and that there is a free group Π through which the affine holonomy homomorphism h factors:

$$\pi \xrightarrow{\phi} \Pi \xrightarrow{\bar{h}} \text{Aff}(E)$$

Choose a graph G with fundamental group Π ; then there exists a map $f : M \rightarrow G$ inducing the homomorphism

$$\pi = \pi_1(M) \xrightarrow{\phi} \pi_1(G) = \Pi.$$

By general position, there exist points $s_1, \dots, s_k \in G$ such that f is transverse to s_i and the complement $G - \{s_1, \dots, s_k\}$ is connected and simply connected. Let H_i denote the inverse image $f^{-1}(s_i)$ and let $H = \cup_i H_i$ denote their disjoint union. Then H is an oriented closed smooth hypersurface such that the complement $M - H \subset M$ has trivial holonomy. Let $M|H$ denote the manifold with boundary obtained by *splitting* M along H ; that is, $M|H$ has two boundary components H_i^+, H_i^- for each H_i and there exist diffeomorphisms $H_i^+ \xrightarrow{g_i} H_i^-$ (generating Π) such that M is the quotient of $M|H$ by the identifications g_i . There is a canonical diffeomorphism of $M - H$ with the interior of $M|H$.

Let ω_E be a parallel volume form on E ; then there exists a parallel volume form ω_M on M such that

$$\Pi^* \omega_M = \text{dev}^* \omega_E.$$

Since $H^n(E) = 0$, there exists an $(n-1)$ -form η on E such that $d\eta = \omega_E$. For any immersion $S \xrightarrow{f} E$ of an oriented closed $(n-1)$ -manifold S , the integral

$$\int_S f^* \eta$$

is independent of the choice of η satisfying $d\eta = \omega_E$. Since $H^{n-1}(E)$, any other η' must satisfy $\eta' = \eta + d\theta$ and

$$\int_S f^* \eta' - \int_S f^* \eta = \int_S d(f^* \theta) = 0.$$

Since $M - H$ has trivial holonomy there is a developing map

$$M - H \xrightarrow{\text{dev}} E$$

and its restriction to $M - H$ extends to a developing map $M|H \xrightarrow{\text{dev}} E$ such that

$$\text{dev}|_{H_i^+} = \bar{h}(g_i) \circ \text{dev}|_{H_i^-}$$

and the normal orientations of H_i^+, H_i^- induced from $M|H$ are opposite. Since $h(g_i)$ preserves the volume form ω_E ,

$$d(h(g_i)^*\eta) = d(\eta) = \omega$$

and

$$\int_{H_i^+} \text{dev}^*\eta = \int_{H_i^+} \text{dev}^*h(g_i)^*\eta = - \int_{H_i^-} \text{dev}^*\eta$$

since the normal orientations of H_i^\pm are opposite. We now compute the ω_M -volume of M :

$$\begin{aligned} \text{vol}(M) &= \int_M \omega_M = \int_{M|H} \text{dev}^*\omega_E \\ &= \int_{\partial(M|H)} \eta = \sum_{i=1}^k \left(\int_{H_i^+} \eta + \int_{H_i^-} \eta \right) = 0, \end{aligned}$$

a contradiction. □

One basic method of finding a primitive η for ω_E involves a radiant vector field ρ . Since ρ expands volume, specifically,

$$d\iota_\rho\omega_E = n\omega_E,$$

and

$$\eta = \frac{1}{n} \iota_\rho\omega_E$$

is a primitive for ω_E . An affine manifold is *radiant* if and only if it possesses a radiant vector field if and only if the affine structure comes from an $(E, \mathbf{GL}(E))$ -structure if and only if its affine holonomy has a fixed point in E . The following result generalizes the above theorem:

THEOREM 11.3.3 (Smillie). *Let M be a closed affine manifold with a parallel exterior differential k -form which has nontrivial de Rham cohomology class. Suppose \mathcal{U} is an open covering of M such that for each $U \in \mathcal{U}$, the affine structure induced on U is radiant. Then $\dim \mathcal{U} \geq k$; that is, there exist $k + 1$ distinct open sets*

$$U_1, \dots, U_{k+1} \in \mathcal{U}$$

such that the intersection

$$U_1 \cap \cdots \cap U_{k+1} \neq \emptyset.$$

(Equivalently the nerve of \mathcal{U} has dimension at least k .)

A published proof of this theorem can be found in Goldman-Hirsch [132].

Using these ideas, Carrière, dAl’bo and Meignez [57] have proved that a nontrivial Seifert 3-manifold with hyperbolic base cannot have an affine structure with parallel volume. This implies that the 3-dimensional Brieskorn manifolds $M(p, q, r)$ with

$$p^{-1} + q^{-1} + r^{-1} < 1$$

admit no affine structure whatsoever. (Compare Milnor [223].)

There is a large class of discrete groups Γ for which every affine representation $\Gamma \rightarrow \text{Aff}(E)$ is conjugate to a representation factoring through $\text{SL}(E)$, that is,

$$\Gamma \rightarrow \text{SL}(E) \subset \text{Aff}(E).$$

For example finite groups have this property, and the above theorem gives an alternate proof that the holonomy of a compact affine manifold must be infinite. Another class of groups having this property are the *Margulis-superrigid groups*, that is, irreducible lattices Γ in semisimple Lie groups G of \mathbb{R} -rank greater than one (for example, $\text{SL}(n, \mathbb{Z})$ for $n > 2$). Margulis proved [207] that every unbounded finite-dimensional linear representation of Γ extends to a representation of G . It then follows that the affine holonomy of a compact affine manifold cannot factor through a Margulis-superrigid group. However, since $\text{SL}(n; \mathbb{R})$ does admit a left-invariant $\mathbb{R}\text{P}^{n^2-1}$ -structure, it follows that if $\Gamma \subset \text{SL}(n; \mathbb{R})$ is a torsion-free cocompact lattice, then there exists a compact affine manifold with holonomy group $\Gamma \times \mathbb{Z}$ although Γ itself is not the holonomy group of an affine structure.

11.4. Fried’s classification of closed similarity manifolds

Fried [106] gives a sharp classification of closed similarity manifolds; this was announced earlier by Kuiper [188], although the proof contains a gap. Later Reischer and Vaisman [270] proved this, using a completely different set of ideas. Miner [227] extended Fried’s theorem to manifolds modeled on the Heisenberg group and its group of similarity transformations. Recently this has been extended to the boundary geometry of any rank one symmetric space by Raphaël Alexandre [4]. Later, in §??, the ideas in Fried’s proof are related to Thurston’s parametrization of $\mathbb{C}\text{P}^1$ -structures and the Kulkarni-Pinkall theory of flat conformal structures [194, 195].

11.4.1. Completeness versus radiance. Fried's theorem is a prototype of a theorem about geometric structures on *closed manifolds*. Here $X = \mathbf{E}^n$ and $G = \text{Sim}(\mathbf{E}^n)$. Namely, Fried shows that a (G, X) -structure on a closed manifold M must reduce to one of two special types, corresponding to *subgeometries* $(G', X') \rightsquigarrow (G, X)$. Specifically, a closed similarity manifold must be one of the following two types:

- A *Euclidean manifold* where $X' = X = \mathbf{E}^n$ and

$$G' = \text{Isom}(\mathbf{E}^n) \hookrightarrow \text{Sim}(\mathbf{E}^n).$$

This is precisely the case when the underlying affine structure on M is complete;

- A finite quotient of a *Hopf manifold* where $X' = \mathbf{E}^n \setminus \{\mathbf{0}\}$ and $G' = \text{Sim}_0(\mathbf{E}^n) \hookrightarrow \text{Sim}(\mathbf{E}^n)$, the group of linear similarity (or conformal) transformations of \mathbf{E}^n . This is precisely the case when the underlying affine structure is incomplete.

The complete case is easy to handle, since in that case Γ acts freely, and any similarity transformation which is not isometric must fix a point. When M is incomplete, very little can be said in general, and the compactness hypothesis must be crucially used.

EXERCISE 11.4.1. *Prove that a complete similarity manifold is a Euclidean manifold, and diffeomorphic to a finite quotient of a product $\mathbb{T}^r \times \mathbf{E}^{n-r}$.*

The other extreme — radiant similarity manifolds — were discussed in §6.5.2.1 of Chapter 6.

The recurrence of an incomplete geodesic on a compact manifold guarantees a divergent sequence in the affine holonomy group Γ . This *holonomy sequence* converges to a singular projective transformation ϕ as in §2.6. The condition that $\Gamma \subset \text{Sim}(\mathbf{E}^n)$ strongly restricts ϕ ; in particular it has *rank one* or its limits are *proximal*, in that most points approach a single point, which Fried shows must lie in \mathbf{E}^n . From this he deduces radiance, and finds that the structure is modeled on $\mathbf{E}^n \setminus \{\mathbf{0}\}$. We closely follow Fried's proof (which we highly recommend reading), but insert more details and coordinate the notation with the rest of this document.

11.4.2. Canonical metrics and incompleteness. Choose a Euclidean metric \mathbf{g}_E on \mathbf{E}^n ; the pullback $\text{dev}^* \mathbf{g}_E$ is a Euclidean metric on \widetilde{M} . Unless M is a Euclidean manifold, this metric is *not* invariant under π . rather it transforms by the *scale factor homomorphism*: $\pi \xrightarrow{\lambda^{\text{oh}}} \mathbb{R}^+$:

$$(54) \quad \gamma^*(\text{dev}^* \mathbf{g}_E) = \lambda \circ \mathbf{h}(\gamma) \text{dev}^* \mathbf{g}_E.$$

defined in §1.4.1.

EXERCISE 11.4.2. *Relate the scale factor $\lambda \circ \mathbf{h}$ to the volume obstruction ν_M .*

Unless M is Euclidean, then $\text{dev}^* \mathbf{g}_E$ is incomplete. Thus we assume that $(\widetilde{M}, \text{dev}^* \mathbf{g}_E)$ is an *incomplete Euclidean manifold* with distance function $\widetilde{M} \times \widetilde{M} \xrightarrow{\widetilde{d}_E} \mathbb{R}$, *non-bijective* developing map $\widetilde{M} \xrightarrow{\text{dev}} E$ and *nontrivial* scale factor homomorphism $\pi_1(M) \xrightarrow{\lambda \circ \mathbf{h}} \mathbb{R}^+$.

We begin with some general facts about an incomplete Euclidean manifold N with trivial holonomy. We apply these facts to the case when N is the universal covering \widetilde{M} of a compact incomplete similarity manifold M .

EXERCISE 11.4.3. *Let N be a Euclidean manifold with trivial holonomy. Choose a developing map $N \xrightarrow{\text{dev}} E$. Let $B \subset N$ be an open subset. The following conditions are equivalent:*

- B is an open ball in N , that is, $\exists c \in N, r > 0$ such that $B = \mathbf{B}_r(c)$.
- B develops to an open ball in E , that is, $\exists c \in E, r > 0$ such that the restriction $\text{dev}|_B$ is a diffeomorphism $B \rightarrow \mathbf{B}_r(c) \subset E$;
- B is the exponential image of a metric ball in the tangent space $\mathbf{T}_c N$, that is, $\exists c \in N, r > 0$ such that the restriction $\text{Exp}|_{\mathbf{B}_r(\mathbf{0}_c)}$ is a diffeomorphism $\mathbf{B}_r(\mathbf{0}_c) \rightarrow B$.

Under these conditions, the maps

$$\mathbf{B}_r(\mathbf{0}_c) \xrightarrow{\text{Exp}_c} B \xrightarrow{\text{dev}} \mathbf{B}_r(\text{dev}(c))$$

are isometries with respect to the restrictions of the Euclidean metrics on $\mathbf{T}_c N$, N and E , respectively.

DEFINITION 11.4.4. *A maximal ball in N is an open ball which is maximal among open balls with respect to inclusion.*

EXERCISE 11.4.5. *A Euclidean manifold with trivial holonomy is complete (that is, is isomorphic to Euclidean space) if and only if no ball is maximal.*

EXERCISE 11.4.6. *Suppose N is an incomplete Euclidean manifold with trivial holonomy.*

- Let $B \subset N$ be a maximal ball and let c be its center. Then B is maximal among open balls centered at c .
- Every open ball lies in a maximal ball.
- Every $x \in N$ is the center of a unique maximal ball $\mathcal{B}(x)$.

- For each x , not every point on $\partial\mathcal{B}(x)$ is visible from x .

DEFINITION 11.4.7. Let N be an incomplete Euclidean manifold with trivial holonomy. For each $x \in N$, let $R(x) < \infty$ be the radius of the maximal ball $\mathcal{B}(x) \subset N$ centered at x .

EXERCISE 11.4.8. The maximal ball $\mathcal{B}(x) = \mathbf{B}_{R(x)}(x)$. Moreover $R(x)$ is the supremum of r such that $\mathbf{B}_r(\mathbf{0}_x) \subset \mathcal{E}_x$, where $\mathcal{E}_x \subset \mathbb{T}_x N$ denotes the domain of \mathbf{Exp}_x defined in §8.3 of Chapter 8.

LEMMA 11.4.9. The function R is Lipschitz:

$$(55) \quad |R(x) - R(y)| \leq \tilde{\mathbf{d}}(x, y)$$

if $x, y \in N$ are sufficiently close. In particular R is continuous.

PROOF. Suppose that $x \in N$ and ϵ such that

$$\epsilon < \sup (R(x), R(y)).$$

Choose $r < R(x)$ so that $\mathbf{B}_r(\mathbf{0}_x) \subset \mathcal{E}_x$. First we show that if $\tilde{\mathbf{d}}(x, y) < \epsilon$, then

$$(56) \quad r < \tilde{\mathbf{d}}(x, y) + R(y)$$

Choose $u \in \mathcal{B}(x)$ such that $\tilde{\mathbf{d}}(x, u) = r$. Suppose that $\tilde{\mathbf{d}}(x, y) < \epsilon$. Then closed ball $\mathbf{B}_r(x)$ lies in the convex set $\mathcal{B}(x)$ which also contains y . Thus $u \in \partial\mathbf{B}_r(x)$ is visible from y , whence

$$\tilde{\mathbf{d}}(y, u) < R(y).$$

Thus

$$r = \tilde{\mathbf{d}}(x, u) \leq \tilde{\mathbf{d}}(x, y) + \tilde{\mathbf{d}}(y, u) < \tilde{\mathbf{d}}(x, y) + R(y),$$

proving (56). Taking the supremum over r yields:

$$R(x) < \tilde{\mathbf{d}}(x, y) + R(y),$$

so $R(x) - R(y) < \tilde{\mathbf{d}}(x, y)$ if $\tilde{\mathbf{d}}(x, y) < \epsilon$. Similarly, symmetry of $\tilde{\mathbf{d}}$ implies that $R(x) - R(y) < \tilde{\mathbf{d}}(x, y)$ if $\tilde{\mathbf{d}}(x, y) < \epsilon$ which implies (55). \square

We return to the case that M is a compact incomplete similarity manifold. Choose a universal covering $N \xrightarrow{\Pi} M$, a developing map $N \xrightarrow{\text{dev}} \mathbf{E}$, and a holonomy representation $\pi_1(M) \xrightarrow{\mathbf{h}} \mathbf{Sim}(\mathbf{E}^n)$. If $\phi \in \pi_1(M)$ is a deck transformation (also denoted $N \xrightarrow{\phi} N$), then $\phi(\mathbf{B}_{R(\tilde{p})})$ is a maximal ball at $\phi(\tilde{p})$, so:

$$(57) \quad R(\phi\tilde{p}) = \lambda \circ \mathbf{h}(\phi)R(\tilde{p})$$

This leads to a natural conformal Riemannian structure on N which descends to a conformal Riemannian structure on M .

This will be the canonical Riemannian structure on a radiant similarity manifold. If M is closed and incomplete, then M is finitely covered by a Hopf manifold M' homeomorphic to $S^{n-1} \times S^1$. The induced Riemannian structure on M' is the Cartesian product of a spherical metric on S^{n-1} with a Euclidean metric on S^1 . By (57), the Riemannian metric $\tilde{\mathbf{g}}$ on \tilde{M} defined by:

$$(58) \quad \tilde{\mathbf{g}}(\tilde{p}) := R(\tilde{p})^{-1} \text{dev}^* \mathbf{g}_E$$

is $\pi_1(M)$ -invariant. Therefore $\tilde{\mathbf{g}}$ passes down to a Riemannian metric \mathbf{g}_M on M , that is, $\Pi^* \mathbf{g}_M = \tilde{\mathbf{g}}$.

The Riemannian structure \mathbf{g}_M has the property that its unit ball is maximal inside the domain \mathcal{E} of the exponential map Exp . When M is closed, even more is true:

PROPOSITION 11.4.10. *Let M be a compact incomplete similarity manifold with universal covering $N \xrightarrow{\Pi} M$. Then $\exists \xi \in \text{Vec}(M)$ which is Π -related to a vector field $\tilde{\xi} \in \text{Vec}(N)$ such that:*

- $\|\xi\|_{\mathbf{g}_M} = \|\tilde{\xi}\|_{\tilde{\mathbf{g}}} = 1$;
- *The halfspace*

$$\mathcal{H}_x := \{\mathbf{v} \in \mathbb{T}_x N \mid \tilde{\mathbf{g}}(\mathbf{v}, \xi) < 1\}$$

lies in \mathcal{E}_x for all $x \in N$.

In particular

$$H_x := \text{Exp}_x(\mathcal{H}_x) \subset N$$

is a natural halfspace neighborhood of x .

By analyzing H_x , we shall prove that ξ is a radiant vector field and M is (covered by) a Hopf manifold. A key ingredient is a *holonomy sequence* $\mathbf{h}(\phi_{ij}) \in \text{Sim}(\mathbf{E}^n)$, where $\phi_{ij} \in \pi_1(M)$, which contracts to $\mathbf{0}$ as $j \nearrow \infty$ (Proposition 11.4.13). The proof of Proposition 11.4.10 will be given in §11.4.4, following several preliminary lemmas needed in the proof.

11.4.3. Incomplete geodesics recur. Fried makes a detailed analysis of an incomplete geodesic $[0, 1) \xrightarrow{\gamma} M$. That is, $\gamma(t) = \text{Exp}(t\mathbf{v})$, where $\mathbf{v} \in \mathbb{T}_x M$ but $t\mathbf{v} \in \mathcal{E}_x \iff t < 1$.

Since M is compact (and $[0, 1)$ isn't compact), the path $\gamma(t)$, *accumulates*. That is, for some sequence $t_n \nearrow 1$ a sequence $\gamma(t_n) \in M$ converges in M as $n \nearrow +\infty$. Denote

$$(59) \quad p := \lim_{n \rightarrow +\infty} \gamma(t_n)$$

Next, we use the recurrence of $\gamma(t)$ to obtain a *holonomy sequence* converging to a singular projective transformation as in §2.6. To that

end, we pass to a specific universal covering space and a developing map. Employ $p \in M$ as a basepoint to define a universal covering space $\widetilde{M} \xrightarrow{\Pi} M$. The total space \widetilde{M} comprises relative homotopy classes of paths $[0, T] \xrightarrow{\gamma} M$ with $\gamma(0) = p$ and the projection is:

$$\begin{aligned} \widetilde{M} &\xrightarrow{\Pi} M \\ [\gamma] &\longrightarrow \gamma(T) \end{aligned}$$

The constant path defines a basepoint $\tilde{p} \in \widetilde{M}$ with $\Pi(\tilde{p}) = p$. The group of deck transformations is $\pi_1(M, p)$ consisting of relative homotopy classes of loops in M based at p .

Lift the incomplete geodesic $[0, 1) \xrightarrow{\gamma} M$ to an incomplete geodesic $[0, 1) \xrightarrow{\tilde{\gamma}} N$ so that

$$\lim_{n \rightarrow +\infty} \tilde{\gamma}(t_n) = \tilde{p}$$

and let $\tilde{x} := \tilde{\gamma}(0)$ be the initial endpoint of $\tilde{\gamma}$ and $\tilde{\mathbf{v}} := \tilde{\gamma}'(0) \in \mathbb{T}_{\tilde{x}}N$ the initial velocity.

Choose a developing map $\widetilde{M} \xrightarrow{\text{dev}} \mathbb{E}^n$. A Euclidean metric tensor $\mathbf{g}_{\mathbb{E}}$ on \mathbb{E} induces a Euclidean metric tensor $\text{dev}^*\mathbf{g}$ on N . By rescaling, we may assume that $\text{dev}^*\mathbf{g}(\tilde{\mathbf{v}}, \tilde{\mathbf{v}}) = 1$. Denote the corresponding distance function by $N \times N \xrightarrow{d} \mathbb{R}$. Choose a coordinate patch $U \ni p$ such that the restriction $\text{dev}|_{\tilde{U}}$ is injective, where $\tilde{U} \subset \widetilde{M}$ is the component of $\Pi^{-1}(U)$ containing \tilde{p} . Choose $\epsilon > 0$ such that:

- $\mathcal{B}_\epsilon := \mathbb{B}_\epsilon(\mathbf{0}_{\tilde{p}}) \subset \mathcal{E}_{\tilde{p}}$;
- The ball $B := \text{Exp}_{\tilde{p}}(\mathcal{B}_\epsilon)$ lies in \tilde{U} ;
- $\epsilon < \frac{1}{2}$.

In particular, the restriction $\text{dev}|_B$ is injective.

LEMMA 11.4.11. *Let $N \xrightarrow{R} \mathbb{R}^+$ be the radius function (defined in 11.4.7). Then*

$$R(\tilde{\gamma}(t)) = 1 - t.$$

EXERCISE 11.4.12. *Prove Lemma 11.4.11.*

After possibly passing to a subsequence, (59) implies that $\gamma(t_i) \in B$. Let

$$s_i := d(\gamma(t_i), p)$$

and $[0, s_i] \xrightarrow{\eta_i} M$ the unit-speed geodesic in B with $\eta_i(0) = p$ and $\eta_i(s_i) = \gamma(t_i)$. Lift η_i to

$$[0, s_i] \xrightarrow{\tilde{\eta}_i} \widetilde{M}$$

with $\tilde{\eta}_i(s_i) = \tilde{\gamma}(t_i)$. Let $\tilde{p}_i := \tilde{\eta}_i(0)$. For $i < j$ define

$$\phi_{ij} := [\eta_j^{-1} \star \gamma|_{[t_i, t_j]} \star \eta_i] \in \pi_1(M, p).$$

A crucial fact is that the scale factors decrease to 0 along the incomplete geodesic $\gamma(t)$. In particular the limit of the holonomy sequence is a singular projective transformation whose image is a single point.

PROPOSITION 11.4.13. *Fix $i \in \mathbb{N}$ and $\delta > 0$. Then $\exists J(i)$ such that the scale factor $\lambda \circ \mathbf{h}(\phi_{ij}) < \delta$ for $j \geq J(i)$.*

PROOF. Denote the $\tilde{\mathbf{g}}$ -length of $\tilde{\eta}_i$ by

$$l_i := \tilde{\mathbf{d}}(\tilde{p}_i, \text{dev } \tilde{\gamma}(t_i)).$$

First, we claim that:

$$(60) \quad \epsilon > \frac{l_i}{l_i + R(\tilde{\gamma}(t_i))}.$$

If $0 \leq s \leq s_i$, then $\tilde{\mathbf{d}}(\tilde{\gamma}(t_i), \tilde{\eta}_i(s)) \leq l_i$. Lemma 11.4.9 implies:

$$R(\tilde{\eta}_i(s)) \leq R(\tilde{\gamma}(t_i)) + l_i$$

so the $\tilde{\mathbf{g}}$ -length of $\tilde{\eta}_i$ equals:

$$\begin{aligned} \tilde{\mathbf{d}}(\tilde{p}_i, \tilde{\gamma}(t_i)) &= \int_0^{s_i} \frac{(\tilde{\eta}_i)^* ds}{R(\tilde{\eta}_i(s))} \\ &\geq \frac{l_i}{l_i + R(\tilde{\gamma}(t_i))} \end{aligned}$$

Thus

$$\frac{l_i}{l_i + R(\tilde{\gamma}(t_i))} \leq \tilde{\mathbf{d}}(\tilde{p}_i, \tilde{\gamma}(t_i)) = \mathbf{d}(x, \gamma_i(t_i)) < \epsilon$$

as desired, proving (60). Next we prove:

$$(61) \quad l_i < 2\epsilon(1 - t_i).$$

In general, $\epsilon > l/(l + R)$ implies that $l < R\epsilon/(1 - \epsilon)$. Furthermore $\epsilon < 1/2$ implies that $\epsilon/(1 - \epsilon)R < 2\epsilon R$ so

$$l_i < 2\epsilon R(\tilde{\gamma}(t_i)) = 2\epsilon R$$

by Lemma 11.4.11, thereby establishing (61).

LEMMA 11.4.14.

$$1 - 2\epsilon < \frac{R(\tilde{p}_i)}{1 - t_i} < 1 + 2\epsilon$$

PROOF. Lemma 11.4.9 implies

$$|R(\tilde{p}_i) - R(\tilde{\gamma}(t_i))| \leq d(\text{dev}(\tilde{p}_i), \text{dev}(\tilde{\gamma}(t_i))) = l_i.$$

Lemma 11.4.11 and (61) together imply

$$|R(\tilde{p}_i) - (1 - t_i)| < 2\epsilon(1 - t_i).$$

Now divide by $1 - t_i$. □

LEMMA 11.4.15. For $i < j$,

$$\frac{R(\tilde{x}_j)}{R(\tilde{p}_i)} < \frac{1 + 2\epsilon}{1 - 2\epsilon} \frac{1 - t_j}{1 - t_i}$$

PROOF. Apply Lemma 11.4.14 to obtain:

$$(62) \quad R(\tilde{x}_j) < (1 + 2\epsilon)(1 - t_j)$$

and

$$(1 - 2\epsilon)(1 - t_i) < R(\tilde{p}_i)$$

that is,

$$(63) \quad \frac{1}{R(\tilde{p}_i)} < \frac{1}{(1 - 2\epsilon)(1 - t_i)}.$$

Multiplying (62) and (63) implies Lemma 11.4.15. □

Since $t_j \nearrow 1$, for any fixed i , there exists $J(i)$ such that $j > J(i)$ implies

$$\frac{1 + 2\epsilon}{1 - 2\epsilon} \frac{1 - t_j}{1 - t_i} < \delta.$$

Since $\tilde{x}_j = \phi_{ij} \tilde{p}_i$, (57) and (54) imply:

$$\frac{R(\tilde{x}_j)}{R(\tilde{p}_i)} = \lambda \circ h(\phi_{ij}).$$

Now apply Lemma 11.4.15 to complete the proof of Proposition 11.4.13. □

Recall from Exercise 2.6.6 of Chapter 2, that a sequence of similarity transformations accumulates to either:

- The *zero* affine transformation (undefined at the ideal hyperplane, otherwise constant);
- A singular projective transformation of rank one, taking values at an ideal point.

The contraction of scale factors (Proposition 11.4.13) implies that the second case cannot occur.

11.4.4. Existence of halfspace neighborhood. The contraction of scale factors (Proposition 11.4.13) easily implies the existence of halfspace neighborhoods (Proposition 11.4.10).

The proof uses the following elementary fact in Euclidean geometry:

EXERCISE 11.4.16. *Let $y \in \mathbb{E}^n$ be a point and $\mathbf{v} \in \mathbb{T}_y \mathbb{E}^n \xleftrightarrow{\sim} \mathbb{R}^n$ be a tangent vector. Choosing coordinates, we may assume that $\text{Exp}_y(\mathbf{v}) = \mathbf{0}$ is the origin. Let \mathcal{S}_t be a one-parameter family of homotheties approaching zero:*

$$\begin{aligned} \mathbb{E}^n &\xrightarrow{\mathcal{S}_t} \mathbb{E}^n \\ p &\longmapsto e^{-t}p \end{aligned}$$

Let B be the ball centered at y of radius $R = \|\mathbf{v}\|$:

$$B := \{x \in \mathbb{E}^n \mid d(y, x) < R\}$$

If $t_n \nearrow \infty$, then the union of $\mathcal{S}_{t_n}(B)$ is the halfspace H containing y and orthogonal to the line segment $y\overleftarrow{\mathbf{0}}$:

$$H(y, \mathbf{0}) := \{y + \mathbf{w} \mid \mathbf{w} \cdot \mathbf{v} < R\}.$$

Let \mathcal{A}_p denote the extension of $\text{dev} \circ \text{Exp}_p$ to $\mathbb{T}_p N \rightarrow \mathbb{A}$ described in Proposition 8.3.6.

CONCLUSION OF PROOF OF PROPOSITION 11.4.10. Let $x \in M$ and lift to $\tilde{x} \in N$. Since M is incomplete, a maximal ball $\mathcal{B}(\tilde{x}) \subset \mathcal{E}_{\tilde{x}}$ exists, and denote its radius by $R := R(\tilde{x})$. Furthermore $\partial\mathcal{B}(\tilde{x})$ contains a vector $\tilde{\mathbf{v}}$ of length R such that $\tilde{\mathbf{v}} \notin \mathcal{E}_{\tilde{x}}$ but $t\tilde{\mathbf{v}} \in \mathcal{E}_{\tilde{x}}$ for $|t| < 1$. (Soon we shall see that $\tilde{\mathbf{v}}$ is unique, and will be the value $\xi(\tilde{x})$.)

Apply this construction to go from the incomplete geodesic

$$\gamma(t) := \text{Exp}_x(t\mathbf{v}), \quad (0 \leq t < R),$$

in M to a holonomy sequence $\gamma_{ij} \in \pi_1(M, p)$ satisfying Proposition 11.4.13.

Define the halfspace

$$\begin{aligned} \mathcal{H}_{\tilde{x}} &:= \{Y \in \mathbb{T}_{\tilde{x}} N \mid \text{dev}^* \mathbf{g}_{\mathbb{E}}(\tilde{\mathbf{v}}, Y) < R\} \\ &= \{Y \in \mathbb{T}_{\tilde{x}} N \mid \mathbf{g}_N(\tilde{\mathbf{v}}, Y) < 1\}. \end{aligned}$$

We show $\mathcal{H}_{\tilde{x}} \subset \mathcal{E}_{\tilde{x}}$. Suppose that Y is a tangent vector at \tilde{x} such that

$$\text{dev}^* \mathbf{g}_{\mathbb{E}}(\tilde{\mathbf{v}}, Y) < R.$$

By Proposition 11.4.13, for $j \gg 1$ and fixed $i < j$, the holonomy $\mathbf{h}(\gamma_{ij})$ is a very sharp contraction with rotational component close to the identity — that is, it's very close to a strong homothety about $\mathbf{0}$. By Exercise 11.4.16 above, $(D\gamma_{ij})_{\tilde{x}}$ maps Y to a vector in $\mathcal{B}_{\gamma_{ij}(\tilde{x})} \subset \mathcal{E}_{\gamma_{ij}(\tilde{x})}$

for $j \gg i$. Thus $(D\gamma_{ij})_{\tilde{x}}(Y)$ is visible from $\gamma_{ij}(\tilde{x})$ so Y is visible from \tilde{x} , as claimed.

In particular every point in $\partial\mathcal{B}(\tilde{x}) \setminus \{X\}$ lies in $\mathcal{H}(\tilde{x})$, and therefore \tilde{x} uniquely determines X , so we write $X =: \xi(\tilde{x})$. □

11.4.5. Visible points on the boundary of the halfspace.

Bounding the halfspace $\mathcal{H}_{\tilde{x}}$ is the affine hyperplane

$$\partial\mathcal{H}_{\tilde{x}} := \{Y \in \mathbb{T}_{\tilde{x}} \mid \mathfrak{g}_N(\tilde{\mathbf{v}}, Y) < 1\},$$

which decomposes as the disjoint union of two subsets:

- The *visible set* $\partial\mathcal{H}_{\tilde{x}} \cap \mathcal{E}_{\tilde{x}}$;
- The *invisible set* $\partial\mathcal{H}_{\tilde{x}} \setminus \mathcal{E}_{\tilde{x}}$.

The invisible set is nonempty, since it contains $\xi(\tilde{x})$.

LEMMA 11.4.17. *The visible set $\partial\mathcal{H}_{\tilde{x}} \cap \mathcal{E}_{\tilde{x}}$ is nonempty.*

PROOF. Suppose every point of $\partial\mathcal{H}_{\tilde{x}}$ is invisible. Then $\mathcal{H}_{\tilde{x}}$ is a closed subset of $\mathcal{E}_{\tilde{x}}$. By construction, $\mathcal{H}_{\tilde{x}}$ is open and connected, so $\mathcal{H}_{\tilde{x}} = \mathcal{E}_{\tilde{x}}$.

Thus the corresponding subset $H_{\tilde{x}} = \text{Exp}(\mathcal{H}_{\tilde{x}}x)$ of N equals all of N , so M is a quotient of a halfspace. The contradiction now follows from the following exercise. □

EXERCISE 11.4.18. *Let $H \subset \mathbb{E}^n$ be an open halfspace, and let $\Gamma < \text{Sim}(\mathbb{E}^n)$ be a discrete subgroup stabilizing H and acting properly on H . Then the quotient $\Gamma \backslash H$ is not compact.*

11.4.6. The invisible set is affine and locally constant. This uses the variation of $H_{\tilde{x}}$ as \tilde{x} varies. The key is Lemma 1 of Fried [106], which says that if $y = \text{Exp}_{\tilde{x}}(Y)$, where $Y \in \partial\mathcal{H}_{\tilde{x}} \cap \mathcal{E}_{\tilde{x}}$ is a visible vector, then $\xi(\tilde{x})$ “is also invisible” from y . That is, the vector in $\mathbb{T}_y N$ whose parallel transport $\mathbb{P}_{y,x}$ to x equals $X \in \mathbb{T}_{\tilde{x}}(N)$ is invisible from y .

Now apply the strong contraction γ_{ij} to show that the halfplane $H(y)$ is moved closer and closer to $\mathbf{0}$. If it contains $\mathbf{0}$, then $X = \xi(x)$ is visible, a contradiction. If $\mathbf{0} \notin \overline{H(y)}$, then for $j \gg i$, the maximal ball $\mathcal{B}(y)$ is contained in a visible halfspace from x , contradicting maximality. Thus $\xi(\tilde{x})$ is also invisible from y .

This shows that \tilde{x} uniquely determines the vector X , and the map $\tilde{x} \mapsto X$ is locally constant.

EXERCISE 11.4.19. *Prove this map is affine and deduce that the invisible subspace $\partial\mathcal{H}_{\tilde{x}} \setminus \mathcal{E}_{\tilde{x}}$ is an affine subspace.*

Now we return to the case that M is a closed affine manifold with holonomy covering space N .

Fried concludes the proof by observing now that the vector field corresponds to an affine projection from the ambient affine space to the invisible subspace $\partial\mathcal{H}_{\tilde{x}} \setminus \mathcal{E}_{\tilde{x}}$. Since it is invariant under deck transformations it descends to a vector field on the compact quotient M .

Recall the *divergence* of a vector field (see §1.7.2) which measures the infinitesimal distortion of volume:

EXERCISE 11.4.20. *The divergence of the vector field X equals $\dim\partial\mathcal{H}_{\tilde{x}} \setminus \mathcal{E}_{\tilde{x}}$. Since M is closed, the vector field has divergence zero.*

Thus $\partial\mathcal{H}_{\tilde{x}} \cap \mathcal{E}_{\tilde{x}}$ is a single point, and M is radiant, as desired.

CHAPTER 12

Hyperbolicity

The opposite of geodesic completeness is *hyperbolicity* in the sense of Vey [275, 274] and Kobayashi [179, 177], which is equivalent to the following notion: An affine manifold M is *completely incomplete* if and only if every affine map $\mathbb{R} \rightarrow M$ is constant, that is, M admits *no* complete geodesic. As noted by the author (see Kobayashi [179]), the combined results of Kobayashi [179], Wu [291], and Vey [277, 276] imply:

THEOREM. *Let M be a closed affine manifold. Suppose that M is completely incomplete. Then M is a quotient of a properly convex cone.*

I find it very striking that the two extreme cases for closed affine manifolds, the developing map is an embedding. That is, the developing map for a *complete* affine manifold is a diffeomorphism, whereas the developing map for a *completely incomplete* affine manifold embeds the universal covering as a sharp convex cone.

In particular a completely incomplete affine manifold M is radiant. Furthermore it fibers over S^1 as a radiant suspension of an automorphism of a projective manifold of codimension one. Topological consequences are that the Euler characteristic $\chi(M) = 0$ and the first Betti number $b_1(M) > 0$.

Along the way we will also show that complete incompleteness is equivalent to the nonexistence of nonconstant *projective* maps $\mathbb{R} \rightarrow M$.

This striking result uses *intrinsic metrics* for affine and projective manifolds, developed by Vey [275, 277, 274, 276] and Kobayashi [179, 177]. Their constructions were in turn inspired by the intrinsic metrics of Carathéodory and Kobayashi for holomorphic mappings between complex manifolds.

We begin by discussing Kobayashi's pseudometric for domains in projective space, and then extend this construction to projective manifolds.

12.1. The Kobayashi metric

To motivate Kobayashi's construction, consider the basic case of intervals in \mathbb{P}^1 . (Compare the discussion in Exercise 2.5.7 on the cross-ratio.)

There are several natural choices to take, for example, the interval of positive real numbers $\mathbb{R}^+ = (0, \infty)$ or the open unit interval $\mathbf{I} = (-1, 1)$. They relate via the projective transformation $\mathbf{I} \xrightarrow{\tau} \mathbb{R}^+$

$$x = \tau(u) = \frac{1+u}{1-u}$$

mapping $-1 < u < 1$ to $0 < x < \infty$ with $\tau(0) = 1$. The corresponding Hilbert metrics are given by

$$(64) \quad d_{\mathbb{R}^+}(x_1, x_2) = \log \left| \frac{x_1}{x_2} \right|$$

$$(65) \quad d_{\mathbf{I}}(u_1, u_2) = 2 \left| \tanh^{-1}(u_1) - \tanh^{-1}(u_2) \right|.$$

This follows from the fact that τ pulls back the parametrization corresponding to Haar measure on \mathbb{R}^+ :

$$\frac{|dx|}{x} = |d \log x|$$

to the *Poincaré metric* on \mathbf{I} :

$$ds_{\mathbf{I}} = \frac{2 |du|}{1-u^2} = 2 |d \tanh^{-1} u|.$$

A slight generalization of this will be useful in §12.2 in the proof of the projective Brody Lemma 12.2.18:

EXERCISE 12.1.1. *Let $r > 0$ and denote by $\mathbf{I}(r)$ the open interval $(-r, r) \subset \mathbb{R}$. Show that the diffeomorphism*

$$\begin{aligned} \mathbf{I} &\longrightarrow \mathbf{I}(r) \\ u &\longmapsto v = ru \end{aligned}$$

maps $0 \mapsto 0$, takes $(\frac{d}{du})_0$ to $r(\frac{d}{dv})_0$, and, dually,

$$r ds_{\mathbf{I}} = \frac{2r |du|}{1-u^2} \longleftarrow \frac{2r^2 |dv|}{r^2-v^2}.$$

Show that the infinitesimal form of the Hilbert metric on $\mathbf{I}(r)$ is:

$$ds_{\mathbf{I}(r)} = \frac{2r |dv|}{r^2-v^2} = 2 d \tanh^{-1}(v/r)$$

for $v \in \mathbf{I}(r)$ and $v = r \tanh(s/2)$.

EXERCISE 12.1.2. Let $x_-, x_+ \in \mathbb{R} \setminus \{0\}$ be distinct. Show that the projective map mapping

$$\begin{aligned} -1 &\longmapsto x_- \\ 0 &\longmapsto 0 \\ 1 &\longmapsto x_+ \end{aligned}$$

is given by:

$$t \longmapsto \frac{2(x_-x_+)t}{(t+1)x_- + (t-1)x_+},$$

and a projective automorphism of \mathbf{I} by

$$t \longmapsto \frac{\cosh(s)t + \sinh(s)}{\sinh(s)t + \cosh(s)} = \frac{t + \tanh(s)}{1 + \tanh(s)t}$$

for $s \in \mathbb{R}$.

In terms of the Poincaré metric on \mathbf{I} the Hilbert distance $\mathbf{d}(x, y)$ can be characterized as an infimum over all projective maps $\mathbf{I} \rightarrow \Omega$:

$$\mathbf{d}(x, y) = \inf \left\{ \mathbf{d}_{\mathbf{I}}(a, b) \mid f \in \text{Proj}(\mathbf{I}, \Omega), a, b \in \mathbf{I}, f(a) = x, f(b) = y \right\}$$

We now define the Kobayashi pseudometric for any domain Ω and, more generally, any manifold with a projective structure (§12.2). This proceeds by a general universal construction forcing two properties:

- The *triangle inequality*: $\mathbf{d}(a, c) \leq \mathbf{d}(a, b) + \mathbf{d}(b, c)$;
- The *projective Schwarz lemma*: Projective maps do not increase distance.

However, the resulting pseudometric may not be positive; indeed for many domains it is identically zero.

Let $\Omega \subset \mathbf{P}$ be a domain and $x, y \in \Omega$. A (projective) *chain* from x to y is a sequence C of projective maps $f_1, \dots, f_m \in \text{Proj}(\mathbf{I}, \Omega)$ and pairs $a_i, b_i \in \mathbf{I}$, for $i = 1, \dots, m$ such that:

$$\begin{aligned} f_1(a_1) = x, f_1(b_1) = f_2(a_2), \dots, \\ f_{m-1}(b_{m-1}) = f_m(a_m), f_m(b_m) = y. \end{aligned}$$

Denote the set of all projective chains from x to y by $\text{Chain}(x \rightsquigarrow y)$. Define *length* of a projective chain by:

$$\ell(C) = \sum_{i=1}^m \mathbf{d}_{\mathbf{I}}(a_i, b_i).$$

and the *Kobayashi pseudodistance* $\mathbf{d}^{\text{Kob}}(x, y)$:

$$\mathbf{d}^{\text{Kob}}(x, y) = \inf \left\{ \ell(C) \mid C \in \text{Chain}(x \rightsquigarrow y) \right\}.$$

The resulting function enjoys the following obvious properties:

- $d^{\text{Kob}}(x, y) \geq 0$;
- $d^{\text{Kob}}(x, x) = 0$;
- $d^{\text{Kob}}(x, y) = d^{\text{Kob}}(y, x)$;
- (Triangle inequality) $d^{\text{Kob}}(x, y) \leq d^{\text{Kob}}(y, z) + d^{\text{Kob}}(z, x)$.
- (Projective Schwarz lemma) If Ω, Ω' are two domains in projective spaces with Kobayashi pseudometrics d, d' respectively and

$$\Omega \xrightarrow{f} \Omega'$$

is a projective map, then

$$d'(f(x), f(y)) \leq d(x, y).$$

- The Kobayashi pseudometric on the interval \mathbf{I} equals the Hilbert metric on \mathbf{I} .
- d^{Kob} is invariant under the group $\text{Aut}(\Omega)$ consisting of all collineations of \mathbf{P} preserving Ω .

PROPOSITION 12.1.3 (Kobayashi [179]). *Let $\Omega \subset \mathbf{P}$ be properly convex. If $x, y \in \Omega$, then*

$$d^{\text{Hilb}}(x, y) = d^{\text{Kob}}(x, y)$$

COROLLARY 12.1.4. *The function $d^{\text{Hilb}} : \Omega \times \Omega \rightarrow \mathbb{R}$ is a complete metric on Ω .*

PROOF OF PROPOSITION 12.1.3. Let $x, y \in \Omega$ be distinct points and let $l = \overleftrightarrow{xy}$ be the line incident to them. Now

$$d_{\Omega}^{\text{Hilb}}(x, y) = d_{l \cap \Omega}^{\text{Hilb}}(x, y) = d_{l \cap \Omega}^{\text{Kob}}(x, y) \leq d_{\Omega}^{\text{Kob}}(x, y)$$

by the Schwarz lemma applied to the projective map $l \cap \Omega \hookrightarrow \Omega$. For the opposite inequality, let S be the intersection of a supporting hyperplane to Ω at x_{∞} and a supporting hyperplane to Ω at y_{∞} . Projection from S to l defines a projective map

$$\Pi_{S,l} \Omega \rightarrow l \cap \Omega$$

which retracts Ω onto $l \cap \Omega$. Thus

$$d_{\Omega}^{\text{Kob}}(x, y) \leq d_{l \cap \Omega}^{\text{Kob}}(x, y) = d_{\Omega}^{\text{Hilb}}(x, y)$$

(again using the Schwarz lemma) as desired. \square

COROLLARY 12.1.5. *Line segments in Ω are geodesics. If $\Omega \subset \mathbf{P}$ is properly convex, $x, y \in \Omega$, then the chain consisting of a single projective isomorphism*

$$\mathbf{I} \rightarrow \overleftrightarrow{xy} \cap \Omega$$

minimizes the length among all chains in $\text{Chain}(x \rightsquigarrow y)$.

EXERCISE 12.1.6. *Prove that the geodesics in Ω with respect to this metric are straight lines.*

An affine (respectively projective) manifold is *hyperbolic* if the Kobayashi pseudometric \mathbf{d}^{Kob} is a metric, that is, if $\mathbf{d}^{\text{Kob}}(x, y) > 0$ for $x \neq y$. A compact affine manifold M is hyperbolic if and only if it is a quotient of a properly convex cone; a compact projective manifold is hyperbolic if and only if it is a quotient of a properly convex domain in projective space. The tameness of developing maps of hyperbolic affine and projective structures suggests, when the pseudometric \mathbf{d}^{Kob} fails to be a metric, that \mathbf{d}^{Kob} may provide a useful tool to understand pathological developing maps.

12.2. Kobayashi hyperbolicity

Now we discuss intrinsic metrics on affine and projective manifolds. The case of domains was discussed in §??.

Recall from §12.1 the open unit interval $\mathbf{I} = (-1, 1)$ with Poincaré metric

$$\mathbf{g}_{\mathbf{I}} := \frac{4 du^2}{(1 - u^2)^2} = \left(ds_{\mathbf{I}}\right)^2,$$

where

$$ds_{\mathbf{I}} = \sqrt{\mathbf{g}_{\mathbf{I}}} := \frac{2 du}{1 - u^2} = d(2 \tanh^{-1}(u))$$

defines the associated norm on the tangent spaces. As in §12.1, the natural parameter s for arc length on \mathbf{I} relates to the Euclidean coordinate u on $\mathbf{I} \subset \mathbb{R}$ by:

$$u = \tanh(s/2).$$

For projective manifolds M , one defines a “universal” pseudometric

$$M \times M \xrightarrow{\mathbf{d}_M^{\text{Kob}}} \mathbb{R}$$

such that affine (respectively projective) maps $\mathbf{I} \rightarrow M$ are distance non-increasing with respect to $ds_{\mathbf{I}}$. This generalizes the Kobayashi metric for projective domains discussed in §12.1.

The definition of $\mathbf{d}_M^{\text{Kob}}$ for an arbitrary $\mathbb{R}\mathbf{P}^n$ -manifold M enforces the triangle inequality and Schwarz lemma by taking the infimum of $g_{\mathbf{I}}$ -distances over *chains* in M , as in §12.1. Recall that if $x, y \in M$, a (projective) *chain* from x to y is a sequence of projective maps $f_1, \dots, f_m \in \text{Proj}(\mathbf{I}, M)$ and pairs $a_i, b_i \in \mathbf{I}$, for $i = 1, \dots, m$ such that:

$$\begin{aligned} f_1(a_1) = x, f_1(b_1) = f_2(a_2), \dots, \\ f_{m-1}(b_{m-1}) = f_m(a_m), f_m(b_m) = y. \end{aligned}$$

Denote the set of all chains from x to y by $\text{Chain}(x \rightsquigarrow y)$ and define *length*:

$$\begin{aligned} \text{Chain}(x \rightsquigarrow y) & \xrightarrow{\ell} \mathbb{R}_{\geq 0} \\ \left(((a_1, b_1), f_1), \dots, ((a_m, b_m), f_m) \right) & \mapsto \sum_{i=1}^m d_{\mathbf{I}}(a_i, b_i), \end{aligned}$$

where $d_{\mathbf{I}}$ is the distance function on the Riemannian 1-manifold $(\mathbf{I}, \mathbf{g}_{\mathbf{I}})$. Now define the *Kobayashi pseudodistance* $d^{\text{Kob}}(x, y)$ as:

$$\begin{aligned} M \times M & \xrightarrow{d^{\text{Kob}}} \mathbb{R}_{\geq 0} \\ (x, y) & \mapsto \inf \left\{ \ell(C) \mid C \in \text{Chain}(x \rightsquigarrow y) \right\}. \end{aligned}$$

just as in as in §12.1. Just as in the case of domains, d^{Kob} satisfies the triangle inequality and the Projective Schwarz lemma:

LEMMA 12.2.1 (Projective Schwarz Lemma). *Projective maps do not increase pseudodistances: If $x, y \in N$, and $f \in \text{Proj}(N, M)$, then*

$$d_M^{\text{Kob}}(f(x), f(y)) \leq d_N^{\text{Kob}}(x, y).$$

DEFINITION 12.2.2. *Let M be an $\mathbb{R}\mathbb{P}^n$ -manifold. Then M is (projectively) hyperbolic if and only if $d_M^{\text{Kob}} > 0$, that is, if (M, d_M^{Kob}) is a metric space. Say that M is complete hyperbolic if if the metric space (M, d_M^{Kob}) is complete.*

12.2.1. Complete hyperbolicity and convexity. The following convexity theorem is due, independently, to Vey [274, 275], and Kobayashi [179], from somewhat different viewpoints. We closely follow Kobayashi [179]; see also [177, 178].

PROPOSITION 12.2.3. *Let M be a complete hyperbolic projective manifold. Then M is properly convex, that is, M is isomorphic to a quotient of a properly convex domain by a discrete group of collineations.*

The proof will be based on the following fundamental compactness property of projective maps (compare Vey [275], Proposition IV, Chapitre II):

LEMMA 12.2.4. *Suppose that M, N are projective manifolds, where M is complete hyperbolic. Let $p \in N$ and $K \subset\subset M$. Then*

$$\text{Proj}_{p,K}(N, M) := \{ f \in \text{Proj}(N, M) \mid f(p) \in K \}$$

is compact.

PROOF. Apply the Projective Schwarz Lemma 12.2.1 to Lemma C.2.1 (Theorem 3.1 of Chapter V of Kobayashi [180]), discussed in Appendix C. \square

PROOF OF PROPOSITION 12.2.3. We show that M is *geodesically convex*, that is, if $\forall p, q \in M$, every path $p \rightsquigarrow q$ is relatively homotopic to a geodesic path from p to q . We may assume that M is simply connected.

For $p \in M$, let $M(p) \subset M$ be the union of geodesic segments in M beginning at p . Exercise 8.3.4 implies $M(p)$ is open. Since M is connected, it suffices to show $M(p)$ is closed.

Suppose that $q_n \in M(p)$ for $n = 1, 2, \dots$ be a convergent sequence in M with $q = \lim_{n \rightarrow \infty} q_n$. We show that $q \in M(p)$.

Let $0 < a < 1$ and $f_n \in \text{Proj}(\mathbf{I}, M)$ with $f_n(0) = p$ and $f_n(a) = q_n$. Lemma 12.2.4 guarantees a subsequence of f_n converging to a projective map $f \in \text{Proj}(\mathbf{I}, M)$ with $f(0) = p$. Then

$$q = \lim_{n \rightarrow \infty} q_n = \lim_{n \rightarrow \infty} f_n(a) = f(a) \in M(p)$$

and $M(p)$ is closed, as desired. \square

12.2.2. The infinitesimal form. Kobayashi's pseudometric d^{Kob} has an infinitesimal form Φ^{Kob} defined by a function $TM \xrightarrow{\Phi^{\text{Kob}}} \mathbb{R}$. That is, $d^{\text{Kob}}(p, q)$ is the infimum of the *pseudolengths*

$$\ell(\gamma) := \int_{\gamma} \Phi^{\text{Kob}}(\gamma')$$

over piecewise C^1 paths $p \rightsquigarrow q$. For $x \in M$ and $\xi \in T_x M$, define:

$$(66) \quad \Phi^{\text{Kob}}(\xi) := \inf \left\{ |ds_{\mathbf{I}}(\mathbf{v})| \mid f \in \text{Proj}(\mathbf{I}, M), f(u) = x, (Df)_u(\mathbf{v}) = \xi \right\}$$

where $u \in \mathbf{I}$ and $ds_{\mathbf{I}}(\mathbf{v})$ denotes the norm of $\mathbf{v} \in T_u \mathbf{I}$ with respect to the Poincaré metric $(ds_{\mathbf{I}})^2$ on \mathbf{I} .

EXERCISE 12.2.5. For affine manifolds, completeness is equivalent to $\Phi^{\text{Kob}} \equiv 0$. For a Hopf manifold, $d^{\text{Kob}} \equiv 0$ but $\Phi^{\text{Kob}} \neq 0$. Indeed $\Phi^{\text{Kob}}(\mathbf{R}) = 1$ where \mathbf{R} is the radiant vector field.

EXERCISE 12.2.6. Show that Φ^{Kob} is homogeneous of degree one, that is,

$$\Phi^{\text{Kob}}(r\xi) = r \Phi^{\text{Kob}}(\xi)$$

for $r \geq 0$. Deduce that $\ell(\gamma)$ is independent of the parametrization of γ .

PROPOSITION 12.2.7. Φ^{Kob} is upper semicontinuous.

Recall that a function $X \xrightarrow{f} \mathbb{R}$ is upper semicontinuous at $x \in X$ if and only if $\forall \epsilon > 0$,

$$f(y) < f(x) + \epsilon$$

for y in an open neighborhood of x . That is, the values of f cannot “jump down” in limits.

$$\lim_{n \rightarrow \infty} f(\xi_n) \leq f\left(\lim_{n \rightarrow \infty} \xi_n\right)$$

for convergent sequences ξ_n . Equivalently, f is a continuous mapping from X to \mathbb{R} , where \mathbb{R} is given the topology whose open sets are intervals $(-\infty, a)$ where $a \in \mathbb{R}$. The indicator function of a closed set is upper semicontinuous. Semicontinuous functions are further discussed in Appendix D.

PROOF OF PROPOSITION 12.2.7. Let $x \in M$ and $\xi \in \mathbb{T}_x M$ and write

$$\Phi^{\text{Kob}}(\xi) = k.$$

Let $\epsilon > 0$. Then $\exists f \in \text{Proj}(\mathbf{I}, M)$ with $f(u) = x$, and $\mathbf{v} \in \mathbb{T}_u \mathbf{I}$ with $(Df)_u(\mathbf{v}) = \xi$ and

$$\|\mathbf{v}\| < k + \epsilon/2.$$

Lift f to $\tilde{f} \in \text{Proj}(\mathbf{I}, \tilde{M})$ and extend \tilde{f} to $\tilde{F} \in \text{Proj}(\mathbb{B}, \tilde{M})$. Let $\|\cdot\|_{\mathbb{B}}$ the corresponding norm for the intrinsic metric on \mathbb{B} defined in §3.3. We may assume that $\|\xi\|_{\mathbb{B}} = \|\xi\|$. Then an open neighborhood \mathcal{N} of $\xi \in \mathbb{T}\tilde{M}$ exists so that if $\xi' \in \mathcal{N}$, then:

- $\xi' \in \mathbb{T}_{x'} \tilde{M}$ where x' lies in the image $\tilde{F}(\mathbb{B}) \subset \tilde{M}$;
- $\|(D\tilde{F})^{-1}(\xi')\|_{\mathbb{B}} < k + \epsilon$.

Since projective maps do not increase distance, $\|\xi'\|_{\tilde{M}} < k + \epsilon$, as desired. \square

COROLLARY 12.2.8 (Proposition 5.16 of Kobayashi [179]). *If M is a complete hyperbolic projective manifold, then $\mathbb{T}M \xrightarrow{\Phi^{\text{Kob}}} \mathbb{R}$ is continuous.*

The proof uses the following lemma (Lemma 5.17 of Kobayashi [179]), stating that the infimum in the definition of Φ^{Kob} is actually achieved.

LEMMA 12.2.9. *Suppose M is complete hyperbolic and $\xi \in \mathbb{T}_p M$. Then $\exists f \in \text{Proj}(\mathbf{I}, M)$ and $\mathbf{v} \in \mathbb{T}_0 \mathbf{I}$ with $f(0) = p$ and $(Df)_0(\mathbf{v}) = \xi$ such that $\Phi^{\text{Kob}}(\xi) = \|\mathbf{v}\|$.*

PROOF. Lemma 12.2.4 implies that the subset of $\text{Proj}(\mathbf{I}, M)$ comprising projective maps $\mathbf{I} \xrightarrow{f} M$ with $f(0) = p$ is compact. Thus the set of $\|(\mathbf{D}f)^{-1}(\xi)\|$ is a compact subset of \mathbb{R}^+ , so its infimum $\Phi^{\text{Kob}}(\xi)$ is positive. \square

PROOF OF COROLLARY 12.2.8. Suppose, for $k = 1, 2, \dots$, that $\xi_k \in \mathbb{T}_{p_k}M$, defines a sequence converging to $\xi_\infty \in \mathbb{T}_{p_\infty}M$. We show that $\lim_{k \rightarrow \infty} \Phi^{\text{Kob}}(\xi_k) = \Phi^{\text{Kob}}(\xi_\infty)$.

Proposition 12.2.7 (semicontinuity of Φ^{Kob}) implies that

$$(67) \quad \Phi^{\text{Kob}}(\xi_\infty) \geq \lim_{k \rightarrow \infty} \Phi^{\text{Kob}}(\xi_k),$$

so it suffices to show that $\Phi^{\text{Kob}}(\xi_\infty) \leq \lim_{k \rightarrow \infty} \Phi^{\text{Kob}}(\xi_k)$.

The above lemma guarantees $f_k \in \text{Proj}(\mathbf{I}, M)$ with $f_k(0) = p_k$ and

$$\mathbf{v}_k := (\mathbf{D}f_k)^{-1}(\xi_k) \in \mathbb{T}_0\mathbf{I}$$

such that $\Phi^{\text{Kob}}(\xi_k) = \|\mathbf{v}_k\|$. By (67), \mathbf{v}_k contains a convergent subsequence, and let $\mathbf{v}_\infty := \lim_{k \rightarrow \infty} \mathbf{v}_k$. Lemma 12.2.4 guarantees that by passing to a further subsequence, we may assume that f_k converges to $f_\infty \in \text{Proj}(\mathbf{I}, M)$ with $f_\infty(0) = p_\infty$ and $\mathbf{D}f_\infty(\mathbf{v}_\infty) = \xi_\infty$. By the definition of Φ^{Kob} ,

$$\Phi^{\text{Kob}}(\xi_\infty) \leq \|\mathbf{v}_\infty\| = \lim_{k \rightarrow \infty} \|\mathbf{v}_k\| = \lim_{k \rightarrow \infty} \Phi^{\text{Kob}}(\xi_k)$$

as desired. \square

EXERCISE 12.2.10. Find an example of a projectively hyperbolic domain for which Φ^{Kob} is not continuous.

EXERCISE 12.2.11. Find an example of a domain Ω for which:

- $\Phi^{\text{Kob}}(\xi) < \infty$ for all nonzero $\xi \in \mathbb{T}\Omega$;
- Ω contains no complete geodesic rays.

THEOREM 12.2.12. Φ^{Kob} is the infinitesimal form of the Kobayashi pseudometric $\mathbf{d}_M^{\text{Kob}}$, that is,

$$\mathbf{d}_M^{\text{Kob}}(x, y) = \inf \left\{ \int_a^b \Phi^{\text{Kob}}(\gamma'(t)) dt \mid \gamma \in \text{Path}(x \rightsquigarrow y) \right\}$$

where $\text{Path}(x \rightsquigarrow y)$ denotes the set of piecewise C^1 paths $[a, b] \xrightarrow{\gamma} M$ with $\gamma(a) = x$, $\gamma(b) = y$.

Since semicontinuous functions are bounded on compact sets (Exercise D.2.3) and measurable (Exercise D.1.2), the above integral is well-defined.

PROOF OF THEOREM 12.2.12. Define

$$\delta^{\text{Kob}}(p, q) := \inf \left\{ \int_a^b \Phi^{\text{Kob}}(\gamma'(t)) dt \mid \gamma \in \text{Path}(p \rightsquigarrow q) \right\}$$

We must prove that $\delta^{\text{Kob}} = \mathbf{d}^{\text{Kob}}$.

To prove $\delta^{\text{Kob}} \leq \mathbf{d}^{\text{Kob}}$, note that every chain $C \in \text{Chain}(x \rightsquigarrow y)$ determine a piecewise C^1 path $\gamma_C \in \text{Path}(p \rightsquigarrow q)$. Since $\mathbf{g}_{\mathbf{I}}$ is the infinitesimal form of $\mathbf{d}_{\mathbf{I}}$, the path γ_C has shorter length than the chain C , that is, their lengths satisfy $\ell(\gamma_C) \leq \ell(C)$. Taking infima implies $\delta^{\text{Kob}} \leq \mathbf{d}^{\text{Kob}}$ as desired.

We prove $\delta^{\text{Kob}} \geq \mathbf{d}^{\text{Kob}}$. Suppose that $\gamma \in \text{Path}(p \rightsquigarrow q)$ as above. Suppose $\epsilon > 0$. We seek a chain $C \in \text{Chain}(p \rightsquigarrow q)$ such that

$$(68) \quad \ell(C) \leq \ell(\gamma) + \epsilon.$$

Proposition 12.2.7 implies that the function

$$\begin{aligned} [a, b] &\xrightarrow{\phi} \mathbb{R} \\ u &\longmapsto \Phi^{\text{Kob}}(\gamma'(t)) \end{aligned}$$

is upper semicontinuous. Exercise D.2.3 implies ϕ is bounded from above. Apply Proposition D.2.4 to conclude that ϕ is the limit of a monotonically decreasing sequence of nonnegative continuous functions.

Apply Lebesgue's monotone convergence theorem (Rudin [243], 1.26) to find a continuous function $[a, b] \xrightarrow{h} \mathbb{R}$ such that:

$$(69) \quad \phi(t) < h(t) \text{ for } a \leq t \leq b$$

$$(70) \quad \int_{[a, b]} h < \ell(\gamma) + \epsilon.$$

for $a \leq t \leq b$.

We claim that for each $s \in [a, b]$,

$$(71) \quad \int_s^t \phi(u) du \leq (1 + \epsilon) h(s) |s - t|$$

for t in an interval I_s centered at s .

To this end, first assume that γ is C^1 ; We choose the open neighborhood I_s of s in three steps:

First, choose a convex ball W_s containing $\gamma(s)$, so that $\gamma(t) \in W_s$ for $t \in I_s$. Let $f_t \in \text{Proj}(\mathbf{I}, M)$ extend the geodesic in W_s joining $\gamma(s)$ to $\gamma(t)$. Then (69) implies that $\|(\mathbf{D}f_t)^{-1}(\gamma'(s))\| < h$. Since γ is C^1 ,

$$(72) \quad \|(\mathbf{D}f_t)^{-1}(\gamma'(t))\| < h$$

for t sufficiently near s .

Next, choose I_s so that (72) holds for $t \in I_s$. This implies that

$$(73) \quad \int_s^t h(u)du \leq (1 + \epsilon) h(s) |s - t|$$

for t sufficiently near s .

Finally, choose I_s so that (73) holds for $t \in I_s$.

Combining (73) with (69) implies

$$\int_s^t \phi(u)du < \int_s^t h(u)du \leq (1 + \epsilon) h(s) |s - t|,$$

establishing the claim when γ is C^1 . Extending (73) to the case that γ is only *piecewise* C^1 is a routine exercise.

Continuing to follow Wu [291]), we pick up the argument of Royden [242]. Let $\eta > 0$ be a *Lebesgue number* for the open cover $\{I_s \mid s \in \mathbf{I}\}$ of \mathbf{I} , that is, every closed interval of length $< \eta$ lies in some I_s . (For the reader's convenience, a proof of the existence of the Lebesgue number is given in Appendix C.3.) Thus there exists a subdivision $a = t_0 < t_1 < \dots < t_k = b$ exists with $t_i - t_{i-1} < \eta$; let s_i be such that $[t_{i-1}, t_i] \subset I_{s_i}$. Continuity of h and (70) imply

$$(74) \quad \sum_{i=1}^k h(s_i)(t_i - t_{i-1}) < \int_a^b h(u) < \ell(\gamma) + \epsilon.$$

By (71),

$$\begin{aligned} d^{\text{Kob}}(\gamma(s_i), \gamma(s_{i-1})) &\leq d^{\text{Kob}}(\gamma(s_i), \gamma(s_{i-1})) + d^{\text{Kob}}(\gamma(s_i), \gamma(s_{i-1})) \\ &\leq (1 + \epsilon) (h(s_i)(s_i - t_i) + h(s_{i-1})(t_i - s_{i-1})) \end{aligned}$$

Apply (73) and (74), obtaining:

$$\begin{aligned} d^{\text{Kob}}(p, q) &\leq \sum_{i=1}^k d^{\text{Kob}}(\gamma(s_i), \gamma(s_{i-1})) \\ &\leq \sum_{i=1}^k h(s_i)(t_i - t_{i-1}) \\ &< \ell(\gamma) + \epsilon. \end{aligned}$$

Now the chain $C \in \text{Chain}(p \rightsquigarrow q)$ defined by:

$$C := \left((s_1, s_2, f_1), \dots, (s_{k-1}, s_k, f_k) \right)$$

has length

$$\ell(C) = \sum_{i=1}^{k-1} d^{\text{Kob}}(\gamma(s_i, s_{i+1}))$$

and (68) follows. Since $\epsilon > 0$ is arbitrary, $d^{\text{Kob}}(p, q) \leq \ell(\gamma) \leq \delta$ as desired. \square

Wu [291]) actually proves a much stronger statement, valid for affine connections which are *not necessarily* flat. His proof is based on the analog for the Kobayashi pseudometric for complex manifolds, due to Royden [242].

Closely related is the universal property of Φ^{Kob} among infinitesimal pseudometrics for which projective maps are infinitesimally nonincreasing (Kobayashi [179], Proposition 5.5):

EXERCISE 12.2.13. *Let M be an $\mathbb{R}\mathbb{P}^n$ -manifold and $M \xrightarrow{\Phi} \mathbb{R}_{\geq 0}$ a function such that $\forall f \in \text{Proj}(\mathbf{I}, M)$*

$$\Phi((Df)_a(\xi)) \leq \|\xi\|_a,$$

where $-1 < a < 1$ and $\xi \in T_a\mathbf{I}$. Then $\Phi \leq \Phi^{\text{Kob}}$.

12.2.3. Completely incomplete manifolds.

EXERCISE 12.2.14. *Let M be a affine manifold. Suppose that every geodesic ray is incomplete. Then M is noncompact.*

THEOREM 12.2.15. *Suppose that M is an $\mathbb{R}\mathbb{P}^n$ -manifold. Then $d_M^{\text{Kob}} > 0$ if and only if every projective map $\mathbb{R} \rightarrow M$ is constant.*

More specifically, we prove (following Kobayashi [179]):

PROPOSITION 12.2.16. *Let M be a projective manifold, $p \in M$ and $\xi \in T_pM$. Suppose $\Phi^{\text{Kob}}(\xi) = 0$. Then $\exists f \in \text{Proj}(\mathbb{R}, M)$ with $f(0) = p$ and $f'(0) = \xi$.*

Since $\Phi^{\text{Kob}}(p, \xi) = 0$, there is a sequence $j_m \in \text{Proj}(\mathbf{I}, M)$ with $j_m(0) = p$ and a sequence $a_m > 0$ with a_m decreasing, $a_m \searrow 0$ such that the differential $(Dj_m)_0$ of j_m at $0 \in \mathbf{I}$ maps:

$$\begin{array}{ccc} T_0\mathbf{I} & \xrightarrow{(Dj_m)_0} & T_pM \\ a_m \left(\frac{d}{du} \right)_0 & \mapsto & \xi. \end{array}$$

Let $r_m = 1/a_m$ so that

$$(75) \quad \left(\frac{d}{du} \right)_0 \xrightarrow{(Dj_m)_0} r_m \xi$$

with $r_m \nearrow +\infty$ monotonically.

As in Exercise 12.1.1, let $\mathbf{I}(r)$ denotes the open interval $(-r, r) \subset \mathbb{R}$. Then

$$\mathbf{I}(r_1) \subset \mathbf{I}(r_2) \subset \cdots \subset \mathbf{I}(r_m) \subset \cdots \subset \mathbb{R}$$

and $\bigcup_{m=1}^{\infty} \mathbf{I}(r_m) = \mathbb{R}$.

The strategy of the proof is to reparametrize the maps j_m to obtain a subsequence of projective maps

$$h_m \in \text{Proj}(\mathbf{I}(r_m), M),$$

such that the restriction of h_m to h_l equals h_l for $l \leq m$.

First renormalize j_m to a projective map $f_m \in \text{Proj}(\mathbf{I}(r_m), M)$:

$$\begin{aligned} \mathbf{I}(r_m) &\xrightarrow{f_m} M \\ u &\longmapsto j_m(u/r_m) \end{aligned}$$

LEMMA 12.2.17. *The differential $(Df_m)_0$ of f_m at 0 maps the tangent vector $(\frac{d}{du})_0 \in \mathbb{T}_0(\mathbf{I}(r_m))$ to ξ .*

PROOF. f_m equals the composition of $\mathbf{I} \xrightarrow{f} M$ with the contraction

$$\begin{aligned} \mathbf{I}(r_m) &\longrightarrow \mathbf{I} \\ u &\longmapsto u/r_m \\ 0 &\longmapsto 0 \end{aligned}$$

whose differential at 0 is multiplication by $(r_m)^{-1}$. Now apply the chain rule and (75). □

Next reparametrize the maps f_m using the following analog of Brody’s reparametrization lemma for holomorphic mappings (Brody [47]), whose proof is given later.

As in Exercise 12.1.1, the infinitesimal norm for $\mathbf{I}(r)$ equals:

$$ds_{\mathbf{I}(r)} = \frac{2r|du|}{r^2 - u^2},$$

a function on $\mathbf{I}(r)$. As in Kobayashi [179]¹ choose a Riemannian metric \mathbf{g} on M such that $\mathbf{g}(\xi) = 1$. The corresponding norm on $\mathbb{T}M$ is $\sqrt{\mathbf{g}}$ and pulls back to a norm $f^*\sqrt{\mathbf{g}}$ on $\mathbb{T}_u\mathbf{I}(r)$. Then there is a continuous function

$$\mathbf{I}(r) \xrightarrow{\mathcal{W}_f} \mathbb{R}^+$$

such that

$$(76) \quad f^*\sqrt{\mathbf{g}}(u) = \mathcal{W}_f(u) ds_{\mathbf{I}(r)}(u).$$

LEMMA 12.2.18 (Reparametrization Lemma). *Let M be an projective manifold, $p \in M$ and $\xi \in \mathbb{T}_pM$ nonzero. Suppose that $f \in \text{Proj}(\mathbf{I}(r), M)$ with $f(0) = p$ and $f'(0) = \xi$. Choose c such that $\mathcal{W}_f(0) >$*

¹Remark 5.27 after Theorem 5.22, pp.145–146

$c > 0$. Then $\exists a, b$ with $0 < a < 1$ and $b \in \text{Aut}(\mathbf{I}(r))$ such that $h \in \text{Proj}(\mathbf{I}_r, M)$ defined by:

$$h(u) := f(ab(u))$$

satisfies

- $\mathcal{W}_h(u) \leq c$;
- $\mathcal{W}_h(0) = c$,

where \mathcal{W}_h is defined in (76)

CONCLUSION OF PROOF OF PROPOSITION 12.2.16 ASSUMING LEMMA 12.2.18. Applying Lemma 12.2.18 to f_m , $\exists h_m \in \text{Proj}(\mathbf{I}(r_m), M)$ and $c_m > 0$ such that

$$(77) \quad \mathcal{W}_{h_m}(u) \leq c_m \text{ and } \mathcal{W}_{h_m}(0) = c$$

and the image of f_m contains the image of h_m .

Denote the restriction of h_m to \mathbf{I}_{r_l} by $h_{l,m}$. Equation (77) implies that, for each $l \in \mathbb{N}$, the family

$$\mathcal{F}_l := \{h_{l,m} \mid m \geq l\}$$

is equicontinuous.

We construct the projective map $h \in \text{Proj}(\mathbb{R}, M)$ by consecutive extensions $h_l \in \text{Proj}(\mathbf{I}_{r_l}, M)$ to $\mathbf{I}_{r_l} \supset \mathbf{I}_{r_{l-1}}$ as follows.

Beginning with $l = 1$, the Arzelà-Ascoli theorem guarantees a convergent subsequence $h_{1,m}$ in $\text{Proj}(\mathbf{I}_{r_1}, M)$. Write

$$h_1 = \lim_{m \rightarrow \infty} h_{1,m} \in \text{Proj}(\mathbf{I}_{r_1}, M).$$

Suppose inductively that $h_l \in \text{Proj}(\mathbf{I}_{r_l}, M)$ has been defined such that h_l extends h_k for all $k \leq l$. Since \mathcal{F}_l is equicontinuous, the Arzelà-Ascoli theorem guarantees a convergent subsequence of $h_{l,m}$. Define

$$h_l := \lim_{m \rightarrow \infty} h_{l,m}.$$

The value of $h_m^* \sqrt{g}$ at $u = 0$ equals $2c \, du \neq 0$. Since this is the value of $h^* \sqrt{g} = h_l^* \sqrt{g}$ at $u = 0$, the map h is nonconstant. This concludes the proof of Proposition 12.2.16 assuming Lemma 12.2.18. \square

PROOF OF LEMMA 12.2.18. For $0 \leq t \leq 1$, consider the projective map

$$\begin{aligned} \mathbf{I}(r) &\xrightarrow{f_t} M \\ u &\longmapsto f(tu). \end{aligned}$$

Then the corresponding function $\mathcal{W}_t := \mathcal{W}_{f_t}$ (defined as in (76))

$$\mathbf{I}(r) \xrightarrow{\mathcal{W}_t} \mathbb{R}^+$$

satisfies the following elementary properties, whose proofs are left as exercises:

$$(78) \quad \mathcal{W}_t(u) = \mathcal{W}_f(tu) \frac{t(r^2 - u^2)}{r^2 - t^2u^2}$$

$$(79) \quad \mathcal{W}_t(u) \geq 0 \text{ and } \mathcal{W}_t(u) = 0 \iff t = 0.$$

$$(80) \quad \lim_{u \rightarrow \pm r} \mathcal{W}_t(u) = 0.$$

The function

$$A(t) := \sup_{u \in \mathbf{I}(r)} \mathcal{W}_t(u)$$

of $t \in [0, 1]$ satisfies the following elementary properties, whose proofs are also left as exercises:

- $A(t) \leq \infty$
- $[0, 1] \xrightarrow{A} \mathbb{R}^+$ is continuous.
- A is monotone-increasing.

Furthermore $A(0) = 0$ and $A(1) = c$, so the Intermediate Value Theorem guarantees $\exists a \in [0, 1]$ such that $A(a) = c$. Thus

$$c = \sup_{u \in \mathbf{I}(r)} \mathcal{W}_a(u).$$

By (80), \mathcal{W}_a assumes its maximum on $u_0 \in \mathbf{I}(r)$; let $b \in \text{Aut}(\mathbf{I}(r))$ take 0 to u_0 . The proof of Lemma 12.2.18 is complete. □

12.3. Hessian manifolds

When M is affine, then Corollary 4.3.2 implies that M is a quotient of a properly convex cone Ω by a discrete group of collineations acting properly on Ω . In addition to the Hilbert metric, Ω enjoys the natural Riemannian metric introduced by Vinberg [278], Koszul [184, 183, 187, 186] and Vesentini [273]. (Compare §??.) In particular Koszul and Vinberg observe that this Riemannian structure is the covariant differential $\nabla\omega$ of a closed 1-form ω . In particular ω is everywhere nonzero, so by Tischler [267], M fibers over S^1 .

This implies Koszul's beautiful theorem [187] that the holonomy mapping hol (described in Chapter 7, §7.2) embeds the space of convex structures onto an open subset of the representation variety. This

has recently been extended to noncompact manifolds by Cooper-Long-Tillmann [79].

Hyperbolic affine manifolds closely relate to *Hessian manifolds*. If ω is a closed 1-form, then its covariant differential $\nabla\omega$ is a symmetric 2-form. Since closed forms are locally exact, $\omega = df$ for some function; in that case $\nabla\omega$ equals the *Hessian* d^2f . Koszul [187] showed that hyperbolicity is equivalent to the existence of a closed 1-form ω whose covariant differential $\nabla\omega$ is positive definite, that is, a Riemannian metric. More generally, Shima [250] considered Riemannian metrics on an affine manifold which are locally Hessians of functions, and proved that such a closed *Hessian* manifold is a quotient of a convex domain, thus generalizing Koszul's result.

We briefly sketch some of the ideas in Koszul's paper. We recommend Shima's book [250] for a very accessible and comprehensive exposition of these and related ideas.

Let (M, ∇) be an affine manifold with connection ∇ . Let $x \in M$ and $\mathcal{E}_x \in \mathbb{T}_x M$ denote the domain of exponential map as in §8.3.1. For $\xi \in \mathbb{T}_x M$, let

$$\lambda(\xi) := \sup\{t \in \mathbb{R} \mid t\xi \in \mathcal{E}_x\} \in (0, \infty],$$

so that the intersection of the line $\mathbb{R}\xi$ with \mathcal{E}_x equals $(-\lambda(-\xi), \lambda(\xi))\xi$. Suppose that ω is a closed 1-form as above, such that the covariant differential $\nabla\omega > 0$. Koszul's theory is based on the two lemmas below. For notational simplicity, write

$$\begin{aligned} (-\lambda(-\xi), \lambda(\xi)) &\xrightarrow{\gamma} M \\ t &\longmapsto \text{Exp}_x(t\xi) \end{aligned}$$

for the maximal geodesic with velocity $\xi = \gamma'(0)$ at time $t = 0$. Observe that for any $-\lambda(-\xi) < t < \lambda(\xi)$, the velocity vector at time t is

$$\gamma'(t) = \mathbb{P}_{\gamma_t}(\xi)$$

where $x \xrightarrow{\gamma_t} \gamma(t)$ is the restriction of γ and

$$\mathbb{T}_x M \xrightarrow{\mathbb{P}_{\gamma_t}} \mathbb{T}_{\gamma(t)} M$$

denotes parallel transport along γ_t . The two basic lemmas are:

LEMMA 12.3.1. *If $\omega(\xi) > 0$, then $\lambda(\xi) < \infty$.*

LEMMA 12.3.2. *If $\lambda(\xi) < \infty$, then*

$$\int_0^{\lambda(\xi)} \omega(\gamma'(t)) dt = +\infty.$$

The role of positivity is apparent from the simple 1-dimensional example when $M = \mathbb{R}^+ \subset \mathbb{R}$. To develop intuition for these conditions, we work out these lemmas in the basic example when $M = \mathbb{R}^+ \subset \mathbb{R}$.

First we show (Lemma 12.3.1) that that $\omega(\xi) > 0$ implies the the geodesic ray γ is incomplete. The cone Ω^* dual to M consists of all $\psi > 0$ and the characteristic function is

$$M \xrightarrow{f} \mathbb{R}$$

$$x \longmapsto \int_{\Omega^*} e^{-\psi x} d\psi = \int_0^\infty e^{-\psi x} d\psi = \frac{1}{x}$$

and the logarithmic differential equals

$$\omega = d \log f = -\frac{dx}{x}.$$

If $\xi = y_0 \partial_x \in T_{x_0} M$ and $\xi \neq 0$, then $\gamma(t) = x_0 + ty_0$ and

$$\lambda(\xi) = \begin{cases} -x_0/y_0 & \text{if } y_0 < 0 \\ \infty & \text{if } y_0 > 0 \end{cases}$$

Similarly

$$\omega(\gamma'(t)) = (-dx/(x_0 + ty_0))(y_0 \partial_x|_{x=x_0+ty_0}) = -y_0/(x_0 + ty_0)$$

so if $\omega(\gamma'(t)) > 0$, then $y_0 < 0$ and $\lambda(\xi) < \infty$ as desired.

Now we verify (Lemma 12.3.2) that along an incomplete geodesic ray, the integral of $\omega(\gamma'(t))$ diverges. Suppose $\lambda(\xi) < \infty$. Then $y_0 < 0$ and

$$\int_0^{\lambda(\xi)} \omega(\gamma'(t)) dt = \log \frac{x_0}{x_0 + ty_0} \Big|_{t=0}^{-x_0/y_0} = \infty$$

as desired.

Projective structures on surfaces

\mathbb{RP}^2 -manifolds are relatively well understood, due to intense activity in recent years. Rather than give an detailed description of this theory, we only summarize the results, and refer to the literature. In particular we recommend the recent book by Casella-Tate-Tillmann [58].

Aside from the two structures with finite fundamental group (\mathbb{RP}^2 itself, and its double cover S^2), this class of geometric structures includes affine structures on surfaces, some new \mathbb{RP}^2 -structures on tori (first analyzed by Sullivan-Thurston [262], Smillie [253] and the author [114] in 1976–1977). as well as convex structures (which are *hyperbolic* in the sense of Kobayashi and Vey; see §12.2). Strikingly the answer is much more satisfactory for surfaces with $\chi < 0$, aside from \mathbb{RP}^2 -structures on tori and Klein bottles which are *not* affine, this chapter concentrates on surfaces of $\chi < 0$. For these, the convex structures play a fundamental role.

13.1. Classification in higher genus

The deformation space $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ of convex \mathbb{RP}^2 structures was calculated by the author [122] in 1985, using the analog of Fenchel-Nielsen coordinates. Shortly thereafter, in his doctoral thesis, Suhyoung Choi proved his *Convex Decomposition Theorem* [66, 68, 68], expressing that on a closed surface of $\chi < 0$, every \mathbb{RP}^2 is obtained from a convex surface by grafting annuli. We summarize their classification as follows (compare Choi-Goldman [70]):

THEOREM. *Let Σ be a closed orientable surface of genus $g > 1$.*

- *The deformation space $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ of marked convex \mathbb{RP}^2 -structures on Σ is homeomorphic to \mathbb{R}^{16g-16} , upon which $\text{Mod}(\Sigma)$ acts properly.*
- *The holonomy map hol embeds $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ as a connected component of $\text{Hom}(\pi, G)/\text{Inn}(G)$ where $G = \text{PGL}(3, \mathbb{R})$.*
- *The deformation space of marked \mathbb{RP}^2 -structures on Σ is homeomorphic to $\mathbb{R}^{16g-16} \times \mathbb{N}$, upon which $\text{Mod}(\Sigma)$ acts properly.*

The first proof that $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ is a cell of dimension $16g - 16$ (in [122]) involves a more general statement, valid when $\partial\Sigma \neq \emptyset$ but with some boundary conditions. The proof introduces an extension of the Fenchel-Nielsen coordinates on the Fricke space — the deformation space of hyperbolic structures on Σ — as described in §4. A particularly tractable and suggestive set of coordinates is due to Fock and Goncharov in [101, 102], and based on parametrizations of hyperbolic structures due to Thurston (see Bonahon [42]) and Penner [234, 235]. Bonahon-Kim [45] describe the relationship between the author’s original extended Fenchel-Nielsen coordinates and Fock-Goncharov’s extended Penner-Thurston coordinates. While the former uses decompositions of Σ into three-holed spheres (*pants decompositions*), the latter uses ideal triangulations.

The symplectic geometry of $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ is described in Choi-Jung-Kim [73] and another approach using affine connections is discussed in Goldman [123].

Choi’s convex decomposition theorem uses a grafting construction, which he extended to nonorientable surfaces [67]. Choi-Goldman [71] extends the calculation to 2-dimensional orbifolds. Unfortunately for lack of space, we do not describe the deep ideas in the convex decomposition theorem.

Somewhat curiously, the diversity of affine structures on the torus gives a much less clean classification for $\chi = 0$ than the geometrically and analytically more interesting case where $\chi < 0$.

13.2. Convex \mathbb{RP}^2 -structures

In general if Ω/Γ is a convex \mathbb{RP}^2 -manifold which is a *closed* surface S with $\chi(S) < 0$, then either $\partial\Omega$ is a conic, or $\partial\Omega$ is a C^1 convex curve (Benzécri [34] which is not C^2 (Kuiper [192])). The key general point is that, if Σ is a closed surface of $\chi < 0$, then the dynamics of the holonomy group $\Gamma \cong \pi_1(\Sigma)$ is standard: If $1 \neq \gamma \in \Gamma < \mathbf{SL}(3, \mathbb{R})$, then γ has one repelling fixed point p_+ on $\partial\Omega$ and one attracting fixed point p_- on $\partial\Omega$. Furthermore, since $\partial\Omega$ is C^1 , the projective lines l_{\pm} tangent to $\partial\Omega$ at p_{\pm} respectively are also γ -invariant. It follows that the intersection $p_0 := l_+ \cap l_-$ is a third fixed point for γ , which has saddle point dynamics. This implies that γ is represented by a 3×3 diagonal matrix with real and distinct eigenvalues. (Compare Fig. 13.1.)

Such examples seem to be conjectured *not* to exist in the original articles of Ehresmann [96] and Benzécri [34]. They are analogous to




FIGURE 13.1. A projective transformation leaving invariant a closed convex curve.

quasi-Fuchsian surface groups in $\mathrm{PSL}(2, \mathbb{C})$, although $\partial\Omega$ is considerably more regular than in the classical case, where the limit circle is not even rectifiable.

In fact the derivative of $\partial\Omega$ is Hölder continuous with Hölder exponent strictly between 1 and 2. The Hölder exponent of the limit circle is a fascinating invariant, which for reasons of space, we do not discuss. We refer to Guichard [140] for some of the first work on this subject. Recently this invariant has been related to the *entropy* of the Hilbert geodesic flow associated to the \mathbb{RP}^2 -structure.

13.2.1. Triangle groups. Figure 3.3 is the first example of a convex \mathbb{RP}^2 -manifold Ω/Γ (actually an orbifold) which is *not* homogeneous (Kac-Vinberg [279]). It arises from a $(3, 3, 4)$ -triangle tessellation, and Γ is the Weyl group of a Kac-Moody Lie algebra of hyperbolic type as follows. Namely the Cartan matrix

$$C = \begin{bmatrix} 2 & -1 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

determines a *Coxeter group*, that is, a group generated by reflections.

For $i = 1, 2, 3$ let E_{ii} denote the elementary matrix having entry 1 in the i -th diagonal slot. Then, for $i = 1, 2, 3$, the reflections

$$\rho_i = I - E_{ii}C$$

generate a discrete subgroup $\Gamma < \mathrm{SL}(3, \mathbb{Z})$ which acts properly on the convex domain depicted in (and appears on the cover of the November 2002 Notices of the American Mathematical Society).

Recall that a reflection ρ in \mathbb{RP}^2 is determined by its fixed set $\mathrm{Fix}(\rho)$, which is a disjoint union $p \sqcup \ell$. Write f_i for the isolated fixed point of ρ_i and ℓ_i for the fixed line of ρ_i . Exercise 2.5.8 imply the relations in the Coxeter group imply conditions on cross ratios of lines passing through f_i .

EXERCISE 13.2.1. Let $p, q, r \in \mathbb{N} \cup \{\infty\}$ and define

$$\begin{aligned} \Gamma(p, q, r) &:= \langle R_1, R_2, R_3 \mid R_1^2 = R_2^2 = R_3^2 = \mathbb{I}, \\ &\quad (R_1 R_2)^p = (R_2 R_3)^q = (R_3 R_1)^r = \mathbb{I} \rangle. \end{aligned}$$

Let $\Delta \subset \mathbb{RP}^2$ with sides s_1, s_2, s_3 respectively, and consider representations $\rho \in \mathrm{Hom}(\Gamma(p, q, r), \mathrm{SL}(3, \mathbb{R}))$ such that $\rho(R_i)$ is reflection fixing the line containing s_i . Furthermore assume that the differential

$$D\rho(R_1, R_2)_{s_1 \cap s_2} \in \mathrm{GL}(\mathbb{T}_{s_1 \cap s_2} \mathbb{RP}^2)$$

is conjugate to a rotation of angle $2\pi/p$, with similar statements for $\rho(R_2 R_3)$ and $\rho(R_3 R_1)$, the respective vertices $s_2 \cap s_3$ and $s_3 \cap s_1$, and exponents p, q, r . Denote by $\mathcal{R}(p, q, r)$ the set of equivalence classes of such representations.

- Show that such a ρ determines a proper free action of $\Gamma(p, q, r)$ on an open domain $\Omega \subset \mathbb{RP}^2$ with fundamental domain Δ .
- Show that $\Gamma(p, q, r)$ admits torsionfree finite index subgroups Γ_f such that Ω/Γ_f is a convex \mathbb{RP}^2 -manifold.
- Compute the dimension of $\mathcal{R}(p, q, r)$.
- If $1/p + 1/q + 1/r \leq 1$, then show $\mathcal{R}(p, q, r)$ is a cell of dimension 0 or 1 depending on the number of p, q, r equal to 2. In particular if $p, q, r \geq 3$, then $\mathcal{R}(p, q, r)$ is homeomorphic to \mathbb{R} .

This is one of the first examples of a *thin subgroup* of a simple Lie group. Compare also Long-Reid-Thistlethwaite [203], where this example is embedded in a (discrete) one-parameter family of subgroups of $\mathrm{SL}(3, \mathbb{Z})$. For the complete classification of convex \mathbb{RP}^2 -orbifolds, see Choi-Goldman [71]

For the deformation theory of convex structures on the three-holed sphere P see [122], which is the first step in the construction of coordinates on $\mathbb{RP}_{\mathrm{convex}}^2(\Sigma)$.

One notable new feature in the projective theory of pants not seen for the classical case of hyperbolic structures is that the geometric

structure on P is not determined by the structure ∂P . The structure at ∂P is determined by the conjugacy classes of the respective holonomies around the boundary components. The conjugacy classes range over a 2-dimensional space, giving 6 dimensions to the structures on ∂P . However the full deformation space has dimension 8, so there are two more *internal parameters* involved in the deformations of a pants. Finding geometric meaning to these internal parameters has been an intriguing and tantalizing problem. Once the boundary parameters (2 dimensions for each of the three boundary components) are prescribed, there are two internal parameters, and the *relative* deformation space is a 2-cell. See also Zhang [292], Wienhard-Zhang [284], Bonahon-Dreyer [44] and Bonahon-Kim [45] for a discussion of the internal parameters.

In terms of the parameters for triangle groups, Guichard [140] estimates the Hölder exponent of the limit set in terms of ratios of eigenvalues of elements, bounding from below. In his Master's Thesis, Lukyanenko [205] conjectures that this bound is obtained for the Coxeter element $\rho(R_1 R_2 R_3)$ in the family described in Exercise 13.2.1.

EXERCISE 13.2.2. *Formulate and prove Lukyanenko's conjecture.*

13.2.2. Generalized Fenchel-Nielsen earthquakes. One would like to develop a theory of twist flows for $\mathbb{R}P^2$ -manifolds analogous to the earthquake flows described in §7.3.2.

As described in §7.3.2, deformations supported in a tubular neighborhood of a curve \mathcal{C} correspond to paths \mathfrak{z}_t in the centralizer of the holonomy $\rho(c)$, where c is a based loop freely homotopic to \mathcal{C} . For example, suppose that $\rho(\gamma) \in \mathbf{SL}(3, \mathbb{R})$ is the diagonal matrix

$$(81) \quad \gamma := \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} \in \mathbf{SL}(3, \mathbb{R})$$

with $\lambda_1 > \lambda_2 > \lambda_3$. The identity component of its centralizer consists of diagonal matrices

$$A(s, t) := \begin{bmatrix} e^s & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^{-s-t} \end{bmatrix}$$

where $s, t \in \mathbb{R}$.

Analogous to the Fenchel-Nielsen earthquake flows discussed in §7.3.2 and the formula for the geodesic length function on $\mathfrak{F}(\Sigma)$ in terms of the invariant function ℓ discussed in Exercise 7.4.5 are functions and flows on $\mathbb{R}P_{\text{convex}}^2(\Sigma)$. If $\rho \in \text{Hom}(\pi, \mathbf{SL}(3, \mathbb{R}))$ is the holonomy

representation of a marked convex \mathbb{RP}^2 -structure on Σ , then for every $c \in \pi \setminus \{1\}$, the the holonomy $\rho(c)$ conjugate to a diagonal matrix

$$(82) \quad \rho(c) \sim \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$$

with $\lambda_1 > \lambda_2 > \lambda_3$. Denote the invariant open set comprising such matrices by \mathfrak{D}^+ , and define an invariant function

$$\begin{aligned} \mathfrak{D}^+ &\xrightarrow{f} \mathbb{R}plus \\ \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} &\longmapsto \log(\lambda_1) - \log(\lambda_3) \end{aligned}$$

EXERCISE 13.2.3. *Let $\rho \in \text{Hom}(\pi, \text{SL}(3, \mathbb{R}))$ be a holonomy representation of a marked convex \mathbb{RP}^2 -manifold M in $\mathbb{RP}_{\text{convex}}^2(\Sigma)$.*

- *Show that the function $f_c(\rho)$ is length of the unique closed geodesic homotopic to \mathcal{C} computed with respect to the Hilbert metric on M (the Hilbert geodesic length function).*
- *Compute the variation function F of f and identify the corresponding flow on $\mathbb{RP}_{\text{convex}}^2(\Sigma)$ in terms of the Hilbert metric on the corresponding \mathbb{RP}^2 -manifold, analogous to Exercise 7.4.6.*

13.2.3. Bulging deformations. We describe here a general construction of such convex domains as limits of *piecewise conic* curves.

If Ω/Γ is a convex \mathbb{RP}^2 -manifold homeomorphic to a closed esurface S with $\chi(S) < 0$, then every element $\gamma \in \Gamma$ is *positive hyperbolic*, that is, conjugate in $\text{SL}(3, \mathbb{R})$ to a diagonal matrix of the form

$$A(s, t) := \begin{bmatrix} e^s & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^{-s-t} \end{bmatrix}.$$

where $s > t > -s - t$. Its centralizer is the *maximal \mathbb{R} -split torus* \mathbf{A} consisting of all diagonal matrices in $\text{SL}(3, \mathbb{R})$. It is isomorphic to a Cartesian product $\mathbb{R}^\times \times \mathbb{R}^\times$ and has four connected components. Its identity component \mathbf{A}^+ consists of diagonal matrices with positive entries.

The orbits of H_t are arcs of conics depicted in Figure 13.2.

Associated to any measured geodesic lamination λ on a hyperbolic surface S is *bulging deformation* as an \mathbb{RP}^2 -surface. Namely, one applies




FIGURE 13.2. Conics tangent to a triangle

a one-parameter group of collineations

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

to the coordinates on either side of a leaf. This extends Thurston’s *earthquake* deformations (the analog of *Fenchel-Nielsen twist deformations* along possibly infinite geodesic laminations), and the *bending deformations* in $\mathrm{PSL}(2, \mathbb{C})$. See Bonahon-Dreyer [44]. This real 2-parameter family is analogous to the *quakebend* deformations defined by McMullen [218].

In general, if S is a convex \mathbb{RP}^2 -manifold, then deformations are determined by a geodesic lamination with a transverse measure taking values in the Weyl chamber of $\mathfrak{sl}(3, \mathbb{R})$. When S is itself a hyperbolic surface, all the deformations in the singular directions become earthquakes and deform $\partial\tilde{S}$ trivially (just as in $\mathrm{PSL}(2, \mathbb{C})$).

EXERCISE 13.2.4. Let M be a marked convex \mathbb{RP}^2 -manifold with holonomy representation $\rho \in \mathrm{Hom}(\pi, \mathrm{SL}(3, \mathbb{R}))$.

- If \mathcal{C} is a simple closed curve, show that the generalized Fenchel-Nielsen flow commutes with the bulging flow, generating an \mathbb{R}^2 -action associated to \mathcal{C} .
- If $\mathcal{C}_1, \mathcal{C}_2$ are disjoint simple closed curves, show that the corresponding \mathbb{R}^2 -actions commute. Therefore a pants decomposition \mathcal{P} of Σ determines an $(\mathbb{R}^2)^{3g-3}$ -action on $\mathbb{RP}^2(\Sigma)$
- Define a mapping

$$\xrightarrow{(\ell, \beta)_{\mathcal{P}}} (\mathbb{R}^+ \times \mathbb{R})^{3g-3}$$

which has the structure of a principal $(\mathbb{R}^2)^{3g-3}$ -bundle with this action.

- *Identify the fibers with Cartesian products of the deformation spaces of marked convex \mathbb{RP}^2 -structures on the components P_i , ($i = 1, \dots, 2g - 2$) of the complement $\Sigma \setminus \mathcal{P}$ with prescribed boundary structures.*

This is the analog of Fenchel-Nielsen coordinates on \mathbb{RP}^2 , once the internal parameters of the P_i are identified.




FIGURE 13.3. Deforming a conic




FIGURE 13.4. A piecewise conic

13.3. Coordinates for convex structures

To describe generalized Fenchel-Nielsen coordinates on $\mathbb{RP}_{\text{convex}}^2(\Sigma)$, one needs coordinates in the case Σ is a 3-holed sphere (a *pair of pants*). Various approaches exist; in




FIGURE 13.5. Bulging data




FIGURE 13.6. The deformed conic




FIGURE 13.7. The conic with its deformation

13.3.1. Fock-Goncharov coordinates. In their paper [101], Fock and Goncharov develop an ambitious program for studying surface

group representations into split \mathbb{R} -forms, and develop natural coordinates on certain components discovered by Hitchin [151]. and studied by Labourie [198].

A version when $G = \mathrm{SL}(3, \mathbb{R})$ is developed in Fock-Goncharov [102], giving coordinates on the deformation space of convex \mathbb{RP}^2 -structures. Compare also Ovsienko-Tabachnikov [231].

A new object in their theory is the *triple ratio*, a projective invariant of three flags in \mathbb{RP}^2 in general position. A *flag* in \mathbb{RP}^2 is an *incident pair* (p, ℓ) , where $p \in \mathbb{RP}^2$ and $\ell \in (\mathbb{RP}^2)^*$. Incidence here simply means that $p \in \ell$. Two flags (p_1, ℓ_1) and (p_2, ℓ_2) are in *general position* if and only if $p_i \notin \ell_j$ for $i \neq j$.

EXERCISE 13.3.1. *Show that the projective group acts transitively on the set of general position pairs of flags.*

Now suppose that Ψ is a polarity. A Ψ -flag is a flag of the form (p, ℓ) , where $\ell = \Psi(p)$.

EXERCISE 13.3.2. *Show that the stabilizer of Ψ in the projective group does not act transitively on the set of general position Ψ -flags, and show that the quotient space is one-dimensional, with a coordinate defined by a cross ratio*

The *triple ratio* of three flags

$$(p_1, \ell_1), (p_2, \ell_2), (p_3, \ell_3)$$

in general position is defined as follows. Find vectors \mathbf{v}_i representing p_i for $i = 1, 2, 3$ and covectors ψ_j representing ℓ_j for $j = 1, 2, 3$.

EXERCISE 13.3.3. *Show that the scalar quantity*

$$\frac{\psi_1(\mathbf{v}_2) \psi_2(\mathbf{v}_3) \psi_3(\mathbf{v}_1)}{\psi_1(\mathbf{v}_3) \psi_2(\mathbf{v}_1) \psi_3(\mathbf{v}_2)}$$

is independent of the choices of ψ_i and \mathbf{v}_j , and describes a complete projective invariant of triples of flags in general position.

Starting with an ideal triangulation τ of surface S , Fock and Goncharov attach parameters to the ideal simplices in τ : two for each side of a simplex (corresponding the hyperbolic conjugacy classes in $\mathrm{SL}(3, \mathbb{R})$) and a triple ratio invariant for each simplex. This gives the correct dimension and

13.3.2. Affine spheres and Labourie-Loftin parametrization.

Another more analytic approach is due independently to Labourie [197] and Loftin [197], which we only briefly mention.

We suppose here that Σ is a closed oriented surface of genus $g > 1$ with a marking $\Sigma \rightarrow M$, where M is a convex \mathbb{RP}^2 -manifold. Fix a holonomy homomorphism

$$\pi_1(M) \xrightarrow{\approx} \Gamma < \mathrm{SL}(3, \mathbb{R})$$

and developing map

$$\widetilde{M} \xrightarrow{\mathrm{dev}} \Omega \subset \mathbb{RP}^2.$$

Associated to a convex \mathbb{RP}^2 -manifold is an Γ -equivariant lift of dev to a convex surface in the convex cone $\Omega' \subset \mathbb{A}^3$ covering Ω . which is an *affine sphere*.

In affine differential geometry (see, e.g. Nomizu-Sasaki [229]), an *affine normal* at a point p in a convex surface $S \subset \mathbb{A}^3$ is the line tangent to the curve γ through p formed by the centroids of sections $S \cap P_t$ where P_t are planes parallel to the affine tangent plane $T_p S$.

EXERCISE 13.3.4. *Show that γ has a natural parametrization which can be characterized in terms of the connection on \mathbb{A}^3 and the geometry of S .*

S is an *affine sphere* if all the affine normals to S concur in \mathbb{A}^3 . (This is the analog of an *umbilic point* in Euclidean differential geometry.)

Solving an projectively invariant Monge-Ampère equation, Loftin and Labourie show that a Γ -equivariant affine sphere in \mathbb{A}^3 exists. Furthermore they show that such an affine sphere is determined by a conformal structure on M (making M a *Riemann surface* X) and a *holomorphic cubic differential* on X . In this way $\mathbb{RP}_{\mathrm{convex}}^2(\Sigma)$ identifies with the holomorphic vector bundle over \mathfrak{T}_g whose fibers comprise holomorphic cubic differentials (Labourie [197], Loftin [197]). In particular this fibration over \mathfrak{T}_g is Mod_g -invariant.

The Vinberg metric constructed in §4 also determines a Riemann surface (by taking the conformal structure underlying the Riemannian metric) and also defines a Mod_g -equivariant mapping $\mathbb{RP}_{\mathrm{convex}}^2(\Sigma) \rightarrow \mathfrak{T}_g$. The relation between the Loftin-Labourie metric and the Vinberg metric seems intriguing.

13.4. Pathological developing maps and grafting

The nonconvex (grafted) structures have *pathological developing maps*, even on T^2 . That is, the developing maps are generally not covering spaces onto their images. The developing maps for such \mathbb{RP}^2 -structures in higher genus are onto \mathbb{RP}^2 , and are not covering spaces. In genus one, these Smillie-Thurston structures have non-covering developing maps onto the complement of three points in \mathbb{RP}^2 . Their radiant

suspensions are affine 3-manifolds whose developing maps surject to $\mathbb{R}^3 \setminus \{\mathbf{0}\}$, or the complements but are not covering spaces (indeed the 3-manifolds are mapping tori of periodic automorphisms of surfaces with $\chi < 0$, and are thus aspherical).

We first describe \mathbb{RP}^2 -manifolds with cyclic holonomy generated by a *positive diagonal matrix*

$$(83) \quad \gamma := \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \in \mathrm{SL}(3, \mathbb{R})$$

with $a > b > c$. The corresponding collineation fixes three points

$$p_3 := \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \quad p_2 := \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad p_1 := \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

and preserves the corresponding lines

$$l_3 := [0 \ 0 \ 1], \quad l_2 := [0 \ 1 \ 0], \quad l_1 := [1 \ 0 \ 0]$$

The dynamics vary in the three corresponding affine patches.

$$(x, y) \xrightarrow{A_3} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix},$$

γ acts as the affine expansion fixing the origin $(x, y) = (0, 0) \longleftrightarrow p_3$:

$$(x, y) \xrightarrow{\gamma} \begin{bmatrix} a/c x \\ b/c y \\ 1 \end{bmatrix}$$

with eigenvalues $a/c > b/c > 1$. Similarly, in the affine chart \mathcal{A}_1 with ideal line l_1 ,

$$(u, v) \xrightarrow{A_1} \begin{bmatrix} 1 \\ u \\ v \end{bmatrix},$$

γ acts as the affine *contraction* fixing the origin $(u, v) = (0, 0) \longleftrightarrow p_1$: having eigenvalues $1 > c/a > b/a$. Denote the corresponding affine patches $\mathbf{A}_3 := \mathbb{RP}^2 \setminus l_3$ and $\mathbf{A}_1 := \mathbb{RP}^2 \setminus l_1$ respectively.

13.4.0.1. *Smillie-Thurston example.* Here is the first example of a closed \mathbb{RP}^2 -manifold whose developing map is not a covering space. (See Sullivan-Thurston [262] and Smillie [253]; actually [253] describes the radiant suspension.)

The Smillie-Thurston example arises as follows. The collineation γ generates a discrete cyclic subgroup $\Gamma := \langle \gamma \rangle < \mathrm{PGL}(3, \mathbb{R})$ which acts properly on the complements $\mathbf{A}_1 \setminus \{p_1\}$ and $\mathbf{A}_3 \setminus \{p_3\}$ respectively.

The quotients are Hopf tori modeled on the affine spaces A_1 and A_3 respectively, which we denote:

$$T_i := (A_i \setminus \{p_i\})/\Gamma \text{ for } i = 1, 3$$

and regard them as $\mathbb{R}P^2$ -manifolds. We may choose developing maps

$$\tilde{T}_i \xrightarrow{\text{dev}_i} \mathbb{R}P^2.$$

The line l_2 is invariant under Γ , and the fixed points p_1, p_3 of Γ separate l_2 into two open intervals. Choose one such interval J ; then the image

$$c_i = \Pi_i(\text{dev}_i^{-1}(J))$$


is a closed geodesic on T_i . There exist tubular neighborhoods \mathcal{N}_i of $c_i \subset T_i$ for $i = 1, 3$ such that a projective isomorphism $\mathcal{N}_1 \xrightarrow{j} \mathcal{N}_3$ of $\mathbb{R}P^2$ -manifolds exists.

Let M be the $\mathbb{R}P^2$ -manifold obtained by *grafting* T_1 to T_3 along j :

$$M := T_1|_{c_1} \bigcup_j T_3|_{c_3}$$

Then M is an $\mathbb{R}P^2$ manifold homeomorphic to a 2-torus with holonomy group Γ and whose developing map dev_M surjects onto the complement $\mathbb{R}P^2 \setminus \{p_1, p_2, p_3\}$.

EXERCISE 13.4.1. *Prove that dev_M is not a covering space onto its image.*






FIGURE 13.8. Pathological development for \mathbb{RP}^2 -torus

This example gives a counterexample to the main technical lemma of [120], (Theorem 2.2) where it is asserted that if M is a closed (G, X) -manifold with holonomy $\Gamma < G$ and $\Omega \subset X$ is a Γ -invariant open subset of X with a Γ -invariant complete Riemannian metric g_Ω , then the region $M_\Omega \subset M$ corresponding to Ω inherits a complete Riemannian metric from g_Ω .

EXERCISE 13.4.2. *Find a counterexample to this assertion inside the Smillie-Thurston example.*

See §14.2 for a correct proof of the main theorem of [120].

Complex-projective structures

From the general viewpoint of locally homogeneous geometric structures, \mathbb{CP}^1 -manifolds occupy a central role. Historically these objects arose from the applying the theory of second-order holomorphic linear differential equations to conformal mapping of plane domains. Theoretically these objects seem to be fundamental in so many homogeneous spaces extend the geometry of \mathbb{CP}^1 . Furthermore \mathbb{CP}^1 -manifolds play a fundamental role in the theory of hyperbolic 3-manifolds and classical Kleinian groups.

A \mathbb{CP}^1 -manifold has the underlying structure as a *Riemann surface*. Starting from a Riemann surface M , a compatible \mathbb{CP}^1 -structure is a (holomorphic) *projective structure* on the Riemann surface M . Remarkably, projective structures on a Riemann surface M admit an extraordinarily clean classification: the deformation space of projective structures on a fixed Riemann surface M is a complex affine space whose underlying vector space is the space $H^0(M; \kappa^2)$ of *holomorphic quadratic differentials* on M . We describe this parametrization, following Gunning [142, 143].

However, the geometry of the developing map can become extremely complicated, despite this clean determination of the deformation space.

An alternate synthetic-geometry parametrization of $\mathbb{CP}^1(\Sigma)$ is due to Thurston, involves locally convex developments into hyperbolic 3-manifolds. In this case $\mathbb{CP}^1(\Sigma)$ identifies with the product of the space $\mathfrak{F}(\Sigma)$ of marked hyperbolic structures on Σ and the *Thurston cone* $\mathcal{ML}(\Sigma)$ of *measured geodesic laminations* on Σ . Although Thurston's parametrization and Poincaré's parametrization have the same crude topological consequence:

$$\mathbb{CP}^1(\Sigma) \approx \mathbb{R}^{12g-12},$$

they are extremely different, both in content and context. For an exposition of Thurston's parametrization, see Kamishima-Tan [165]. This generalizes more directly to higher dimensions, and more closely relates to topological constructions with the developing map, such as *grafting*.

The grafting construction was first developed by Hejhal [146] and Maskit [214] (Theorem 5) and Sullivan-Thurston [262]. As for $\mathbb{R}\mathbb{P}^2$ -surfaces studied in the previous chapter, this construction yields pathological developing maps, typically local homeomorphisms from the universal covering space *onto* all of $\mathbb{C}\mathbb{P}^1$. Grafting is also responsible for the *non-injectivity* of the holonomy mapping

$$\mathbb{C}\mathbb{P}^1(\Sigma) \xrightarrow{\text{hol}} \text{Hom}(\pi_1(\Sigma), \text{PSL}(2, \mathbb{C})) / \text{PSL}(2, \mathbb{C}),$$

that is, when the holonomy representation does *not* determine the structure. This consequence of grafting was already noted in §5.4.5 for closed $\mathbb{R}\mathbb{P}^1$ -manifolds. (Compare also Goldman [120], Gallo-Kapovich-Marden [112] and Baba [15, 14, 13, 12]) and Baba-Gupta [16].

We will only touch on the subject, which has a vast and ever-expanding literature. We refer to the excellent survey article of Dumas [91] for more details. See also Kapovich [167], §7, Hubbard [154], §6.3 and Marden [206], §6-8 for other perspectives on the subject.

These structures extend, in higher dimensions, to *flat conformal structures*, upon which we only discuss briefly in § We refer to the excellent survey article of Matsumoto [215] for more details.

Another direction in which these structures generalize is to *holomorphic projective structures* on complex manifolds. We do not discuss these structures here, referring instead to Klingler [174], Dumitrescu [92], and McKay [217].

14.1. Schwarzian parametrization

For the remainder of this chapter, we denote by X the Riemann surface underlying a $\mathbb{C}\mathbb{P}^1$ -manifold. Denote by \mathbb{P}^1 the complex projective line, with automorphism group $\text{PGL}(2, \mathbb{C})$. Since \mathbb{P}^1 is the Riemann sphere $\mathbb{C} \cup \{\infty\}$, holomorphic mappings $X \rightarrow \mathbb{P}^1$ are simply meromorphic functions on X . Indeed, we choose a marked Riemann surface

$$\Sigma \xrightarrow{\approx} X$$

representing a point $\mathfrak{F}(\Sigma)$ (which we abusively call X) and consider the subset $\mathcal{P}(X)$ of $\mathbb{C}\mathbb{P}^1(\Sigma)$ with underlying marked Riemann surface X . This is the fiber of the forgetful map $\mathbb{C}\mathbb{P}^1(\Sigma) \rightarrow \mathfrak{F}(\Sigma)$ over X .

We show that $\mathcal{P}(X)$ is a (complex) affine space whose underlying vector space identifies with the vector space $H^0(X, \kappa^2)$ comprising *holomorphic quadratic differentials* on X . The key to this construction is the *Schwarzian differential* \mathcal{S} which assigns a holomorphic mapping f defined on a subset $\Omega \subset \mathbb{P}^1$ a quadratic differential $\Phi \in H^0(\Omega, \kappa^2)$. In




FIGURE 14.1. Octagonal fundamental domain for a \mathbb{CP}^1 -surface of genus two

terms of a local coordinate z , the quadratic differential is

$$\Phi = \phi(z)dz^2$$

where $\Omega \xrightarrow{\phi} \mathbb{C}$ is holomorphic. We apply \mathcal{S} to the developing map to obtain the *Schwarzian parameter* Φ . This operator is $\mathrm{PSL}(2, \mathbb{C})$ -invariant and satisfies a *transformation law* making it a cocycle from the pseudogroup of local biholomorphisms f defined on Ω , taking values in the presheaf $H^0(\Omega, \kappa^2)$ of holomorphic quadratic differentials:

$$(84) \quad \mathcal{S}(f \circ g) = g^*\mathcal{S}(f) + \mathcal{S}(f)$$

It defines the *projective sub-pseudogroup* in the sense that $\mathcal{S}(f) = 0$ if and only if f is locally projective.

14.1.1. Affine structures and the complex exponential map.

We begin with the easier case of the differential operator, the *pre-Schwarzian* \mathcal{A} , defining the *afine sub-pseudogroup*. Namely, let $\Omega \subset \mathbb{P}^1$ and $\Omega \xrightarrow{f} \mathbb{P}^1$ a holomorphic mapping which is a *local biholomorphism* at every $z \in \Omega$, that is,

$$f'(z) \neq 0,$$

for all $z \in \Omega$. The differential operator \mathcal{A} associates to f the *Abelian differential*, that is, the holomorphic 1-form

$$(85) \quad \mathcal{A}(f) := d \log f' = \frac{f''(z)}{f'(z)} dz$$

If g is another local biholomorphism, the Chain Rule implies that wherever the composition $g \circ f$ is defined,

$$(g \circ f)' = (g' \circ f) \cdot f',$$

so

$$\log(g \circ f)' = \log(g' \circ f) + \log f',$$

and differentiating

$$(86) \quad \mathcal{A}(f \circ g) = g^* \mathcal{A}(f) + \mathcal{A}(f)$$

where $g^*(\phi(z)dz)$ denotes the natural action of g on the Abelian differential $\phi(z)dz$:

$$g^*(\phi(z)dz) := \phi(g(z))g'(z)dz$$

The transformation law (86) asserts that \mathcal{A} is a *cocycle* from the pseudogroup of biholomorphisms to the presheaf $\Omega \mapsto H^0(\Omega, \kappa)$ of Abelian differentials.

The cocycle condition (86) asserts $\text{Ker}(\mathcal{A})$ is closed under composition (wherever defined) and defines a subpseudogroup. These are the *locally affine* mappings defined by $f''(z) = 0$.

Furthermore, if X is a Riemann surface and $X \xrightarrow{f} \mathbf{P}^1$ is a local biholomorphism, (86) implies that the restrictions $\mathcal{A}(f|_U)$ to coordinate patches $U \subset \Omega$ extend to a globally defined Abelian differential $\mathcal{A}(f) \in H^0(X; \kappa)$.

14.1.2. Projective structures and quadratic differentials.

Now we deduce the form of the Schwarzian from its projective invariance, using the cocycle property of the differential operator \mathcal{A} and the Bruhat decomposition of the projective group $G := \text{PSL}(2, \mathbb{C})$ by reducing the calculation of $\mathcal{A}(g)$ for general $g \in G$ to that of $\mathcal{A}(\mathcal{J})$ where $\mathcal{J}(z) := 1/z$

EXERCISE 14.1.1. *The projective group G is generated by the affine group $\mathfrak{B} := \text{Aff}(1, \mathbb{C})$ (its Borel subgroup) and the inversion*

$$z \xrightarrow{\mathcal{J}} -1/z,$$

(the generator of the Weyl group).

- Deduce the Bruhat decomposition

$$G = \mathfrak{B} \coprod \mathfrak{B} \mathfrak{J} \mathfrak{B}$$

explicitly from the following identity in $\mathrm{SL}(2, \mathbb{C})$:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1/c & a \\ c & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & d/c \\ 0 & 1 \end{bmatrix}$$

when $c \neq 0$ and $ad - bc = 1$.

- Show that

$$\mathcal{A}(\mathfrak{J}) = \frac{-2}{z} dz$$

- Suppose

$$g(z) = \frac{az + b}{cz + d}.$$

Then

$$\mathcal{A}(g) = \frac{-2}{z + d/c} dz$$

- Define a differential operator:

$$\begin{aligned} H^0(\Omega, \kappa) &\xrightarrow{\mathrm{D}} H^0(\Omega, \kappa^2) \\ \phi(z)dz &\longmapsto \phi'(z)dz^2 \end{aligned}$$

Show that

$$(87) \quad \mathrm{D}\mathcal{A}(g) - \frac{1}{2}\mathcal{A}(g)^2 = 0$$

The formula (87) is reminiscent of the expression of the curvature of a connection in terms of a connection 1-form and the *Fundamental Theorem of Calculus* for Lie group-valued differential forms (Sharpe [249]).

Now define the *Schwarzian differential*

$$\begin{aligned} \mathrm{Bihol}(\Omega, \mathbb{P}^1) &\xrightarrow{\mathfrak{S}} H^0(\Omega, \kappa^2) \\ g &\longmapsto \mathrm{D}\mathcal{A}(g) - \frac{1}{2}\mathcal{A}(g)^2 = \left\{ \left(\frac{g''}{g'}\right)' - \frac{1}{2}\left(\frac{g''}{g'}\right)^2 \right\} dz^2 \end{aligned}$$

where $\mathrm{Bihol}(\Omega, \mathbb{P}^1)$ denotes the space of holomorphic maps $\Omega \rightarrow \mathbb{P}^1$ which are *local biholomorphisms*, that is meromorphic functions g on Ω for which $g'(z) \neq 0$ everywhere on Ω . (The usual *Schwarzian derivative*, denoted classically by $\{g, z\}$, is the coefficient

$$\left(\frac{g''}{g'}\right)' - \frac{1}{2}\left(\frac{g''}{g'}\right)^2 = \left(\frac{g'''}{g'}\right)' - \frac{3}{2}\left(\frac{g''}{g'}\right)^2$$

of the Schwarzian derivative $\mathfrak{S}(g)$.)

EXERCISE 14.1.2. Prove the cocycle property (84).

Although (84) follows from a slightly messy but straightforward calculation (see Gunning [142] for example), a more conceptual treatment is discussed in Hubbard [154]. If $g \in \text{Bihol}(\Omega, \mathbb{P}^1)$ is a local biholomorphism, and $z_0 \in \Omega$, denote the element of G which agrees with $g \in G$ to second order at z_0 by $\mathcal{O}(g)_{z_0} \in G$, the *osculating Möbius transformation* for g at z_0 .

Explicitly, write

$$g(z) = a_0 + a_1(z - z_0) + \frac{a_2}{2}(z - z_0)^2 + \frac{a_3}{6}(z - z_0)^3 + \cdots + \frac{a_n}{n!}(z - z_0)^n + \cdots$$

with $a_0 = g(z_0)$, $a_1 = g'(z_0) \neq 0$, $a_2 = g''(z_0)$ and $a_3 = g'''(z_0)$. The Möbius transformation

$$z \xrightarrow{\mathcal{O}(g)_{z_0}} \frac{a_1 z}{1 + (a_2/a_1)z}$$

osculates g at z_0 and

$$f(z) = \mathcal{O}(g)_{z_0}(z) + \left(\frac{a_3}{6} - \frac{a_2^2}{4a_1} \right) z^3 + \cdots$$

The leading term equals $1/a_1$ times the coefficient of $\mathcal{S}(g)$ at z_0 , that is, the Schwarzian derivative of g at z_0 . From this conceptual description of $\mathcal{S}(g)$ follows the cocycle property (84).

14.1.3. Solving the Schwarzian equation. To show that $\mathcal{P}(X)$ is an $H^0(X, \kappa^2)$ -torsor, one must show that for any holomorphic quadratic differential $\Phi = \phi(z)dz^2$, a meromorphic function $f \in \text{Bihol}(X, \mathbb{P}^1)$ exists with $\mathcal{S}(f) = \Phi$. Furthermore f is unique up to a Möbius transformation. That is, given ϕ , find f solving the nonlinear third order differential equation

$$(88) \quad \left(\frac{f''}{f'} \right)' - \frac{1}{2} \left(\frac{f''}{f'} \right)^2 = \phi.$$

The solution involves relating this nonlinear equation to the second order *linear* differential equation

$$(89) \quad u''(z) + \frac{1}{2}\phi(z)u(z) = 0$$

often called Hill's equation or a Sturm-Liouville equation.

EXERCISE 14.1.3. Define the projective solution $\tilde{X} \xrightarrow{f} \mathbb{P}^1$ of (89) as follows. Every $x \in X$ has a neighborhood Ω such that the solutions

of the linear equation (89) on Ω is a 2-dimensional (complex) vector space. Choose a basis $u_1(z), u_2(z)$ of this vector space and define

$$\begin{aligned} \Omega &\xrightarrow{f} \mathbf{P}^1 \\ z &\longmapsto u_1(z)/u_2(z) \end{aligned}$$

- Define the monodromy representation

$$\pi_1(X) \longrightarrow \mathrm{GL}(2, \mathbb{C})$$

of (89) by analytically continuing the local solutions u_1, u_2 over loops in X .

- f is well-defined up to composition with a projective automorphism $\mathbf{P}^1 \rightarrow \mathbf{P}^1$.
- The derivative $f'(z) \neq 0$ for all $z \in \Omega$.
- $\mathcal{S}(f) = \phi(z)dz^2$.
- The projective solution f on Ω analytically continues to a local biholomorphism

$$\tilde{X} \xrightarrow{\tilde{f}} \mathbf{P}^1$$

equivariant with respect to the projectivization $\pi_1(X) \longrightarrow \mathrm{PGL}(2, \mathbb{C})$ of the monodromy representation of (89).

14.2. Fuchsian holonomy

A related idea is the classification of projective structures with Fuchsian holonomy [120]. This is the converse to the *grafting construction*, whereby grafting is the only construction yielding geometric structures with the same holonomy. Recall that a *Fuchsian representation* of a surface group π is an embedding of π as a discrete subgroup of the group $\mathrm{PGL}(2, \mathbb{R}) \cong \mathrm{Isom}(\mathbb{H}^2)$. Equivalently, ρ is the holonomy representation of a hyperbolic structure on a surface Σ with $\pi_1(\Sigma) \cong \pi$. The main result is:

THEOREM 14.2.1 (Grafting Theorem). *Let M be a closed $\mathbb{C}\mathbf{P}^1$ -manifold whose holonomy representation $\pi_1(M) \xrightarrow{\rho} \mathrm{PSL}(2, \mathbb{C})$ is a composition*

$$\pi_1(M) \xrightarrow{\rho_0} \mathrm{PGL}(2, \mathbb{R}) \hookrightarrow \mathrm{PSL}(2, \mathbb{C})$$

where ρ_0 is Fuchsian representation. Let M_0 be a hyperbolic structure with holonomy representation ρ_0 , regarded as a $\mathbb{C}\mathbf{P}^1$ -manifold. Then there is a unique multicurve $S \subset M_0$ such that M is obtained from M_0 by grafting along S .

However the proof contained a gap, which I first learned from M. Kapovich, who pointed me to the paper of Kuiper [189]. (The same

gap can be found in papers of Goldman-Kamishima [113] and Faltings [99].) This gap was later filled by Choi-Lee [75]. Here we give a correct proof, based on Kulkarni-Pinkall [194] (Theorem 4.2), communicated to me by Daniele Alessandrini. See also Dupont [93].

The grafting theorem uses the setup of Exercise 5.2.8, which we briefly recall. Let M be a connected smooth manifold with a universal covering space $\widetilde{M} \xrightarrow{\Pi} M$ with covering group $\pi = \pi_1(M)$. Give M a (G, X) -structure and let (dev, ρ) be a developing pair:

- $\widetilde{M} \xrightarrow{\text{dev}} X$ denotes the developing map;
- $\pi \xrightarrow{\rho} \Gamma < G$ denotes the holonomy representation.

Suppose that $\Omega \subset X$ is a Γ -invariant open subset and

$$M_\Omega := \Pi(\text{dev}^{-1}\Omega) \subset M$$

the subdomain of M corresponding to Ω as in Exercise 5.2.8.

The author is grateful to Daniele Alessandrini for patiently explaining the details of the following basic result. Recall from §2.6.4 that the *normality domain* $\text{Nor}(\Gamma, X)$ consists of points having open neighborhoods U such that

$$\Gamma|_U := \{\gamma|_U \mid g \in \Gamma\} \subset \subset \text{Map}(U, X).$$

THEOREM 14.2.2 (Kulkarni-Pinkall [194], Theorem 4.2). *Let M be a closed (G, X) -manifold with holonomy group $\Gamma < G$. Let $\Omega \subset \text{Nor}(\Gamma, X)$ be a Γ -invariant subset of the normality domain, and $M_\Omega \subset M$ the corresponding region of M . Then for each component $W \subset M_\Omega$, and each component \widetilde{W} of $\Pi^{-1}(W)$, the restriction*

$$\widetilde{W} \xrightarrow{\text{dev}|_{\widetilde{W}}} \Omega$$

is a covering space. In particular $\text{dev}|_{\widetilde{W}}$ is onto.

The proof of Theorem 14.2.2 breaks into a sequence of lemmas. We show that $\text{dev}|_{\widetilde{W}}$ satisfies the *path-lifting criterion* for covering spaces: every path in Ω lifts to a path in \widetilde{W} . Specifically, let

$$[0, 1] \xrightarrow{\gamma} \Omega$$

be a path in Ω and consider a point

$$\tilde{w}_0 \in \Pi^{-1}(\gamma(0)) \cap \widetilde{W}$$

We seek a path

$$[0, 1] \xrightarrow{\tilde{c}} \widetilde{W}$$

satisfying:

- $\tilde{c}(0) = \tilde{w}_0$;

- $\text{dev} \circ \tilde{c} = \gamma$.

Since dev is a local homeomorphism, the set T of $t \in [0, 1]$ such that $[0, t] \xrightarrow{\gamma|_{[0,t]}} \Omega$ lifts to

$$[0, t] \xrightarrow{\tilde{c}|_{[0,t]}} \tilde{W}$$

is open. Since dev is a local homeomorphism, any extension $\tilde{c}|_{[0,t]}$ of the lift to $[0, t]$ is necessarily unique. Thus T is a *connected* open neighborhood of 0 in $[0, 1]$. By reparametrizing \tilde{c} , we may assume that \tilde{c} is defined on $[0, 1)$. It suffices to show that \tilde{c} lifts to $[0, 1]$.

Let $c = \Pi \circ \tilde{c}$ be the curve in M . Since M is compact and

$$[0, 1) \xrightarrow{c} \Pi(\tilde{W}) \subset M,$$

the curve c *accumulates* in M . That is, a sequence $t_n \in [0, 1)$ with $t_n \nearrow 1$ and $z \in M$ exists, such that

$$\lim_{n \rightarrow \infty} c(t_n) = z.$$

Employ z as the basepoint in M . Fix the corresponding universal covering space $\tilde{M} \xrightarrow{\Pi} M$, where \tilde{M} comprises relative homotopy of paths γ in M starting at z . Recall that the deck transformation of \tilde{M} corresponding to the relative homotopy class $[\beta] \in \pi_1(M, z)$ of a loop β based at z is:

$$[\gamma] \mapsto [\gamma \star \beta].$$

Choose a developing map $\tilde{M} \xrightarrow{\text{dev}} X$.

Let $U \ni z$ be an evenly covered coordinate patch in M such that the restriction of dev to some (and hence every) component of $\Pi^{-1}(U)$ is a homeomorphism. Passing to a subsequence if necessary, we may assume that $c(t_n) \in U$ for all n . Choose paths $z \xrightarrow{\alpha_n} c(t_n)$ in U . Then the concatenation

$$\alpha_1 \star c|_{[t_1, t_n]} \star \alpha_n^{-1}$$

is a based loop β_n in M having relative homotopy class $[\beta_n] \in \pi_1(M, z)$. Let \tilde{U} be the component of $\Pi^{-1}(U)$ containing $\tilde{c}(t_1)$, and \tilde{z} the unique element of $\tilde{U} \cap \Pi^{-1}(z)$. To simplify notation, denote the deck transformation $[\beta_n]$ by β_n .

LEMMA 14.2.3.

$$\lim_{n \rightarrow \infty} \rho(\beta_n)^{-1} \gamma(t_n) = \text{dev}(\tilde{z}).$$

PROOF. For $n \gg 1$, each $c(t_n) \in U$. The definition of the deck transformation β_n implies that $\tilde{c}(t_n) \in \beta_n \tilde{U}$. Hence

$$\beta_n^{-1} \tilde{c}(t_n) \in \tilde{U}.$$

Since $c(t_n) \rightarrow z$ and $\Pi|_{\tilde{U}}$ is bijective, $\lim_{n \rightarrow \infty} \beta_n^{-1} \tilde{c}(t_n) = \tilde{z}$. Continuity of \mathbf{dev} implies:

$$\begin{aligned} \lim_{n \rightarrow \infty} \rho(\beta_n^{-1})\gamma(t_n) &= \lim_{n \rightarrow \infty} \rho(\beta_n^{-1})\mathbf{dev}(\tilde{c}(t_n)) = \lim_{n \rightarrow \infty} \mathbf{dev}(\beta_n^{-1}\tilde{c}(t_n)) \\ &= \mathbf{dev}\left(\lim_{n \rightarrow \infty} \beta_n^{-1}\tilde{c}(t_n)\right) = \mathbf{dev}(\tilde{z}), \end{aligned}$$

as claimed. \square

CONCLUSION OF THE PROOF OF THEOREM 14.2.2. Apply the condition of normality to the images

$$\rho(\beta_n)^{-1} \circ \gamma$$

of the curve $[0, 1] \xrightarrow{\gamma} \Omega$. By the definition of Ω , each $\gamma(s)$ has an open neighborhood U_s for which the set of restrictions $\rho(\beta_n)|_{U_s}$ is precompact in $\mathbf{Map}(U_s, X)$. Compactness of $[0, 1]$ guarantees finitely many s_i exist so that the U_{s_i} cover $[0, 1]$. It follows that the images $\rho(\beta_n)^{-1} \circ \gamma$ form a precompact sequence in $\mathbf{Map}([0, 1], X)$. After passing to a subsequence, we may assume that $\rho(\beta_n)^{-1} \circ \gamma$ converges uniformly to a continuous map $[0, 1] \xrightarrow{\delta} X$.

Now apply Lemma 14.2.3, using the uniform convergence

$$\rho(\beta_n)^{-1} \circ \gamma \rightrightarrows \delta,$$

obtaining

$$\lim_{t \rightarrow 1} \mathbf{dev}(\tilde{c}(t)) = \delta(1) \in \mathbf{dev}(\tilde{U}).$$

Since $\mathbf{dev}|_{\tilde{U}}$ is injective, defining

$$\tilde{c}(1) := (\mathbf{dev}|_{\beta_N \tilde{U}})^{-1}(\delta(1))$$

is the desired continuous extension of \tilde{c} . The proof of Theorem 14.2.2 is complete. \square

CHAPTER 15

Geometric structures on 3-manifolds

This final chapter we collect a few results on geometric structures on closed 3-manifolds.

However, the theory is very much in its infancy and certain innocent-sounding questions seem (at least now) to be inaccessible.

Two exceptions are the theory of complete affine structures on 3-manifolds, and the classification of projective structures on 3-manifolds with solvable fundamental group.

We begin with a brief overview of flat conformal and spherical CR-structures, where many examples exist in dimension three; flat conformal structures naturally extend the \mathbb{CP}^1 -structures discussed in the previous chapter. Then we survey the classification of complete affine 3-manifolds, and close with Dupont's classification of affine structures on 3-manifolds with solvable fundamental group.

15.1. Higher dimensions: flat conformal and spherical CR-structures

These structures generalize to (G, X) -structures where G is a semisimple Lie group and $X = G/P$, where $P \subset G$ is a parabolic subgroup. The simplest generalization occurs when $G = \mathbf{SO}(n+1, 1)$ and $X = S^n$. The conformal automorphisms of S^n are just Möbius transformations. In this case X is the model space for *conformal (Euclidean) geometry* and a (G, X) -structure is a *flat conformal structure*, that is, a conformal equivalence class of conformally flat Riemannian metrics.

A key point in this identification is the famous result of Liouville, that, in dimensions > 2 , a conformal map from a nonempty connected domain in S^n is the restriction of a unique Möbius transformation of S^n .

Furthermore this is the boundary structure for hyperbolic structures in dimension $n+1$, since $S^n = \partial\mathbf{H}_{\mathbb{R}}^{n+1}$ and the group of isometries of \mathbf{H}^{n+1} restricts to the group of conformal automorphisms of S^n .

A good general survey of this subject is Matsumoto [215]). Kami-shima-Tan [165] and Dumas [91]) describe the unpublished construction of Thurston (using hyperbolic geometry) which identifies a flat conformal structure with a hyperbolic structure with the extra structure of a measured geodesic lamination; roughly speaking the $\mathbb{C}P^1$ -structure is identified with an equivariant map of the universal covering into \mathbb{H}^3 which is locally convex and *pleated* (piecewise totally geodesic). This has been extended to higher dimensional flat conformal structures by Kulkarni and Pinkall [194, 195].

Some of the most interesting examples are due to Gromov-Lawson-Thurston [136] and Kuiper [193]. While products $\Sigma \times S^1$ (where Σ is a closed hyperbolic surface) admit flat conformal structures, 3-dimensional nilmanifolds and hyperbolic torus bundles do *not* admit such structures (Goldman [117]). However, [136, 193] produce examples of flat conformal structures on *twisted* oriented S^1 -bundles over closed hyperbolic surfaces.

Another interesting example is *spherical CR-geometry*, the boundary structure for complex hyperbolic geometry is a *spherical CR-structure*, where $G = \mathrm{PU}(n, 1)$ and $X = \partial\mathbb{H}_{\mathbb{C}}^n \approx S^{2n-1}$.

One of the first papers on this subject is Burns-Shnider [53] which computed the homogenous domains. This geometry is extensively discussed in Goldman [124]. For more information on this very active field of research see the papers of Schwartz [245], Parker, Falbel, Deraux, Paupert, Will and others.

While [117] shows that T^3 and hyperbolic torus bundles do not admit spherical CR-structures, some twisted S^1 -bundles over closed surfaces do. Ananin, Grossi and Gusevskii [5] produce surprising examples of spherical CR-structures on products $S \times S^1$.

EXERCISE 15.1.1. *Find examples of closed 3-manifolds with flat conformal (respectively spherical CR-manifolds) whose developing maps are surjective but not covering spaces.*

The classification of complete affine structures on *closed* 3-manifolds has been understood since the early 1980's, see Fried-Goldman [121, 110]. The considerably more interesting case of *noncompact* complete 3-manifolds has only been understood recently. The big breakthrough came in the early 1980's with Margulis's resolution [208] of Milnor's question [225]; see Abels [1] and Goldman [127, 128] for expositions. For a summary of this theory, starting with its historical origins and leading to current research, see the article for the Feitschrift honoring the seventieth birthday of G. Margulis by Danciger-Drumm-Goldman-Smilga [83]. We only give a brief summary of these developments here.

Drumm's thesis introduced a more geometric approach to these questions involving hypersurfaces called *crooked planes*. He constructed Schottky groups with fundamental domains bounded by crooked planes to give many examples of Margulis spacetimes. Now through the remarkable work of Danciger-Guéritaud-Kassel [84, 85] (based on an analysis two-generator groups in [60, 61, 51, 62]). (Compare also Guéritaud's survey [139].) Many of these results extend to the broader (and quite fascinating) study of constant curvature Lorentzian manifolds. In particular the study of 3-dimensional *anti-de Sitter manifolds*, and its extension to flat conformal Lorentzian 3-manifolds is quite fascinating. Anti-de Sitter 3-space is the Lorentzian analog of hyperbolic 3-space. A suggestive model arises from the universal covering space $X := \widetilde{\mathrm{SL}(2, \mathbb{R})}$ and

$$G := (\widetilde{\mathrm{SL}(2, \mathbb{R})} \times \widetilde{\mathrm{SL}(2, \mathbb{R})}) / Z,$$

where $Z \subset \widetilde{\mathrm{SL}(2, \mathbb{R})} \times \widetilde{\mathrm{SL}(2, \mathbb{R})}$ is the diagonally embedding of the infinite cyclic group

$$\mathrm{center}(\widetilde{\mathrm{SL}(2, \mathbb{R})}) \cong \pi_1(\mathrm{SL}(2, \mathbb{R})) \cong \mathbb{Z}.$$

For this geometry, G is the group $\mathrm{SO}(2, 2)$ acting on but for reasons of space, we do not discuss these structures, instead we refer to articles by Schlenker, Tholozan, and Danciger-Guéritaud and Kassel. This geometry also lies in the flat conformal Lorentzian geometry, where G is the *Einstein Universe*, consisting of null lines in a Lorentzian vector space; compare Barbot-Charette-Drumm-Goldman [20].

For possibly incomplete structures, much less is known. We then describe a few cases where one has definitive information, including the case of closed affine 3-manifolds with nilpotent holonomy and the beautiful classification of Serge Dupont [94] of affine structures on hyperbolic torus bundles.

Finally we discuss a few results concerning geometric structures on closed 3-manifolds, in particular \mathbb{RP}^3 -structures, flat conformal structures, and spherical CR-structures.

Cooper-Goldman [77] show that the connected sum $\mathbb{RP}^3 \# \mathbb{RP}^3$ fails to admit an \mathbb{RP}^3 -structure. To the author's knowledge, this is the only closed 3-manifold known *not* to admit a projective structure.

15.2. Complete affine 3-manifolds

A more extensive recent survey of this subject is Danciger-Drumm-Goldman-Smilga [83].

Complete affine structures on 3-manifolds were classified by Fried-Goldman [110]. They are finitely covered by complete affine solvmanifolds (see §8.6.2 of Chapter 8) and thus relate to left-invariant affine structures on 3-dimensional Lie groups (see Chapter 10).

The first step in the classification is the Milnor-Auslander question: Namely, if $M^3 = \mathbf{A}^3/\Gamma$, show that Γ is solvable. Let $A(\Gamma)$ denote the Zariski closure of Γ in $\text{Aff}(\mathbf{A}^3)$; clearly it suffices to show that $A(\Gamma)$ is solvable. This is equivalent to showing that the Zariski closure $A(L(\Gamma))$ in $\text{GL}(\mathbb{R}^3)$ of the linear holonomy group $L(\Gamma) \subset \text{GL}(\mathbb{R}^3)$ is solvable. The proof is a case-by-case analysis of the possible Levi factors of $A(L(\Gamma))$.

Complete affine structures on Euclidean 3-manifolds are classified using 3-dimensional commutative associative algebras; see Chapter 10, Exercise 10.5.3.

In his 1977 paper [225], Milnor set the record straight caused by the confusion surrounding Auslander’s flawed proof of Conjecture 8.6.2. Influenced by Tits’s work [269] on free subgroups of linear groups and amenability, Milnor observed, that for an affine space \mathbf{A} of given dimension, the following conditions are all equivalent:

- Every discrete subgroup of $\text{Aff}(\mathbf{A})$ which acts properly on \mathbf{A} is amenable.
- Every discrete subgroup of $\text{Aff}(\mathbf{A})$ which acts properly on \mathbf{A} is virtually solvable.
- Every discrete subgroup of $\text{Aff}(\mathbf{A})$ which acts properly on \mathbf{A} is virtually polycyclic.
- A nonabelian free subgroup of $\text{Aff}(\mathbf{A})$ admits no proper action on \mathbf{A} .
- The Euler characteristic $\chi(\Gamma \backslash \mathbf{A})$ (when defined) of a complete affine manifold $\Gamma \backslash \mathbf{A}$ must vanish (unless $\Gamma = \{1\}$ of course).
- A complete affine manifold $\Gamma \backslash \mathbf{A}$ has finitely generated fundamental group Γ .

(If these conditions were met, one would have a satisfying structure theory, similar to, but somewhat more involved, than the Bieberbach structure theory for flat Riemannian manifolds.)

In [225], Milnor provides abundant “evidence” for this “conjecture”. For example, the *infinitesimal version*: Namely, let $G \subset \text{Aff}(\mathbf{A})$ be a connected Lie group which acts properly on \mathbf{A} . Then G must be an amenable Lie group, which simply means that it is a compact extension of a solvable Lie group. (Equivalently, its Levi subgroup is compact.) Furthermore, he provides a *converse*: Milnor shows that every virtually polycyclic group admits a proper affine action. (However, Milnor’s actions do *not* have compact quotient. Benoist [24, 26]

found finitely generated nilpotent groups which admit no affine crystallographic action. Benoist's examples are 11-dimensional.)

However convincing as his "evidence" is, Milnor still proposes a possible way of constructing counterexamples:

"Start with a free discrete subgroup of $O(2, 1)$ and add translation components to obtain a group of affine transformations which acts freely. However it seems difficult to decide whether the resulting group action is properly discontinuous."

This is clearly a geometric problem: As Schottky showed in 1907, free groups act properly by isometries on hyperbolic 3-space, and hence by diffeomorphisms of A^3 . These actions are *not* affine.

One might try to construct a proper affine action of a free group by a construction like Schottky's. Recall that a *Schottky group of rank g* is defined by a system of g open half-spaces H_1, \dots, H_g and isometries A_1, \dots, A_g such that the $2g$ half-spaces

$$H_1, \dots, H_g, A_1(H_1^c), \dots, A_g(H_g^c)$$

are all disjoint (where H^c denotes the *complement* of the closure \bar{H} of H). The *slab*

$$\text{Slab}_i := H_i^c \cap A_i(H_i)$$

is a fundamental domain for the action of the cyclic group $\langle A_i \rangle$. The *ping-pong lemma* then asserts that the intersection of all the slabs

$$\Delta := \text{Slab}_1 \cap \dots \cap \text{Slab}_g$$

is a fundamental domain for the group $\Gamma := \langle A_1, \dots, A_g \rangle$. Furthermore Γ is freely generated by A_1, \dots, A_g . The basic idea is the following. Let $B_i^+ := A_i(H_i^c)$ (respectively $B_i^- := H_i$) denote the *attracting basin* for A_i (respectively A_i^{-1}). That is, A_i maps all of H_i^c to B_i^+ and A_i^{-1} maps all of $A_i(H_i)$ to B_i^- . Let $w(a_1, \dots, a_g)$ be a reduced word in abstract generators a_1, \dots, a_g , with initial letter a_i^\pm . Then

$$w(A_1, \dots, A_g)(\Delta) \subset B_i^\pm.$$

Since all the basins B_i^\pm are disjoint, $w(A_1, \dots, A_g)$ maps Δ off itself. Therefore $w(A_1, \dots, A_g) \neq 1$.

Freely acting discrete cyclic groups of affine transformations have fundamental domains which are *parallel slabs*, that is, regions bounded by two parallel affine hyperplanes. One might try to combine such slabs to form "affine Schottky groups", but immediately one sees this idea is doomed, if one uses parallel slabs for Schottky's construction: parallel slabs have disjoint complements only if they are parallel to each other, in which case the group is necessarily cyclic anyway. From

this viewpoint, a discrete group of affine transformations seems very unlikely to act properly.

15.3. Margulis spacetimes

In the early 1980's Margulis, while trying to prove that a nonabelian free group can't act properly by affine transformations, discovered that discrete free groups of affine transformations can indeed act properly!

Around the same time, David Fried and I were also working on these questions, and reduced Milnor's question in dimension three to what seemed at the time to be one annoying case which we could not handle. Namely, we showed the following: Let A be a three-dimensional affine space and $\Gamma \subset \text{Aff}(A)$. Suppose that Γ acts properly on A . Then either Γ is polycyclic or the restriction of the linear holonomy homomorphism

$$\Gamma \xrightarrow{L} \text{GL}(A)$$

discretely embeds Γ onto a subgroup of $\text{GL}(A)$ conjugate to the orthogonal group $\text{O}(2, 1)$.

In particular the complete affine manifold $M^3 = \Gamma \backslash A$ is a *complete flat Lorentz 3-manifold* after one passes to a finite-index torsion-free subgroup of Γ to ensure that Γ acts freely. In particular the restriction $L|_{\Gamma}$ defines a free properly discrete isometric action of Γ on the hyperbolic plane \mathbb{H}^2 and the quotient $\Sigma^2 := \mathbb{H}^2/L(\Gamma)$ is a complete hyperbolic surface with a homotopy equivalence

$$M^3 := \Gamma \backslash A \simeq \mathbb{H}^2/L(\Gamma) =: \Sigma^2.$$

Already this excludes the case when M^3 is compact, since Γ is the fundamental group of a closed aspherical 3-manifold (and has cohomological dimension 3) and the fundamental group of a hyperbolic surface (and has cohomological dimension ≤ 2). This is a crucial step in the proof of Conjecture 8.6.2 in dimension 3.

That the hyperbolic surface Σ^2 is *noncompact* is a much deeper result due to Geoffrey Mess [220]. Later proofs and a generalization have been found by Goldman-Margulis [134] and Labourie [196]. (Compare the discussion in §15.3.3.) Since the fundamental group of a noncompact surface is free, Γ is a free group. Furthermore $L|_{\Gamma}$ embeds Γ as a free discrete group of isometries of hyperbolic space. Thus Milnor's suggestion is the *only* way to construct nonsolvable examples *in dimension three*.

15.3.1. Affine boosts and crooked planes. Since L embeds Γ_0 as the fundamental group of a hyperbolic surface, $L(\gamma)$ is elliptic only

if $\gamma = 1$. Thus, if $\gamma \neq 1$, then $L(\gamma)$ is either hyperbolic or parabolic. Furthermore $L(\gamma)$ is hyperbolic for most $\gamma \in \Gamma_0$.

When $L(\gamma)$ is hyperbolic, γ is an *affine boost*, that is, it has the form

$$(90) \quad \gamma = \begin{bmatrix} e^{\ell(\gamma)} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-\ell(\gamma)} \end{bmatrix} \begin{bmatrix} 0 \\ \alpha(\gamma) \\ 0 \end{bmatrix}$$

in a suitable coordinate system. (Here the 3×3 matrix represents the linear part, and the column 3-vector represents the translational part.) γ leaves invariant a unique (spacelike) line C_γ (the second coordinate line in (90)). Its image in $E^{2,1}/\Gamma$ is a *closed geodesic* $C_\gamma/\langle\gamma\rangle$. Just as for hyperbolic surfaces, most loops in M^3 are freely homotopic to such closed geodesics. (For a more direct relationship between the dynamics of the geodesic flows on Σ^2 and M^3 , compare Goldman-Labourie

Margulis observed that C_γ inherits a natural orientation and metric, arising from an orientation on A , as follows. Choose repelling and attracting eigenvectors $L(\gamma)^\pm$ for $L(\gamma)$ respectively; choose them so they lie in the same component of the nullcone. Then the orientation and metric on C_γ is determined by a choice of nonzero vector $L(\gamma)^0$ spanning $\text{Fix}(L(\gamma))$. this vector is uniquely specified by requiring that:

- $L(\gamma)^0 \cdot L(\gamma)^0 = 1$;
- $(L(\gamma)^0, L(\gamma)^-, L(\gamma)^+)$ is a positively oriented basis.

The restriction of γ to C_γ is a translation by displacement $\alpha(\gamma)$ with respect to this natural orientation and metric.

Compare this to the more familiar *geodesic length function* $\ell(\gamma)$ associated to a class γ of closed curves on the hyperbolic surface Σ . The linear part $L(\gamma)$ acts by *transvection* along a geodesic $c_{L(\gamma)} \subset H^2$. The quantity $\ell(\gamma) > 0$ measures how far $L(\gamma)$ moves points of $c_{L(\gamma)}$.

This pair of quantities

$$(\ell(\gamma), \alpha(\gamma)) \in \mathbb{R}_+ \times \mathbb{R}$$

is a complete invariant of the isometry type of the *flat Lorentz cylinder* $A/\langle\gamma\rangle$. The absolute value $|\alpha(\gamma)|$ is the length of the unique primitive closed geodesic in $A/\langle\gamma\rangle$.

A fundamental domain is the parallel slab

$$(\Pi_{C_\gamma})^{-1} (p_0 + [0, \alpha(\gamma)] \gamma^0)$$

where

$$A \xrightarrow{\Pi_{C_\gamma}} C_\gamma$$

denotes orthogonal projection onto

$$C_\gamma = p_0 + \mathbb{R}\gamma^0.$$

As noted above, however, parallel slabs can't be combined to form fundamental domains for Schottky groups, since their complementary half-spaces are rarely disjoint.

In retrospect this is believable, since these fundamental domains are fashioned from the dynamics of the translational part (using the projection Π_{C_γ}). While the effect of the translational part is properness, the dynamical behavior affecting most points is influenced by the *linear part*: While points on C_γ are displaced by γ at a polynomial rate, all other points move at an exponential rate.

Furthermore, parallel slabs are less robust than slabs in \mathbb{H}^2 : while small perturbations of one boundary component extend to fundamental domains, this is no longer true for parallel slabs. Thus one might look for other types of fundamental domains better adapted to the exponential growth dynamics given by the linear holonomy $L(\gamma)$.

Todd Drumm, in his 1990 Maryland thesis [89, 90], defined more flexible polyhedral surfaces, which can be combined to form fundamental domains for *Schottky groups* of 3-dimensional affine transformations. A *crooked plane* is a PL surface in \mathbf{A} , separating \mathbf{A} into two *crooked halfspaces*. The complement of two disjoint crooked halfspace is a *crooked slab*, which forms a fundamental domain for a cyclic group generated by an affine boost. Drumm proved the remarkable theorem that if S_1, \dots, S_g are crooked slabs whose complements have disjoint interiors, then given any collection of affine boosts γ_i with S_i as fundamental domain, then the intersection $S_1 \cap \dots \cap S_g$ is a fundamental domain for $\langle \gamma_1, \dots, \gamma_g \rangle$ acting on *all* of \mathbf{A} .

Modeling a crooked fundamental domain for Γ acting on \mathbf{A} on a fundamental polygon for Γ_0 acting on \mathbb{H}^2 , Drumm proved the following sharp result:

THEOREM (Drumm). *Every noncompact torsion-free Fuchsian group Γ_0 admits a proper affine deformation Γ whose quotient is a solid handlebody.*

15.3.2. Marked length spectra. We now combine the geodesic length function $\ell(\gamma)$ describing the geometry of the hyperbolic surface Σ with the Margulis invariant $\alpha(\gamma)$ describing the Lorentzian geometry of the flat affine 3-manifold M .

As noted by Margulis, $\alpha(\gamma) = \alpha(\gamma^{-1})$, and more generally

$$\alpha(\gamma^n) = |n|\alpha(\gamma).$$

The invariant ℓ satisfies the same homogeneity condition, and therefore

$$\frac{\alpha(\gamma^n)}{\ell(\gamma^n)} = \frac{\alpha(\gamma)}{\ell(\gamma)}$$

is constant along hyperbolic cyclic subgroups. Hyperbolic cyclic subgroups correspond to periodic orbits of the geodesic flow ϕ on the unit tangent bundle $U\Sigma$. Periodic orbits, in turn, define ϕ -invariant probability measures on $U\Sigma$. Goldman-Labourie-Margulis prove that, for any affine deformation, this function extends to a continuous function Υ_Γ on the space $\mathcal{C}(\Sigma)$ of ϕ -invariant probability measures on $U\Sigma$. Furthermore when Γ_0 is convex cocompact (that is, contains no parabolic elements), then the affine deformation Γ acts properly if and only if Υ_Γ never vanishes. Since $\mathcal{C}(\Sigma)$ is connected, nonvanishing implies either all $\Upsilon_\Gamma(\mu) > 0$ or all $\Upsilon_\Gamma(\mu) < 0$. From this follows Margulis's *Opposite Sign Lemma*, first proved in to groups with parabolics by Charette and Drumm [59]:

THEOREM (Margulis). *If Γ acts properly, then all of the numbers $\alpha(\gamma)$ have the same sign.*

For an excellent treatment of the original proof of this fact, see the survey article of Abels [1].

15.3.3. Deformations of hyperbolic surfaces. The Margulis invariant may be interpreted in terms of deformations of hyperbolic structures as follows

Suppose Γ_0 is a Fuchsian group with quotient hyperbolic surface $\Sigma_0 = \Gamma_0 \backslash \mathbb{H}^2$. Let \mathfrak{g}_{Ad} be the Γ_0 -module defined by the adjoint representation applied to the embedding $\Gamma_0 \hookrightarrow \mathbf{O}(2, 1)$. The coefficient module \mathfrak{g}_{Ad} corresponds to the Lie algebra of *right-invariant* vector fields on $\mathbf{O}(2, 1)$ with the action of $\mathbf{O}(2, 1)$ by left-multiplication. Geometrically these vector fields correspond to the infinitesimal isometries of \mathbb{H}^2 .

A family of hyperbolic surfaces Σ_t smoothly varying with respect to a parameter t determines an *infinitesimal deformation*, which is a cohomology class $[u] \in H^1(\Gamma_0, \mathfrak{g}_{\text{Ad}})$. The cohomology group $H^1(\Gamma_0, \mathfrak{g}_{\text{Ad}})$ corresponds to *infinitesimal deformations* of the hyperbolic surface Σ_0 . In particular the tangent vector to the path Σ_t of marked hyperbolic structures smoothly varying with respect to a parameter t defines a cohomology class

$$[u] \in H^1(\Gamma_0, \mathfrak{g}_{\text{Ad}}).$$

The same cohomology group parametrizes affine deformations. The translational part u of a linear representations of Γ_0 is a cocycle of the

group Γ_0 taking values in the corresponding Γ_0 -module \mathbf{V} . Moreover two cocycles define affine deformations which are conjugate by a translation if and only if their translational parts are cohomologous cocycles. Therefore translational conjugacy classes of affine deformations form the cohomology group $H^1(\Gamma_0, \mathbf{V})$. Inside $H^1(\Gamma_0, \mathbf{V})$ is the subset **Proper** corresponding to *proper affine deformations*.

The adjoint representation Ad of $\text{O}(2, 1)$ identifies with the orthogonal representation of $\text{O}(2, 1)$ on \mathbf{V} . Therefore the cohomology group $H^1(\Gamma_0, \mathbf{V})$ consisting of translational conjugacy classes of affine deformations of Γ_0 can be identified with the cohomology group $H^1(\Gamma_0, \mathfrak{g}_{\text{Ad}})$ corresponding to infinitesimal deformations of Σ_0 .

THEOREM. *Suppose $u \in Z^1(\Gamma_0, \mathfrak{g}_{\text{Ad}})$ defines an infinitesimal deformation tangent to a smooth deformation Σ_t of Σ .*

- *The marked length spectrum ℓ_t of Σ_t varies smoothly with t .*
- *Margulis's invariant $\alpha_u(\gamma)$ represents the derivative*

$$\left. \frac{d}{dt} \right|_{t=0} \ell_t(\gamma)$$

- *(Opposite Sign Lemma) If $[u] \in \text{Proper}$, then all closed geodesics lengthen (or shorten) under the deformation Σ_t .*

Since closed hyperbolic surfaces do not support deformations in which *every* closed geodesic shortens, such deformations only exist when Σ_0 is noncompact. This leads to a new proof [134] of Mess's theorem that Σ_0 is not compact. (For another, somewhat similar proof, which generalizes to higher dimensions, see Labourie [196].)

The tangent bundle $\text{T}G$ of any Lie group G has a natural structure as a Lie group, where the fibration $\text{T}G \xrightarrow{\Pi} G$ is a homomorphism of Lie groups, and the tangent spaces

$$\text{T}_x G = \Pi^{-1}(x) \subset \text{T}G$$

are vector groups. The deformations of a representation $\Gamma_0 \xrightarrow{\rho_0} G$ correspond to representations $\Gamma_0 \xrightarrow{\rho} \text{T}G$ such that $\Pi \circ \rho = \rho_0$. In our case, affine deformations of $\Gamma_0 \hookrightarrow \text{O}(2, 1)$ correspond to representations in the tangent bundle $\text{TO}(2, 1)$. When G is the group $\mathbf{G}(\mathbb{R})$ of \mathbb{R} -points of an algebraic group \mathbf{G} defined over \mathbb{R} , then

$$\text{T}G \cong \mathbf{G}(\mathbb{R}[\epsilon])$$

where ϵ is an indeterminate with $\epsilon^2 = 0$. (Compare [125].) This is reminiscent of the classical theory of quasi-Fuchsian deformations of

Fuchsian groups, where one deforms a Fuchsian subgroup of $\mathrm{SL}(2, \mathbb{R})$ in

$$\mathrm{SL}(2, \mathbb{C}) = \mathrm{SL}(2, \mathbb{R}[i])$$

where $i^2 = -1$.

15.3.4. Classification. In light of Drumm's theorem, classifying Margulis spacetimes M^3 begins with the classification of hyperbolic structures Σ^2 . Thus the deformation space of Margulis spacetimes maps to the Fricke space $\mathfrak{F}(\Sigma)$ of marked hyperbolic structures on the underlying topology of Σ .

The main result of is that the positivity (or negativity) of Υ_Γ on $\mathcal{C}(\Sigma)$ is necessary and sufficient for properness of Γ . (For simplicity we restrict ourselves to the case when $L(\Gamma)$ contains no parabolics — that is, when Γ_0 is convex cocompact.) Thus the proper affine deformation space **Proper** identifies with the open convex cone in $H^1(\Gamma_0, \mathbb{V})$ defined by the linear functionals Υ_μ , for μ in the compact space $\mathcal{C}(\Sigma)$. These give uncountably many linear conditions on $H^1(\Gamma_0, \mathbb{V})$, one for each $\mu \in \mathcal{C}(\Sigma)$. Since the invariant probability measures arising from periodic orbits are dense in $\mathcal{C}(\Sigma)$, the cone **Proper** is the interior of half-spaces defined by the countable set of functional Υ_γ , where $\gamma \in \Gamma_0$.

The zero level sets $\Upsilon_\gamma^{-1}(0)$ correspond to affine deformations where γ does not act freely. Therefore **Proper** defines a component of the subset of $H^1(\Gamma_0, \mathbb{V})$ corresponding to affine deformations which are *free* actions.

Actually, one may go further. An argument inspired by Using an argument due to Thurston [264], one reduces the consideration to only those measures arising from *multicurves*, that is, unions of disjoint *simple* closed curves. These measures (after scaling) are dense in the *Thurston cone* $\mathcal{ML}(\Sigma)$ of *measured geodesic laminations* on Σ . One sees the combinatorial structure of the Thurston cone replicated on the boundary of **Proper** $\subset H^1(\Gamma_0, \mathbb{V})$.

Two particular cases are notable. When Σ is a 3-holed sphere or a 2-holed cross-surface (real projective plane), then the Thurston cone degenerates to a finite-sided polyhedral cone. In particular properness is characterized by 3 Margulis functionals for the 3-holed sphere, and 4 for the 2-holed cross-surface. Thus the deformation space of equivalence classes of proper affine deformations is either a cone on a triangle or a convex quadrilateral, respectively.

When Σ is a 3-holed sphere, these functionals correspond to the three components of $\partial\Sigma$. The halfspaces defined by the corresponding three Margulis functionals cut off the deformation space (which is a

polyhedral cone with 3 faces). The Margulis functionals for the other curves define halfspaces which strictly contain this cone.

When Σ is a 2-holed cross-surface these functionals correspond to the two components of $\partial\Sigma$ and the two orientation-reversing simple closed curves in the interior of Σ . The four Margulis functionals describe a polyhedral cone with 4 faces. All other closed curves on Σ define halfspace strictly containing this cone.

In both cases, an ideal triangulation for Σ models a crooked fundamental domain for M , and Γ is an affine Schottky group, and M is an open solid handlebody of genus 2 (Charette-Drumm-Goldman depicts these finite-sided deformation spaces).

For the other surfaces where $\pi_1(\Sigma)$ is free of rank two (equivalently $\chi(\Sigma) = -1$), infinitely many functionals Υ_μ are needed to define the deformation space, which necessarily has infinitely many sides. In these cases M^3 admits crooked fundamental domains corresponding to ideal triangulations of Σ , although unlike the preceding cases there is no single ideal triangulation which works for all proper affine deformations. Once again M^3 is a genus two handlebody.

15.4. Dupont's classification of hyperbolic torus bundles

THEOREM 15.4.1 (Dupont[94]). *Let M^3 be an hyperbolic torus bundle with an affine structure. Then M^3 is a quotient $\Gamma \backslash G$ where G is an affine Lie group isomorphic to $\mathbb{R}^2 \rtimes \mathbb{R}^+$ where $\mathbb{R}^+ \cong \mathrm{SO}(1, 1)$. Furthermore the developing image $\Omega \approx G$ is one of the following:*

- All of \mathbb{A}^3 (M is complete);
- A halfspace in \mathbb{A}^3 ;
- The product with \mathbb{A}^1 with a convex parabolic domain (such as $\{y > x^2\}$);
- The product with \mathbb{A}^1 with a concave parabolic domain (such as $\{y < x^2\}$).

In particular, a developing map embeds \widetilde{M} as one of these domains and the holonomy homomorphism is an isomorphism $\pi_1(M) \xrightarrow{\cong} \Gamma < G$.

This builds on Dupont's classification [93] of affine actions of the two-dimensional group $\mathrm{Aff}(\mathbb{R})$ on \mathbb{A}^3 .

Although the class of affine structures on closed 3-manifolds with *nilpotent* holonomy are understood, the general case of *solvable* holonomy remains mysterious. However, Serge Dupont [94] completely classifies affine structures on 3-manifolds with solvable *fundamental group*. (Compare also Dupont [93], in the volume [156].) In terms of the Thurston geometrization, these are the geometric 3-manifolds

modeled on Sol, that is, 3-manifolds finitely covered by *hyperbolic torus bundles*: mapping tori (suspensions) of hyperbolic elements of $\mathrm{GL}(2, \mathbb{Z})$. Dupont shows that all such structures arise from left-invariant affine structures on the corresponding Lie group G , which is the semidirect product of \mathbb{R}^2 by \mathbb{R} , where \mathbb{R} acts on \mathbb{R}^2 as a unimodular hyperbolic one-parameter subgroup (explicitly, G is isomorphic to the identity component in the group of Lorentz isometries of the Minkowski plane).

Two structures are particularly interesting for the behavior of geodesics in light of the results of Vey [277]. Recall that a properly convex domain $\Omega \in \mathbb{A}^n$ is *divisible* if it admits a discrete group $\Gamma < \mathrm{Aut}(\Omega)$ acting properly such that Ω/Γ is compact. (Equivalently, the quotient space Ω/Γ by a discrete subgroup $\Gamma \subset \mathrm{Aut}(\Omega)$ is compact and Hausdorff.) Vey proved that a divisible domain is a cone. However, dropping the properness of the action of Γ on Ω allows counterexamples: the *parabolic cylinder*

$$\Omega := \{(x, y) \in \mathbb{A}^2 \mid y > x^2\}$$

is a properly convex domain which is not a cone, but admits a group Γ of automorphisms such that Ω/Γ is compact but not Hausdorff. (See § 10.4.4 for Lie algebraic properties of Ω .)

Now take the product $\Omega \times \mathbb{R} \subset \mathbb{A}^3$. Let $G < \mathrm{Aff}(\mathbb{A}^3)$ be the subgroup acting simply transitively (isomorphic to the 3-dimensional unimodular exponential non-nilpotent solvable Lie group) discussed in §10.8.1), and let $\Gamma < G$ be a lattice. Then Γ acts properly on $\Omega \times \mathbb{R}$ and:

- The quotient $M = (\Omega \times \mathbb{R})/\Gamma$ is a hyperbolic torus bundle (and in particular compact and Hausdorff);
- $\Omega \times \mathbb{R}$ is not a cone.

Clearly $\Omega \times \mathbb{R}$ is not properly convex, showing that Vey's result is sharp [116].

The Kobayashi pseudometric degenerates along a 1-dimensional foliation of M , and defines the hyperbolic structure transverse to this foliation discussed by Thurston [265], Chapter 4.

APPENDIX A

Transformation groups

A.1. Group actions

We shall consider *left-actions* of a group on a set, unless otherwise noted. Suppose that G is a group acting on a set X , with the (left-) action denoted by:

$$\begin{aligned} G \times X &\xrightarrow{\alpha} X \\ (g, x) &\longmapsto g \cdot x \end{aligned}$$

We refer to X as a (left-) G -set.

The *kernel* of the action α consists of all g such that $\alpha(g, \cdot) = \mathbb{I}$, that is, $g \cdot x = x, \forall x \in X$. Equivalently, this is the kernel of the homomorphism of G into the group of automorphisms of the set X . The action is *effective* (or *faithful*) if its kernel is trivial.

If $x \in X$, its *stabilizer*: is the subgroup:

$$\text{Stab}(x) := \{g \in G \mid g \cdot x = x\}.$$

If $g \in G$, then its *fixed-point set* (or *stationary set*) is the subset:

$$\text{Fix}(g) := \{x \in X \mid g \cdot x = x\}.$$

The action is *free* if and only if $\text{Stab}(x) = \{1\}, \forall x \in X$, or equivalently $\text{Fix}(g) = \emptyset, \forall g \in G, g \neq 1$.

The *orbit* of a point x is the image

$$G \cdot x := \alpha(G \times \{x\}) = \{g \cdot x \mid g \in G\}.$$

The orbits partition X , so that the group action defines an equivalence relation on X . The action is *transitive* if some (and hence every) orbit equals X .

The action is *simply transitive* if it is transitive and free. In terms of the *orbit map* (or *evaluation map*)

$$\begin{aligned} G &\xrightarrow{\alpha_x} X \\ g &\longmapsto \alpha(g, x) = g \cdot x, \end{aligned}$$

- α is free $x \iff \alpha_x$ is injective $\forall x \in X$;
- α is transitive $\iff \alpha_x$ is surjective (for any x);
- α is simply transitive $\iff \alpha_x$ is bijective.

A simply transitive action α makes X into a G -torsor.

EXERCISE A.1.1. Let X be a left G -set and $x \in X$. Let $H = \text{Stab}(x)$.

- The orbit map α_x defines a G -equivariant isomorphism $G/H \rightarrow G \cdot x$ where G acts by left-multiplication on the set G/H of left cosets gH for $g \in G$.
- Suppose that $N < H$ is a nontrivial normal subgroup of G . Then G does not act effectively on G/H .

A.2. Proper and syndetic actions

A convenient context in which to work is that of *locally compact Hausdorff topological spaces* and *topological groups*. Recall that a continuous map $X \xrightarrow{f} Y$ is *proper* if $\forall K \subset\subset Y$, the preimage $f^{-1}(K) \subset\subset X$. Recall the following facts from general topology:

- A closed subset of a compact space is compact.
- The continuous image of a compact space is compact.
- Compact subsets of a Hausdorff space are closed,

It follows that a proper map is closed.

EXERCISE A.2.1. Suppose that X, Y are manifolds and f is a smooth map. Furthermore suppose that f is a local homeomorphism. Then f is a covering space.

Let G be a locally compact Hausdorff topological group and

$$\begin{aligned} G \times X &\xrightarrow{\alpha} X \\ (g, x) &\longmapsto g \cdot x \end{aligned}$$

is a (left) action. We say that the group action α is *proper* if and only if the continuous map

$$\begin{aligned} G \times X &\xrightarrow{f_\alpha} X \times X \\ (g, x) &\longmapsto (g \cdot x, x) \end{aligned}$$

is a proper map.

EXERCISE A.2.2. Show that properness is equivalent to the either of the two following conditions: (for the last condition assume that G is second countable)

- $\forall K_1, K_2 \subset\subset X$, the set

$$G(K_1, K_2) := \{g \in G \mid gK_1 \cap K_2 \neq \emptyset\} \subset\subset G.$$
- $\forall K \subset\subset X$, $G(K, K) \subset\subset G$.

- For all sequences $g_n \in G$, $x_n \in X$, such that $g_n x_n$ converges, the sequence g_n has a convergent subsequence.

For the last condition, we can say that if x_n stays bounded, but $g_n \rightarrow \infty$, then $g_n x_n \rightarrow \infty$.

EXERCISE A.2.3. Suppose that α is a proper action of a locally compact group on a locally compact Hausdorff space X . Then the quotient space $G \backslash X$ is Hausdorff. Is the converse true?

EXERCISE A.2.4. Suppose that Γ is a discrete group and $G \times X \xrightarrow{\alpha} X$ is a proper free action. Then the quotient map $X \rightarrow G \backslash X$ is a covering space.

A group action α is *syndetic* if the quotient $G \backslash X$ is compact.

A.3. Topological transformation groupoids

We must relate the actions of $\text{Aff}(A)$ on $\mathfrak{C}(A)$ and G on $\mathfrak{C}(P)$. Recall that a *topological transformation groupoid* consists of a small category \mathfrak{G} whose objects form a topological space X upon which a topological group G acts such that the morphisms $a \rightarrow b$ consist of all $g \in G$ such that $g(a) = b$. We write $\mathfrak{G} = (G, X)$. A *homomorphism of topological transformation groupoids* is a functor

$$(X, G) \xrightarrow{(f, F)} (X', G')$$

arising from a continuous map $X \xrightarrow{f} X'$ which is equivariant with respect to a continuous homomorphism $G \xrightarrow{F} G'$.

The space of isomorphism classes of objects in a category \mathfrak{G} will be denoted $\text{Iso}(\mathfrak{G})$. We shall say that \mathfrak{G} is *proper* (respectively *syndetic*) if the corresponding action of G on X is proper (respectively syndetic). If \mathfrak{G} and \mathfrak{G}' are topological categories, a functor $\mathfrak{G} \xrightarrow{F} \mathfrak{G}'$ is an *equivalence of topological categories* if the induced map

$$\text{Iso}(\mathfrak{G}) \xrightarrow{\text{Iso}(F)} \text{Iso}(\mathfrak{G}')$$

is a homeomorphism and F is *fully faithful*, that is, for each pair of objects a, b of \mathfrak{G} , the induced map

$$\text{Hom}(a, b) \xrightarrow{F_*} \text{Hom}(F(a), F(b))$$

is a homeomorphism. If F is fully faithful it is enough to prove that $\text{Iso}(F)$ is surjective. (Compare Jacobson [157].) We have the following general proposition:

LEMMA A.3.1. *Suppose that*

$$(X, G) \xrightarrow{(f, F)} (X', G')$$

is a homomorphism of topological transformation groupoids which is an equivalence of groupoids and such that f is an open map.

- *If (X, G) is proper, so is (X', G') .*
- *If (X, G) is syndetic, so is (X', G') .*

PROOF. An equivalence of topological groupoids induces a homeomorphism of quotient spaces

$$X/G \longrightarrow X'/G'$$

so if X'/G' is compact (respectively Hausdorff) so is X/G . Since (X, G) is syndetic if and only if X/G is compact, this proves the assertion about syndeticity. By Koszul [185], p.3, Remark 2, (X, G) is proper if and only if X/G is Hausdorff and the action (X, G) is *wandering* (or *locally proper*): each point $x \in X$ has a neighborhood U such that

$$G(U, U) = \{g \in G \mid g(U) \cap U \neq \emptyset\}$$

is precompact. Since (f, F) is fully faithful, F maps $G(U, U)$ isomorphically onto $G'(f(U), f(U))$. Suppose that (X, G) is proper. Then X/G is Hausdorff and so is X'/G' . We claim that G' acts locally properly on X' . Let $x' \in X'$. Then there exists $g' \in G'$ and $x \in X$ such that $g'f(x) = x'$. Since G acts locally properly on X , there exists a neighborhood U of $x \in X$ such that $G(U, U)$ is precompact. It follows that $U' = g'f(U)$ is a neighborhood of $x' \in X'$ such that $G'(U', U') \cong G(U, U)$ is precompact, as claimed. Thus G' acts properly on X' . \square

APPENDIX B

Affine connections in local coordinates

We summarize some of the basic facts about (not necessarily flat) affine connections on a smooth manifold M . After discussing the expression of general affine connections in local coordinates is a review of the Levi-Civita (or *Riemannian*) connection. As this general theory only uses the nondegeneracy of the metric connection, it applies equally to *pseudo-Riemannian*, structures defined by a (possibly indefinite) metric tensor. Due to its fundamental importance, this appendix ends with a detailed discussion of the Riemannian connection for the hyperbolic plane \mathbf{H}^2 .

B.1. Affine connections in local coordinates

Recall that an *affine connection* on M is a (linear connection) on the tangent bundle $\mathbb{T}M$, which can be defined in several different ways. One of the most familiar ways is that of a *Koszul connection*, that is, a covariant differentiation operation on a vector bundle \mathbb{V} over M , which happens to be isomorphic to the tangent bundle $\mathbb{T}M$. Explicitly we consider an isomorphism of vector bundles over M :

$$\mathbb{T}M \xrightarrow{\cong} \mathbb{V}$$

together with a covariant differentiation operation

$$\begin{aligned} \mathbb{T}_p M \times \Gamma(\mathbb{V}) &\longrightarrow \mathbb{V}_p \\ (X_p, v) &\longmapsto \nabla_{X_p}(v) \end{aligned}$$

which satisfies *tensorial conditions*

$$\nabla_{fX_p}(v) = f(p)\nabla_{X_p}(v), \quad \nabla_{X_p}(fv) = X_p(f)v + f(p)\nabla_{X_p}v$$

for $f \in C^\infty(M)$.

In local coordinates (x^1, \dots, x^n) , the connection is defined by *Christoffel symbols*

$$\nabla_{\partial_i}(\partial_j) := \Gamma_{ij}^k(x)\partial_k.$$

whereby the general covariant derivative of vector fields is:

$$\nabla_{a_i \partial_i} b_j \partial_j = a_i \frac{\partial b_j}{\partial x_i} \Gamma_{ij}^k(x) \partial_k.$$

From an affine connection, define the *torsion* and *curvature* tensors as follows.

EXERCISE B.1.1. Show that the operations on vector fields $X, Y \in \text{Vec}(M)$

$$\begin{aligned}\text{Tor}_\nabla(X, Y) &:= \nabla_X Y - \nabla_Y X - [X, Y] \\ \text{Riem}_\nabla(X, Y) &:= \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]}\end{aligned}$$

are tensorial (that is, multilinear over $C^\infty(M)$), and define smooth sections of the vector bundles $\Lambda^2(\mathbb{T}M) \otimes \mathbb{T}M$ and $\Lambda^2(\mathbb{T}M) \otimes \text{End}(\mathbb{T}M)$, respectively.

In local coordinates on $\mathbb{T}M$, where (x^1, \dots, x^n) are local coordinates on M and (v^1, \dots, v^n) are local coordinates on $\mathbb{T}_p M$, the geodesic equations are:

$$\begin{aligned}\frac{d}{dt}x^k(t) &= v^k(t) \\ \frac{d}{dt}v^i(t) &= -\Gamma_{ij}^k(x)v^i(t)v^j(t)\end{aligned}$$

(for $k = 1, \dots, n$) and $\Gamma_{ij}^k(x)$ are the *Christoffel symbols* defined by:

$$\nabla_{\partial_i}(\partial_j) = \Gamma_{ij}^k(x)\partial_k.$$

In particular the geodesic flow corresponds to the vector field (called the *geodesic spray*):

$$\phi_\Gamma := v^k \frac{\partial}{\partial x^k} - \Gamma_{ij}^k(x)v^i v^j \frac{\partial}{\partial v^k}$$

on $\mathbb{T}M$. See Kobayashi-Nomizu [181], do Carmo [87] or O'Neill [230] for further details.

That ϕ_Γ is *vertically homogeneous of degree one*: That is, it transforms under the one-parameter group of homotheties

$$(p, \mathbf{v}) \xrightarrow{h_t} (p, e^s \mathbf{v})$$

by

$$(91) \quad (h_t)_*(\phi_\Gamma) = e^t \phi_\Gamma$$

Furthermore its trajectories are the velocity vector fields of the *geodesics* on M . Namely, let $p = (x^1, \dots, x^n) \in M$ and $\mathbf{v} = (v^1, \dots, v^n) \in \mathbb{T}_p M$ be an initial condition. Let $\gamma_{p, \mathbf{v}}(t) \in \mathbb{T}M$ denote the trajectory of ϕ_Γ , defined for t in an open interval containing $0 \in \mathbb{R}$, defined by:

$$\left. \frac{\partial}{\partial t} \right|_{t=0} \gamma_{p, \mathbf{v}}(t) = \mathbf{v}, \quad \gamma_{p, \mathbf{v}}(0) = (p, \mathbf{v})$$

Then the local flow Φ^t of ϕ_Γ on $\mathbb{T}M$ satisfies

$$\Phi^t(p, \mathbf{v}) = (\gamma_{p,\mathbf{v}}(t), \gamma'_{p,\mathbf{v}}(t)),$$

where

$$\gamma_{p,\mathbf{v}}(t) := (\Pi \circ \Phi^t)(p, \mathbf{v})$$

and $\mathbb{T}M \xrightarrow{\Pi} M$ denoting the bundle projection. The homogeneity condition (91) implies that

$$\gamma_{p,s\mathbf{v}}(t) = \gamma_{p,\mathbf{v}}(st).$$

We write:

$$(92) \quad \text{Exp}_p(t\mathbf{v}) := \gamma_{p,\mathbf{v}}(t)$$

for $t \in \mathbb{R}$ sufficiently near 0. Then, whenever s, t are sufficiently near 0,

$$\text{Exp}_p((s+t)\mathbf{v}) = \text{Exp}_{\gamma(t)}(s\mathbb{P}(t\mathbf{v}))$$

where, for clarity, we denote $\gamma(t) := \text{Exp}_p(t\mathbf{v})$ and parallel transport $\Pi_p^{\gamma(t)}$ by:

$$\mathbb{T}_p M \xrightarrow{\Pi} \mathbb{T}_{\gamma(t)}.$$

Observe that $\Pi(\mathbf{v}) = \gamma'(t)$.

Compare §8.3 and standard references.

B.2. Projective Equivalence

Projective structures can be defined in terms of affine connections. First we remark that the geodesics of an affine connection ∇ are independent of the torsion. Namely a curve $\gamma(t)$ is a geodesic if and only if

$$\frac{d^2}{dt^2} \gamma^j(t) + \Gamma_{ij}^k \frac{d\gamma^i}{dt} \frac{d\gamma^j}{dt} = 0$$

where the second term is symmetric in i, j . By subtracting $\frac{1}{2}\text{Tor}_\nabla$, we may assume that $\text{Tor}_\nabla = 0$, that is, Γ_{ij}^k is symmetric in i, j .

EXERCISE B.2.1. *Show that two torsionfree affine connections have the same set of parametrized geodesics if and only if they are equal.*

Weyl found an elementary criteria for when two torsionfree affine connections ∇ and $\tilde{\nabla}$ have the same *unparametrized* geodesics. In that case, we say the connections are *projectively equivalent*.

EXERCISE B.2.2. *(Weyl) ∇ and $\tilde{\nabla}$ are projectively equivalent if and only if the difference $\tilde{\nabla} - \nabla$ is the symmetrization of a 1-form ω , that is,*

$$\tilde{\nabla}_X(Y) - \nabla_X Y = \omega(X)Y + \omega(Y)X,$$

$\forall X, Y \in \text{Vec}(M)$. In terms of local coordinates $\omega = \omega_l dx^l$, this means

$$\tilde{\nabla}_{ij}^k - \nabla_{ij}^k = \omega_i \delta_j^k + \omega_j \delta_i^k.$$

For more discussion, see Spivak [258], Addendum 1, Proposition 18 and Ovsienko-Tabachnikov [231], Proposition A.3.2.

EXERCISE B.2.3. In the projective models for affine space and hyperbolic space, the geodesics are segments of projective lines. This gives several examples of projective equivalences. For the invariant affine connection on an affine patches, and the Levi-Civita connection for hyperbolic space in the Beltrami-Klein model, find the 1-forms ω effecting these projective equivalences.

A torsionfree affine connection is *projectively flat* if locally it is projectively equivalent to the standard connection on an affine patch. Projective flatness is detected by a contraction of the curvature tensor. The deformation space $\mathbb{RP}^2(\Sigma)$ can be obtained as the space of projective equivalence classes of projectively flat affine connections. The symplectic structure can be obtained as a double symplectic quotient of the affine space of affine connections on Σ . The first symplectic reduction arises from the action of $\Omega^1(\Sigma)$ generating projective equivalence, and the moment map is Tor . The corresponding symplectic quotient is the space \mathfrak{A} of projective equivalence classes of torsionfree affine connections. The second symplectic reduction arises from the Hamiltonian action of the group $\text{Diff}^0(\Sigma)$ on \mathfrak{A} ; here the moment map is the projective curvature tensor. See Goldman [123] for details.

B.3. The (pseudo-) Riemannian connection

The *fundamental theorem of Riemannian geometry* asserts a pseudo-Riemannian manifold (M, \mathbf{g}) admits a unique affine connection ∇ with natural properties:

- (Orthogonality) $\nabla \mathbf{g} = 0$;
- (Symmetry) $\text{Tor}(\nabla) = 0$.

Orthogonality is equivalent to the condition that the parallel transport operator

$$\mathbb{T}_x M \xrightarrow{\mathbb{P}_\gamma} \mathbb{T}_y M$$

along a path $x \xrightarrow{\gamma} y$ maps $(\mathbb{T}_x, \mathbf{g}_x)$ isometrically to $(\mathbb{T}_y, \mathbf{g}_y)$. When computed with respect to an orthonormal frame, this implies that the Christoffel symbols satisfy:

$$\Gamma_{ij}^k = -\Gamma_{ik}^j,$$

that is, the $n \times n$ -matrix of 1-forms $[\Gamma_{ij}^k dx^1]$ is skew-symmetric.

EXERCISE B.3.1. *The Koszul formula is an remarkable explicit formula for the Levi-Civita connection in terms of the metric tensor \mathbf{g} and the Lie bracket on $\text{Vec}(M)$. Namely, if $X, Y, Z \in \text{Vec}(M)$, then*

$$2\mathbf{g}(\nabla_X Y, Z) = X\mathbf{g}(Y, Z) + Y\mathbf{g}(Z, X) - Z\mathbf{g}(X, Y) \\ - \mathbf{g}(X, [Y, Z]) + \mathbf{g}(Y, [Z, X]) + \mathbf{g}(Z, [X, Y])$$

defines the unique affine ∇ -ortogonal symmetric connection. In local coordinates (x^1, \dots, x^n) ,

$$\Gamma_{ij}^k = \frac{1}{2}\mathbf{g}^{km} \left(\frac{\partial \mathbf{g}_{jm}}{\partial x^i} + \frac{\partial \mathbf{g}_{im}}{\partial x^j} - \frac{\partial \mathbf{g}_{ij}}{\partial x^m} \right)$$

B.4. The Levi-Civita connection for the Poincaré metric

We denote complex numbers $z = x + yi \in \mathbb{C}$ where $x, y \in \mathbb{R}$.

Let \mathbb{H}^2 denote the upper half-plane

$$\{x + iy \mid x, y \in \mathbb{R}, y > 0\}$$

with the *Poincaré metric*:

$$g = \frac{|dz|^2}{y^2}$$

We compute the Levi-Civita connection ∇ with respect to several different frames.

B.4.1. The usual coordinate system on \mathbb{H}^2 . Let ∂_x, ∂_y be the coordinate vector fields, so that:

$$(93) \quad \mathbf{g}(\partial_x, \partial_x) = \mathbf{g}(\partial_y, \partial_y) = y^{-2}$$

$$(94) \quad \mathbf{g}(\partial_x, \partial_y) = \mathbf{g}(\partial_y, \partial_x) = 0.$$

The vector fields

$$\xi := y\partial_x \\ \eta := y\partial_y$$

define an orthonormal frame field. Therefore, for any vector field ϕ ,

$$\phi = \mathbf{g}(\phi, \xi)\xi + \mathbf{g}(\phi, \eta)\eta \\ (95) \quad = y^2 \left(\mathbf{g}(\phi, \partial_x)\partial_x + \mathbf{g}(\phi, \partial_y)\partial_y \right)$$

THEOREM. *In terms of the coordinate frame, the Levi-Civita connection is given by:*

$$(96) \quad \nabla_x \partial_x = y^{-1} \partial_y$$

$$(97) \quad \nabla_x \partial_y = -y^{-1} \partial_x$$

$$(98) \quad \nabla_y \partial_x = -y^{-1} \partial_x$$

$$(99) \quad \nabla_y \partial_y = -y^{-1} \partial_y$$

In terms of the orthonormal frame, the Levi-Civita connection is given by:

$$\nabla_\xi \xi = \eta$$

$$\nabla_\xi \eta = -\xi$$

$$\nabla_\eta \xi = 0$$

$$\nabla_\eta \eta = 0$$

PROOF. Symmetry of ∇ and $[\partial_x, \partial_y] = 0$ implies:

$$(100) \quad \nabla_x \partial_y = \nabla_y \partial_x$$

which implies the equivalence (97) \iff (98).

Differentiate (93) with respect to ∂_x :

$$(101) \quad \mathbf{g}(\nabla_x \partial_x, \partial_x) = 0$$

$$(102) \quad \mathbf{g}(\nabla_x \partial_y, \partial_y) = 0.$$

Combine (102) with (100):

$$(103) \quad \mathbf{g}(\nabla_y \partial_x, \partial_y) = 0$$

Differentiate (93) with respect to ∂_y :

$$2\mathbf{g}(\nabla_y \partial_x, \partial_x) = 2\mathbf{g}(\nabla_y \partial_y, \partial_y) = -2y^{-3},$$

whence

$$(104) \quad \mathbf{g}(\nabla_y \partial_x, \partial_x) = -y^{-3}$$

and

$$(105) \quad \mathbf{g}(\nabla_y \partial_y, \partial_y) = -y^{-3}.$$

Now:

$$(106) \quad \begin{aligned} \mathbf{g}(\nabla_x \partial_x, \partial_y) &= \underbrace{\partial_x \mathbf{g}(\partial_x, \partial_y)}_{0 \text{ by (94)}} - \mathbf{g}(\partial_x, \nabla_x \partial_y) \\ &= -\mathbf{g}(\partial_x, \nabla_y \partial_x) \quad \text{by (100)} \\ &= -\mathbf{g}(\nabla_y \partial_x, \partial_x) = y^{-3} \quad \text{by (103),} \end{aligned}$$

and:

$$(107) \quad \mathbf{g}(\nabla_y \partial_y, \partial_x) = \underbrace{\partial_y \mathbf{g}(\partial_y, \partial_x)}_{0 \text{ by (94)}} - \underbrace{\mathbf{g}(\partial_y, \nabla_y \partial_x)}_{0 \text{ by (103)}} = 0$$

Now we compute the covariant derivatives $\nabla_x \partial_x, \nabla_x \partial_y, \nabla_y \partial_x, \nabla_y \partial_y$ in terms of their inner products:

(98) follows by applying (95) to (103) and (104).

(97) follows by applying (100) to (98), as mentioned above.

(96) follows by applying (95) to (101) and (106).

(99) follows by applying (95) to (107) and (105). □

B.4.2. Connection 1-form for orthonormal frame. Denote the coframe field dual to the orthonormal frame by:

$$\begin{aligned} \xi^* &:= y^{-1} dx \\ \eta^* &:= y^{-1} dy. \end{aligned}$$

The covariant differentials of the orthonormal frame are:

$$\begin{aligned} \nabla \xi &= dx \otimes \partial_y = \xi^* \otimes \eta \\ \nabla \eta &= -dx \otimes \frac{\partial}{\partial x} = -\xi^* \otimes \xi, \end{aligned}$$

which we may write as:

$$\nabla \begin{bmatrix} \xi \\ \eta \end{bmatrix} = \begin{bmatrix} 0 & \xi^* \\ -\xi^* & 0 \end{bmatrix} \otimes \begin{bmatrix} \xi \\ \eta \end{bmatrix} = -\xi^* \mathbf{J} \otimes \begin{bmatrix} \xi \\ \eta \end{bmatrix}$$

where

$$\mathbf{J} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

is the infinitesimal generator of $\text{SO}(2)$. Thus the *connection 1-form* is

$$\mathbf{f} := -y^{-1} dx \mathbf{J}$$

and its derivative is the *curvature 2-form*

$$\mathbf{F} := d\mathbf{f} = -dA \mathbf{J}$$

where $dA := y^{-2} dx \wedge dy$ is the *Poincaré area form*.

B.4.3. Geodesic curvature of a hypercycle. We use these to calculate the geodesic curvature of a hypercycle. The positive imaginary axis $i\mathbb{R}^+$ is a geodesic with endpoints $0, \infty$. Let $-\pi/2 < \theta < \pi/2$. The ray $e^{i\theta}\mathbb{R}^+$ is a hypercycle h_ρ at distance $\rho \in \mathbb{R}$ from the geodesic $i\mathbb{R}^+$, where

$$e^{i\theta} = \cos(\theta) + i \sin(\theta) = \tanh(\rho) + i \operatorname{sech}(\rho)$$

parametrizes the geodesic perpendicular to $i\mathbb{R}^+$ through i with unit speed.

$$\gamma(s) := e^s (\tanh(\rho) + i \operatorname{sech}(\rho))$$

parametrizes h_ρ with unit speed. Its velocity equals

$$\gamma'(s) = e^s (\tanh(\rho)\partial_x + \operatorname{sech}(\rho)\partial_y).$$

Since $\|\gamma'(s)\| = 1$ the acceleration of γ (the vector $\frac{D}{ds}\gamma'(s)$) is orthogonal to γ and the geodesic curvature of γ equals:

$$k_{\mathbf{g}} = \left\| \frac{D}{ds}\gamma(s) \right\| = \tanh(\rho)$$

Similarly the geodesic curvature of a metric circle of radius ρ equals $\coth(\rho)$. To see this, use the Poincare unit disc model: for $|z| < 1$, the metric tensor is:

$$\mathbf{g} := \frac{4|dz|^2}{(1-|z|^2)^2}$$

and writing (hyperbolic polar coordinates)

$$z = e^{i\theta} \tanh(\rho/2)$$

the metric tensor is $\mathbf{g} = d\rho^2 + \sinh^2(\rho)d\theta^2$, with area form $dA = \sinh^2(\rho)d\rho \wedge d\theta$. Consider a disc D_ρ with (hyperbolic) radius ρ . Then its circumference equals $2\pi \sinh(\rho)$ and its area $2\pi(\cosh(\rho) - 1)$. Let $k_{\mathbf{g}}$ be the geodesic curvature of the metric circle ∂D_ρ . Applying the Gauss-Bonnet theorem

$$2\pi\chi(\Sigma) = \int_{\Sigma} K dA + \oint_{\partial\Sigma} k_{\mathbf{g}} ds$$

to $\sigma = D_\rho$ obtaining:

$$2\pi = -2\pi(\cosh(\rho) - 1) + k_{\mathbf{g}}(2\pi \sinh(\rho)),$$

that is,

$$k_{\mathbf{g}} = \coth(\rho)$$

as desired.

APPENDIX C

Facts about metric spaces

This appendix discusses several well-known general facts about metric spaces and pseudometric spaces, which are used in §12 and elsewhere.

C.1. Nonincreasing homeomorphisms are isometries

Here we give the proof of Lemma 4.3.3, which states that a distance non-increasing homeomorphism of a compact metric space (X, d) is an isometry. This fact is used in the proof of Vey's Semisimplicity (Theorem 4.3.1).

The proof given here was found on *Stack Exchange*, submitted by Dap on 1 March 2018.

LEMMA C.1.1. *Suppose (X, d) is a compact metric space and let $X \xrightarrow{f} X$ be a surjective continuous map such that $d(f(x), f(y)) \leq d(x, y)$ for all $x, y \in X$. Then f is an isometry.*

PROOF. Let $\epsilon > 0$. Since X is compact, it admits an $\epsilon/4$ -cover, that is, a set $S := \{s_1, \dots, s_m\} \subset X$ such that $\forall x \in X, \exists s_j \in S$ with $d(s_j, x) < \epsilon/4$. We may assume that S minimizes the number $N(S)$ of pairs $(s_i, s_j) \in S \times S$ with

$$d(s_1, s_2) \geq D := d(x, y) - \epsilon/2.$$

Since f is surjective and distance non-increasing, $f(S)$ is also an $\epsilon/4$ -cover, and $d(s_1, s_2) < D$ implies $d(f(s_1), f(s_2)) < D$. Thus

$$d(f(s_1), f(s_2)) \geq D$$

whenever $d(s_1, s_2) \geq D$ and $s_1, s_2 \in S$.

Picking $s_1, s_2 \in S$ with $d(s_1, x), d(s_2, y) \leq \epsilon/4$ gives $d(s_1, s_2) \geq d(x, y) - \epsilon/2$. As above, $d(f(s_1), f(s_2)) \geq d(x, y) - \epsilon/2$, whence

$$d(f(x), f(y)) \geq d(x, y) - \epsilon.$$

□

C.2. Compactness of distance nonincreasing maps

Recall that a topological space is *separable* if it has a countable dense subset.

LEMMA C.2.1 (Kobayashi [180], Chapter V, Theorem 3.1). *Let (N, d_N) and (M, d_M) be connected locally compact pseudometric spaces. Suppose that (N, d_N) is separable and (M, d_M) is a complete metric space. Then the subset of $\mathbf{Map}(N, M)$ comprising distance-nonincreasing maps*

$$N \xrightarrow{f} M$$

is locally compact. In particular, if $p \in N$ and $K \subset\subset M$, the subset of all such maps with $f(p) \in K$ is compact.

We briefly sketch the proof; see Kobayashi [180] for further details.

Let f_n be a sequence of distance-nonincreasing projective maps taking $p \in N$ to $K \subset\subset M$. Choose a countable dense subset $\{p_1, \dots\} \subset N$; then

$$K_i := \overline{B}_{d_M(K, p_i)}(K) \subset\subset M.$$

Since the f_n are distance-nonincreasing,

$$f_n(p_i) \in K_i.$$

Passing to a subsequence, assume that the sequence $f_n(p_i)$ converges, for each i . Then for each $q \in N$, the sequence $f_n(q)$ is Cauchy. Since N is complete, $f_n(q)$ converges. Finally, this convergence is uniform on compact subsets of N .

C.3. The Lebesgue number of an open covering

LEMMA C.3.1. *Let (X, d) be a compact metric space and \mathcal{U} an open covering of X . Then $\exists \delta > 0$ such that every subset $Y \subset X$ with $\text{diam}(Y) < \delta$ lies in some $U \in \mathcal{U}$.*

PROOF. Because X is compact, \mathcal{U} contains a finite subcovering U_1, \dots, U_n . If some $U_i = X$, then any $\delta > 0$ suffices. Therefore we assume that $\mathcal{U} = \{U_1, \dots, U_n\}$ and every $U_i \subsetneq X$.

Since each $X \setminus U_i$ is closed and nonempty, the map

$$\begin{aligned} X &\xrightarrow{d_i} \mathbb{R} \\ x &\longmapsto d(x, X \setminus U_i) \end{aligned}$$

continuous and positive. If $x \in U_i$ and $j \neq i$, then $d_i(x) > 0$ and $d_j(x) \geq 0$, so

$$\begin{aligned} X &\longrightarrow \mathbb{R} \\ x &\longmapsto \frac{1}{n} \sum_{i=1}^n d_i \end{aligned}$$

is continuous and positive, and has a positive lower bound δ .

Suppose $\text{diam}(Y) < \delta$. The $\exists x_0 \in X$ (for example, take any $x_0 \in Y$) so that $Y \subset B_\delta(x_0)$. Then

$$\delta > \frac{1}{n} \sum_{i=1}^n d_i(x_0)$$

so $\exists i$ such that $d(x_0, X \setminus U_i) = d_i(x_0) > \delta$. Thus $B_\delta(x_0)$ is disjoint from $X \setminus U_i$ and

$$Y \subset B_\delta(x_0) \subset U_i$$

as desired. □

APPENDIX D

Semicontinuous functions

D.1. Definitions and elementary properties

Let X be a topological space and $X \xrightarrow{f} \mathbb{R}$ a function. Then f is *upper semicontinuous* if it satisfies any of the following equivalent conditions:

- For each $x \in X$, f is *upper semicontinuous at x* , that is, for all $\epsilon > 0$,

$$f(y) < f(x) + \epsilon$$

for y in some neighborhood of x .

- f is a continuous mapping from X to \mathbb{R} , where \mathbb{R} is given the topology whose nonempty open sets are intervals $(-\infty, a)$ where $a \in \mathbb{R}$.

Examples of upper semicontinuous functions include the indicator function of a closed set, or the *greatest integer* (or *floor*) function. A function f is *lower semicontinuous* if and only if $-f$ is upper semicontinuous.

EXERCISE D.1.1. *Show that the following conditions are equivalent:*

- f is *lower semicontinuous*.
- $\forall x \in X$ and $\epsilon > 0$,

$$f(y) > f(x) - \epsilon$$

for y in an open neighborhood of x .

- $X \xrightarrow{f} \mathbb{R}$ is continuous where \mathbb{R} is given the topology generated by infinite open intervals (a, ∞) , for $a \in \mathbb{R}$.

EXERCISE D.1.2. *A semicontinuous function on a smooth manifold is Borel. (Hint: see Rudin [243], Theorem 1.12(c))*

D.2. Approximation by continuous functions

Let (X, d) be a metric space, $n > 0$ and $X \xrightarrow{f} \mathbb{R}$ any function.

PROPOSITION D.2.1. *Let $n \geq 0$. The function*

$$X \xrightarrow{h_n} \mathbb{R}$$

$$x \longmapsto \sup \{ f(p) - n \mathbf{d}(p, x) \mid p \in X \}$$

is n -Lipschitz.

PROOF. For any $p, y \in X$,

$$f(p) - n \mathbf{d}(p, y) \leq \sup \{ f(q) - n \mathbf{d}(q, y) \mid q \in X \} = h_n(y)$$

so

$$f(p) \leq h_n(y) + n \mathbf{d}(p, y) \leq h_n(y) + n(\mathbf{d}(p, x) + \mathbf{d}(x, y))$$

whence

$$f(p) - n \mathbf{d}(p, x) \leq h_n(y) + n \mathbf{d}(x, y).$$

Taking the supremum over p ,

$$h_n(x) \leq h_n(y) + n \mathbf{d}(x, y)$$

and

$$h_n(x) - h_n(y) \leq n \mathbf{d}(x, y).$$

Symmetrizing, the result follows. \square

LEMMA D.2.2. *If $m < n$, then $h_m(x) \geq h_n(x)$.*

PROOF. Taking the supremum over p of

$$f(p) - m \mathbf{d}(p, x) \leq f(p) - n \mathbf{d}(p, x),$$

the result follows. \square

Taking $p = x$ in the definition of h_n yields $h_n(x) \geq f(x)$. We now prove that a bounded upper semicontinuous function is the pointwise limit of a monotonically nonincreasing sequence of continuous (in fact Lipschitz) functions:

$$h_n(x) \searrow f(x).$$

EXERCISE D.2.3. *Let f be an upper semicontinuous function on a compact space X . Then f is bounded above, that is, $\exists M < \infty$ such that $f(x) \leq M$ for all $x \in X$.*

PROPOSITION D.2.4. *Suppose that f is an upper semicontinuous function which is bounded above. Let $x \in X$. Then*

$$\lim_{n \rightarrow \infty} h_n(x) = f(x).$$

PROOF. Suppose that $M < \infty$ and $f(y) < M$ for all $y \in X$. Let $\epsilon > 0$. Then it suffices to prove:

$$(108) \quad h_n(x) \leq f(x) + \epsilon$$

for sufficiently large n (depending on x , ϵ and M).

Since f is upper semicontinuous, $\exists \delta > 0$ such that

$$(109) \quad f(y) < f(x) + \epsilon$$

whenever $d(x, y) < \delta$. We claim:

$$(110) \quad f(y) - n d(y, x) < f(x) + \epsilon$$

whenever

$$(111) \quad n > \frac{M - f(x)}{\delta}.$$

If $d(x, y) < \delta$, then (109) implies:

$$f(y) - n d(y, x) < f(y) < f(x) + \epsilon.$$

Otherwise $d(x, y) \geq \delta$ and (111) implies:

$$f(y) - n d(y, x) < M - n\delta \leq f(x) < f(x) + \epsilon,$$

as claimed, proving (110). Taking supremum yields (108), completing the proof of Proposition D.2.4. \square

EXERCISE D.2.5. Let $X = \mathbb{R}$ and f be the indicator function at $0 \in \mathbb{R}$, that is,

$$f(x) := \begin{cases} 1 & \text{if } x = 0 \\ 0 & \text{if } x \neq 0 \end{cases}$$

Then

$$h_n(x) = \max(1 - nx, 0)$$

which is supported on the interval $[-1/n, 1/n]$.




FIGURE D.1. Continuous approximation of indicator function

APPENDIX E

SL(2, \mathbb{C}) and O(3, 1)

We prove the local isomorphism

$$\mathrm{SL}(2, \mathbb{C}) \longrightarrow \mathrm{O}(3, 1)$$

mentioned in §3.3.2. I am grateful to D. Sullivan for raising this question and motivating this material.

Let \mathbf{V} be a two-dimensional vector space over \mathbb{C} , and let $\mathbf{V} \times \mathbf{V} \xrightarrow{\Omega} \mathbb{C}$ be a nonzero (and hence nondegenerate) symplectic structure (that is, a skew-symmetric \mathbb{C} -bilinear form). Then $\mathrm{SL}(2, \mathbb{C})$ equals the automorphism group $\mathrm{Aut}(\mathbf{V}, \Omega)$.

Let $\mathbf{V}_{\mathbb{R}}$ be the underlying real vector space. Then \mathbf{V} corresponds to the pair $(\mathbf{V}_{\mathbb{R}}, \mathbf{J})$ where $\mathbf{V}_{\mathbb{R}} \xrightarrow{\mathbf{J}} \mathbf{V}_{\mathbb{R}}$ is the complex structure.

In terms of $(\mathbf{V}_{\mathbb{R}}, \mathbf{J})$, the \mathbb{C} -symplectic structure Ω on \mathbf{V} is equivalent to a pair (ω, ψ) of symplectic structures $\mathbf{V}_{\mathbb{R}} \times \mathbf{V}_{\mathbb{R}} \rightarrow \mathbb{R}$ on $\mathbf{V}_{\mathbb{R}}$ which are the real and imaginary parts of Ω :

$$\Omega(x, y) = \omega(x, y) + i\psi(x, y)$$

Both ω and ψ are compatible with \mathbf{J} in the following sense:

$$(112) \quad \omega(\mathbf{J}x, \mathbf{J}y) = -\omega(x, y)$$

and

$$\psi(\mathbf{J}x, \mathbf{J}y) = -\psi(x, y).$$

Furthermore \mathbf{J} and ω determine ψ by:

$$(113) \quad \psi(x, y) = -\omega(\mathbf{J}x, y).$$

Thus the pair (\mathbf{V}, Ω) is equivalent to the pair $(\mathbf{V}_{\mathbb{R}}, \mathbf{J}, \omega)$ satisfying (112). Furthermore $\mathrm{SL}(2, \mathbb{C})$ equals the automorphism group $\mathrm{Aut}(\mathbf{V}, \mathbf{J}, \omega)$.

Choose a nonzero element $\mu \in \Lambda^4(\mathbf{V}_{\mathbb{R}})$ of $\mathbf{V}_{\mathbb{R}}$. The second exterior power $\mathbf{W} := \Lambda^2(\mathbf{V}_{\mathbb{R}})$ of $\mathbf{V}_{\mathbb{R}}$ has dimension 6 and admits a nondegenerate symmetric bilinear form

$$\begin{aligned} \Lambda^2(\mathbf{V}_{\mathbb{R}}) \times \Lambda^2(\mathbf{V}_{\mathbb{R}}) &\longrightarrow \mathbb{R} \\ (x, y) &\longmapsto x \cdot y \end{aligned}$$

defined by:

$$x \wedge y = (x \cdot y)\mu.$$

If $g \in \mathbf{GL}(\mathbf{V}_{\mathbb{R}})$, then

$$g(x) \cdot g(y) = \det(g) x \cdot y$$

applied to an orientation-reversing element of $\mathbf{GL}(\mathbf{V}_{\mathbb{R}})$ implies this form is equivalent to its negative and therefore has signature $(3, 3)$. This defines a local isomorphism $\mathbf{SL}(4, \mathbb{R}) \rightarrow \mathbf{O}(3, 3)$.

Now introduce the first symplectic structure $\omega \in \Lambda^2(\mathbf{V}_{\mathbb{R}}^*)$. If $g \in \mathbf{Aut}(\mathbf{V}_{\mathbb{R}})$ stabilizes ω , it stabilizes the dual bivector $\omega^* \in \Lambda^2(\mathbf{V}_{\mathbb{R}}) = \mathbf{W}$. Choosing μ so that $(\omega \wedge \omega)(\mu) < 0$ implies that $\omega^* \cdot \omega^* < 0$. It follows that the orthogonal complement

$$\mathbf{W}_1 := (\omega^*)^\perp \subset \mathbf{W}$$

is a nondegenerate subspace having signature $(3, 2)$. In particular the stabilizer $\mathbf{Aut}(\mathbf{V}_{\mathbb{R}}, \omega) \cong \mathbf{Sp}(4, \mathbb{R})$ preserves \mathbf{W}_1 , defining a local isomorphism $\mathbf{Sp}(4, \mathbb{R}) \rightarrow \mathbf{O}(3, 2)$.

Similarly, the second symplectic structure $\psi \in \Lambda^2(\mathbf{V}_{\mathbb{R}}^*)$ admits a dual bivector $\psi^* \in \Lambda^2(\mathbf{V}_{\mathbb{R}}) = \mathbf{W}$ orthogonal to ω^* also satisfying $\psi^* \cdot \psi^* < 0$. The orthogonal complement $\mathbf{W}_2 := (\omega^*, \psi^*)^\perp \subset \mathbf{W}$ has signature $(3, 1)$.

By (113), the complex structure \mathbf{J} and the symplectic structure ω determine the symplectic structure ψ , so

$$\mathbf{Aut}(\mathbf{V}_{\mathbb{R}}, \omega, \psi) = \mathbf{Aut}(\mathbf{V}, \Omega) \cong \mathbf{SL}(2, \mathbb{C}).$$

The stabilizer $\mathbf{Aut}(\mathbf{V}_{\mathbb{R}}, \omega, \psi)$ preserves \mathbf{W}_2 , defining a local isomorphism $\mathbf{SL}(2, \mathbb{C}) \rightarrow \mathbf{O}(3, 1)$.

Explicitly, choose a basis $\mathbf{e}_1, \mathbf{e}_2$ of \mathbf{V} with $\Omega(\mathbf{e}_1, \mathbf{e}_2) = 1$. Extend to a basis $\mathbf{e}_1, \mathbf{f}_1, \mathbf{e}_2, \mathbf{f}_2$ of $\mathbf{V}_{\mathbb{R}}$ by $\mathbf{f}_j := \mathbf{J}\mathbf{e}_j$. Evaluate ω, ψ on this basis:

$$\omega(\mathbf{e}_1, \mathbf{e}_2) = -\omega(\mathbf{f}_1, \mathbf{f}_2) = 1 \quad \psi(\mathbf{f}_1, \mathbf{e}_2) = -\psi(\mathbf{f}_2, \mathbf{e}_1) = 1$$

(with other values 0) so that ¹

$$\omega^* = \frac{1}{4}\mathbf{e}_1 \wedge \mathbf{e}_2 - \mathbf{f}_1 \wedge \mathbf{f}_2, \quad \psi^* = \frac{1}{4}\mathbf{f}_1 \wedge \mathbf{e}_2 - \mathbf{f}_2 \wedge \mathbf{e}_1.$$

Let $\mu := \mathbf{e}_1 \wedge \mathbf{e}_2 \wedge \mathbf{f}_1 \wedge \mathbf{f}_2$. Then $\omega^* \cdot \omega^* = \psi^* \cdot \psi^* = -1/8$, $\omega^* \cdot \psi^* = 0$. Then $\mathbf{e}_1 \wedge \mathbf{f}_1, \mathbf{e}_2 \wedge \mathbf{f}_2, \mathbf{e}_1 \wedge \mathbf{e}_2 + \mathbf{f}_1 \wedge \mathbf{f}_2, \mathbf{e}_1 \wedge \mathbf{f}_2 - \mathbf{f}_1 \wedge \mathbf{e}_2$ bases \mathbf{W}_2 with Gram matrix (evidently of signature $(3, 1)$):

$$\begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}.$$

¹Convention: $(\alpha \wedge \beta)(X_1, \dots, X_{p+q}) = \frac{1}{p!q!} \sum_{\sigma \in \mathfrak{S}_{p+q}} \alpha(X_{\sigma(1)}, \dots, X_{\sigma(p)}) \beta(X_{\sigma(p+1)}, \dots, X_{\sigma(p+q)})$

Bibliography

1. Herbert Abels, *Properly discontinuous groups of affine transformations: a survey*, Geom. Dedicata **87** (2001), no. 1-3, 309–333. MR 1866854 [15.1](#), [15.3.2](#)
2. William Abikoff, *The real analytic theory of Teichmüller space*, Lecture Notes in Mathematics, vol. 820, Springer, Berlin, 1980. MR 590044 [7.4](#)
3. Lars V. Ahlfors, *Complex analysis*, third ed., McGraw-Hill Book Co., New York, 1978, An introduction to the theory of analytic functions of one complex variable, International Series in Pure and Applied Mathematics. MR 510197 [1](#)
4. Raphaël Alexandre, *Fried's theorem for boundary geometries of rank one symmetric spaces*, Preprint, hal-02293855. [11.4](#)
5. Sasha Ananin, Carlos H. Grossi, and Nikolay Gusevskii, *Complex hyperbolic structures on disc bundles over surfaces*, Int. Math. Res. Not. IMRN (2011), no. 19, 4295–4375. MR 2838042 [15.1](#)
6. D. K. Arrowsmith and P. M. D. Furness, *Locally symmetric spaces*, J. London Math. Soc. (2) **10** (1975), no. 4, 487–499. MR 0467598 (57 #7454) [10.2](#)
7. ———, *Flat affine Klein bottles*, Geometriae Dedicata **5** (1976), no. 1, 109–115. MR 0433358 (55 #6334) [10.2](#)
8. L. Auslander and L. Markus, *Holonomy of flat affinely connected manifolds*, Ann. of Math. (2) **62** (1955), 139–151. MR 0072518 (17,298b) [8.3.2](#), [8.3.9](#)
9. ———, *Flat Lorentz 3-manifolds*, Mem. Amer. Math. Soc. No. **30** (1959), 60. MR 0131842 (24 #A1689) [6.5.2.3](#), [8.6.2](#), [10.7](#)
10. Louis Auslander, *The structure of complete locally affine manifolds*, Topology **3** (1964), no. suppl. 1, 131–139. MR 0161255 [8.6](#), [8.6.2](#), [9.2.3](#)
11. ———, *Simply transitive groups of affine motions*, Amer. J. Math. **99** (1977), no. 4, 809–826. MR 447470 [10.5.1](#), [10.6](#), [10.7](#)

12. Shinpei Baba, *A Schottky decomposition theorem for complex projective structures*, *Geom. Topol.* **14** (2010), no. 1, 117–151. MR 2578302 [14](#)
13. ———, *Complex projective structures with Schottky holonomy*, *Geom. Funct. Anal.* **22** (2012), no. 2, 267–310. MR 2929066 [14](#)
14. ———, *2π -grafting and complex projective structures, I*, *Geom. Topol.* **19** (2015), no. 6, 3233–3287. MR 3447103 [14](#)
15. ———, *2π -grafting and complex projective structures with generic holonomy*, *Geom. Funct. Anal.* **27** (2017), no. 5, 1017–1069. MR 3714716 [14](#)
16. Shinpei Baba and Subhojoy Gupta, *Holonomy map fibers of $\mathbb{C}P^1$ -structures in moduli space*, *J. Topol.* **8** (2015), no. 3, 691–710. MR 3394314 [14](#)
17. Samuel A. Ballas, Daryl Cooper, and Arielle Leitner, *Generalized cusps in real projective manifolds: classification*, *J. Topol.* **13** (2020), no. 4, 1455–1496. MR 4125754 [4](#)
18. Samuel A. Ballas, Jeffrey Danciger, and Gye-Seon Lee, *Convex projective structures on nonhyperbolic three-manifolds*, *Geom. Topol.* **22** (2018), no. 3, 1593–1646. MR 3780442 [4](#)
19. Thierry Barbot, *Variétés affines radiales de dimension 3*, *Bull. Soc. Math. France* **128** (2000), no. 3, 347–389. MR 1792474 [6.5.2.3](#)
20. Thierry Barbot, Virginie Charette, Todd Drumm, William M. Goldman, and Karin Melnick, *A primer on the $(2 + 1)$ Einstein universe*, *Recent developments in pseudo-Riemannian geometry*, *ESI Lect. Math. Phys.*, Eur. Math. Soc., Zürich, 2008, pp. 179–229. MR 2436232 [15.1](#)
21. Oliver Baues, *Varieties of discontinuous groups*, in Igodt et al. [[156](#)], pp. 147–158. MR 1796130 (2001i:58013) [7.2](#)
22. ———, *Deformation spaces for affine crystallographic groups*, *Cohomology of groups and algebraic K -theory*, *Adv. Lect. Math. (ALM)*, vol. 12, Int. Press, Somerville, MA, 2010, pp. 55–129. MR 2655175 (2011f:57066) [8.5](#), [8.5.1](#)
23. ———, *The deformations of flat affine structures on the two-torus*, *Handbook of Teichmüller theory Vol. IV*, *European Mathematical Society*, 2014, pp. 461–537. [5.4](#), [5.4.4](#), [8.5](#), [10](#), [10.2](#)
24. Yves Benoist, *Une nilvariété non affine*, *C. R. Acad. Sci. Paris Sér. I Math.* **315** (1992), no. 9, 983–986. MR 1186933 [15.2](#)
25. ———, *Nilvariétés projectives*, *Comment. Math. Helv.* **69** (1994), no. 3, 447–473. MR 1289337 [11.2](#)
26. ———, *Une nilvariété non affine*, *J. Differential Geom.* **41** (1995), no. 1, 21–52. MR 1316552 [15.2](#)

27. ———, *Automorphismes des cônes convexes*, *Invent. Math.* **141** (2000), no. 1, 149–193. MR 1767272 (2001f:22034) [4.6](#)
28. ———, *Tores affines*, *Crystallographic groups and their generalizations* (Kortrijk, 1999), *Contemp. Math.*, vol. 262, Amer. Math. Soc., Providence, RI, 2000, pp. 1–37. MR 1796124 (2001k:57025) [10.2](#), [11.2](#)
29. ———, *Convexes divisibles. II*, *Duke Math. J.* **120** (2003), no. 1, 97–120. MR 2010735 (2004m:22018) [4.6](#)
30. ———, *Convexes divisibles. I*, *Algebraic groups and arithmetic*, *Tata Inst. Fund. Res.*, Mumbai, 2004, pp. 339–374. MR 2094116 (2005h:37073) [4.6](#)
31. ———, *Convexes divisibles. III*, *Ann. Sci. École Norm. Sup. (4)* **38** (2005), no. 5, 793–832. MR 2195260 (2007b:22011) [4.6](#)
32. ———, *Convexes divisibles. IV. Structure du bord en dimension 3*, *Invent. Math.* **164** (2006), no. 2, 249–278. MR 2218481 (2007g:22007) [4.6](#)
33. ———, *A survey on divisible convex sets*, *Geometry, analysis and topology of discrete groups*, *Adv. Lect. Math. (ALM)*, vol. 6, Int. Press, Somerville, MA, 2008, pp. 1–18. MR 2464391 (2010h:52013) [4.6](#)
34. Jean-Paul Benzécri, *Variétés localement affines*, *Seminaire de Topologie et Géom. Diff.*, Ch. Ehresmann (1958–60) (1959), no. 7, 1–34. ([document](#)), [9.1](#), [13.2](#), [13.2](#)
35. ———, *Sur les variétés localement affines et localement projectives*, *Bulletin de la Société Mathématique de France* **88** (1960), 229–332. [4](#), [4.5.2.3](#), [6.1.2](#), [6.4.3](#), [6.4.6](#), [6.4.7](#)
36. ———, *Sur la classe d'Euler (ou Stiefel-Whitney) de fibrés affins plats*, *C. R. Acad. Sci. Paris* **260** (1965), 5442–5444. MR 0178477 (31 #2734) [9.1](#), [9.1.3](#)
37. Marcel Berger, *Geometry. I*, *Universitext*, Springer-Verlag, Berlin, 1987, Translated from the French by M. Cole and S. Levy. MR 882541 (88a:51001a) [2](#), [4.1.1](#)
38. ———, *Geometry. II*, *Universitext*, Springer-Verlag, Berlin, 1987, Translated from the French by M. Cole and S. Levy. MR 882916 ([document](#)), [2](#), [2.5.2](#)
39. ———, *Geometry I*, *Universitext*, Springer-Verlag, Berlin, 2009, Translated from the 1977 French original by M. Cole and S. Levy, Fourth printing of the 1987 English translation [MR0882541]. MR 2724360 ([document](#)), [2.5.2](#)

40. Marcel Berger, Pierre Pansu, Jean-Pic Berry, and Xavier Saint Raymond, *Problems in geometry*, Problem Books in Mathematics, Springer-Verlag, New York, 1984, Translated from the French by Silvio Levy. MR 772926 ([document](#))
41. N. Bergeron and T. Gelander, *A note on local rigidity*, *Geom. Dedicata* **107** (2004), 111–131. MR 2110758 (2005k:22015) [7.2.1](#)
42. Francis Bonahon, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, *Ann. Fac. Sci. Toulouse Math.* (6) **5** (1996), no. 2, 233–297. MR 1413855 [13.1](#)
43. ———, *Geometric structures on 3-manifolds*, Handbook of geometric topology, North-Holland, Amsterdam, 2002, pp. 93–164. MR 590044 ([document](#))
44. Francis Bonahon and Guillaume Dreyer, *Parameterizing Hitchin components*, *Duke Math. J.* **163** (2014), no. 15, 2935–2975. MR 3285861 [13.2.1](#), [13.2.3](#)
45. Francis Bonahon and In Kang Kim, *The Goldman and Fock-Goncharov coordinates for convex projective structures on surfaces*, *Geom. Dedicata* **192** (2018), 43–55. MR 3749422 [13.1](#), [13.2.1](#)
46. Raoul Bott and Loring W. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, vol. 82, Springer-Verlag, New York-Berlin, 1982. MR 658304 [9.2.1](#)
47. Robert Brody, *Compact manifolds and hyperbolicity*, *Trans. Amer. Math. Soc.* **235** (1978), 213–219. MR 470252 [12.2.3](#)
48. Michelle Bucher and Tsachik Gelander, *The generalized Chern conjecture for manifolds that are locally a product of surfaces*, *Advances in Math.* **228** (2011), 1503–1542. [9.1.3](#)
49. Dietrich Burde, *Simple left-symmetric algebras with solvable Lie algebra*, *Manuscripta Math.* **95** (1998), no. 3, 397–411. MR 1612015 [10.4](#), [10.4.4](#), [10.4.4.2](#)
50. ———, *Left-symmetric algebras, or pre-Lie algebras in geometry and physics*, *Cent. Eur. J. Math.* **4** (2006), no. 3, 323–357 (electronic). MR 2233854 (2007c:17001) [10](#), [10.3.1](#), [10.4.4.2](#)
51. Jean-Philippe Burrelle, Virginie Charette, Todd A. Drumm, and William M. Goldman, *Crooked halfspaces*, *Enseign. Math.* **60** (2014), no. 1-2, 43–78. MR 3262435 [15.1](#)
52. Marc Burger, Alessandra Iozzi, and Anna Wienhard, *Higher Teichmüller spaces: from $SL(2, \mathbb{R})$ to other Lie groups*, Handbook of Teichmüller theory. Vol. IV, IRMA Lect. Math. Theor. Phys., vol. 19, Eur. Math. Soc., Zürich, 2014, pp. 539–618. MR 3289711 [9.1.3](#)

53. D. Burns, Jr. and S. Shnider, *Spherical hypersurfaces in complex manifolds*, Invent. Math. **33** (1976), no. 3, 223–246. MR 419857 [15.1](#)
54. Herbert Busemann and Paul J. Kelly, *Projective geometry and projective metrics*, Academic Press Inc., New York, N. Y., 1953. MR 0054980 (14,1008e) [2](#), [2.5.2](#), [3.2.6](#), [3.2.2](#)
55. R. D. Canary, D. B. A. Epstein, and P. Green, *Notes on notes of Thurston*, Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984), London Math. Soc. Lecture Note Ser., vol. 111, Cambridge Univ. Press, Cambridge, 1987, pp. 3–92. MR 903850 [7.2.1](#)
56. Yves Carrière, *Autour de la conjecture de L. Markus sur les variétés affines*, Invent. Math. **95** (1989), no. 3, 615–628. MR 979369 ([document](#))
57. Yves Carrière, Françoise Dal’bo, and Gaël Meigniez, *Inexistence de structures affines sur les fibrés de Seifert*, Math. Ann. **296** (1993), no. 4, 743–753. MR 1233496 (94h:57031) [11.3](#)
58. Alex Casella, Dominic Tate, and Stephan Tillmann, *Moduli spaces of real projective structures on surfaces*, MSJ Memoirs, vol. 38, Mathematical Society of Japan, Tokyo, [2020] ©2020. MR 4242841 [4.5.3](#), [13](#)
59. Virginie Charette and Todd A. Drumm, *The Margulis invariant for parabolic transformations*, Proc. Amer. Math. Soc. **133** (2005), no. 8, 2439–2447. MR 2138887 [15.3.2](#)
60. Virginie Charette, Todd A. Drumm, and William M. Goldman, *Affine deformations of a three-holed sphere*, Geom. Topol. **14** (2010), no. 3, 1355–1382. MR 2653729 [15.1](#)
61. ———, *Finite-sided deformation spaces of complete affine 3-manifolds*, J. Topol. **7** (2014), no. 1, 225–246. MR 3180618 [15.1](#)
62. ———, *Proper affine deformations of the one-holed torus*, Transform. Groups **21** (2016), no. 4, 953–1002. MR 3569564 [15.1](#)
63. Leonard S. Charlap, *Bieberbach groups and flat manifolds*, Universitext, Springer-Verlag, New York, 1986. MR 862114 [6](#), [6.2.1](#), [8.6](#)
64. S. S. Chern and Phillip Griffiths, *Abel’s theorem and webs*, Jahresber. Deutsch. Math.-Verein. **80** (1978), no. 1-2, 13–110. MR 494957 (80b:53008) [5](#)
65. Shiing-shen Chern, *A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds*, Ann. of Math. (2) **45** (1944), 747–752. MR 0011027 (6,106a) [9.2.1](#)

66. Suhyoung Choi, *Convex decompositions of real projective surfaces. I. π -annuli and convexity*, J. Differential Geom. **40** (1994), no. 1, 165–208. MR 1285533 (95i:57015) [13.1](#)
67. ———, *Convex decompositions of real projective surfaces. III. For closed or nonorientable surfaces*, J. Korean Math. Soc. **33** (1996), no. 4, 1139–1171. MR 1424213 [13.1](#)
68. ———, *The convex and concave decomposition of manifolds with real projective structures*, Mém. Soc. Math. Fr. (N.S.) (1999), no. 78, vi+102. MR 1779499 (2001j:57030) [13.1](#)
69. ———, *The decomposition and classification of radiant affine 3-manifolds*, Mem. Amer. Math. Soc. **154** (2001), no. 730, viii+122. MR 1848866 (2002f:57049) [6.5.2.3](#)
70. Suhyoung Choi and William M. Goldman, *The classification of real projective structures on compact surfaces*, Bull. Amer. Math. Soc. (N.S.) **34** (1997), no. 2, 161–171. MR 1414974 (97m:57020) [13.1](#)
71. ———, *The deformation spaces of convex \mathbb{RP}^2 -structures on 2-orbifolds*, Amer. J. Math. **127** (2005), no. 5, 1019–1102. MR 2170138 (2007k:57031) [13.1](#), [13.2.1](#)
72. Suhyoung Choi, Craig D. Hodgson, and Gye-Seon Lee, *Projective deformations of hyperbolic Coxeter 3-orbifolds*, Geom. Dedicata **159** (2012), 125–167. MR 2944525 [4.6](#)
73. Suhyoung Choi, Hongtaek Jung, and Hong Chan Kim, *Symplectic coordinates on $\mathrm{PSL}_3(\mathbb{R})$ -Hitchin components*, Pure Appl. Math. Q. **16** (2020), no. 5, 1321–1386. MR 4220999 [13.1](#)
74. Suhyoung Choi, Gye-Seon Lee, and Ludovic Marquis, *Deformations of convex real projective manifolds and orbifolds*, Handbook of group actions. Vol. III, Adv. Lect. Math. (ALM), vol. 40, Int. Press, Somerville, MA, 2018, pp. 263–310. MR 3888622 [4](#)
75. Suhyoung Choi and Hyunkoo Lee, *Geometric structures on manifolds and holonomy-invariant metrics*, Forum Math. **9** (1997), no. 2, 247–256. MR 1431123 (98a:53052) [14.2](#)
76. Bruno Colbois and Patrick Verovic, *Rigidity of Hilbert metrics*, Bull. Austral. Math. Soc. **65** (2002), no. 1, 23–34. MR 1889375 [4.6](#)
77. Daryl Cooper and William M. Goldman, *A 3-manifold with no real projective structure*, Ann. Fac. Sci. Toulouse Math. (6) **24** (2015), no. 5, 1219–1238. MR 3485333 [15.1](#)
78. Daryl Cooper, Darren Long, and Stephan Tillmann, *On convex projective manifolds and cusps*, Adv. Math. **277** (2015), 181–251. MR 3336086 ([document](#)), [4](#), [4.5.3](#), [4.5.3](#)

79. ———, *Deforming convex projective manifolds*, *Geom. Topol.* **22** (2018), no. 3, 1349–1404. MR 3780436 [4](#), [4.5.3](#), [12.3](#)
80. H. S. M. Coxeter, *Introduction to geometry*, second ed., John Wiley & Sons Inc., New York, 1969. MR 0346644 (49 #11369) ([document](#)), [2](#), [2.5.2](#), [3](#), [3.2.4](#)
81. ———, *Projective geometry*, second ed., University of Toronto Press, Toronto, Ont., 1974. MR 0346652 (49 #11377) [2](#), [2.5.2](#)
82. ———, *Non-Euclidean geometry*, sixth ed., MAA Spectrum, Mathematical Association of America, Washington, DC, 1998. MR 1628013 [1](#), [3.2.2](#)
83. Jeffrey Danciger, Todd A. Drumm, William M. Goldman, and Iliia Smilga, *Proper Actions of Discrete Groups of Affine Transformations*, *Dynamics, Geometry, Number Theory: the impact of Margulis on Modern Mathematics*, Proceedings, D. Fisher, D. Kleinbock, and G. Soifer, (eds.), University of Chicago Press, 2021. [15.1](#), [15.2](#)
84. Jeffrey Danciger, François Guéritaud, and Fanny Kassel, *Geometry and topology of complete Lorentz spacetimes of constant curvature*, *Ann. Sci. Éc. Norm. Supér. (4)* **49** (2016), no. 1, 1–56. MR 3465975 [15.1](#)
85. ———, *Margulis spacetimes via the arc complex*, *Invent. Math.* **204** (2016), no. 1, 133–193. MR 3480555 [15.1](#)
86. Pierre Deligne and Dennis Sullivan, *Fibrés vectoriels complexes à groupe structural discret*, *C. R. Acad. Sci. Paris Sér. A-B* **281** (1975), no. 24, Ai, A1081–A1083. MR 0397729 (53 #1587) [9.2](#)
87. Manfredo Perdigão do Carmo, *Riemannian geometry*, *Mathematics: Theory & Applications*, Birkhäuser Boston, Inc., Boston, MA, 1992, Translated from the second Portuguese edition by Francis Flaherty. MR 1138207 ([document](#)), [8.1.1](#), [B.1](#)
88. J. Dorfmeister, *Quasiclans*, *Abh. Math. Sem. Univ. Hamburg* **50** (1980), 178–187. MR 593749 [10](#)
89. Todd A. Drumm, *Fundamental polyhedra for Margulis spacetimes*, ProQuest LLC, Ann Arbor, MI, 1990, Thesis (Ph.D.)–University of Maryland, College Park. MR 2638637 [15.3.1](#)
90. ———, *Linear holonomy of Margulis space-times*, *J. Differential Geom.* **38** (1993), no. 3, 679–690. MR 1243791 [15.3.1](#)
91. David Dumas, *Complex projective structures*, *Handbook of Teichmüller theory. Vol. II*, IRMA Lect. Math. Theor. Phys., vol. 13, Eur. Math. Soc., Zürich, 2009, pp. 455–508. MR 2497780 [7.2.1](#), [14](#), [15.1](#)

92. Sorin Dumitrescu, *Structures géométriques sur les courbes et les surfaces complexes*, Ann. Fac. Sci. Toulouse Math. (6) **10** (2001), no. 3, 507–531. MR 1923688 [14](#)
93. Serge Dupont, *Variétés projectives à holonomie dans le groupe $\text{Aff}^+(\mathbf{R})$* , in Igodt et al. [[156](#)], pp. 177–193. MR 1796133 [14.2](#), [15.4](#)
94. ———, *Solvariétés projectives de dimension trois*, Geom. Dedicata **96** (2003), 55–89. MR 1956834 [10](#), [15.1](#), [15.4.1](#), [15.4](#)
95. Clifford J. Earle, *On variation of projective structures*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 87–99. MR 624807 [7.2.1](#)
96. Charles Ehresmann, *Sur les espaces localement homogènes*, L'Ens. Math **35** (1936), 317–333. ([document](#)), [5.2.1](#), [13.2](#)
97. ———, *Les connexions infinitésimales dans un espace fibré différentiable*, in Thone [[263](#)], pp. 29–55. MR 0042768 [7.2.1](#)
98. Y. Eliashberg and N. Mishachev, *Introduction to the h-principle*, Graduate Studies in Mathematics, vol. 48, American Mathematical Society, Providence, RI, 2002. MR 1909245 [6.1.1](#), [7.5](#)
99. Gerd Faltings, *Real projective structures on Riemann surfaces*, Compositio Math. **48** (1983), no. 2, 223–269. MR 700005 [14.2](#)
100. Benson Farb and Dan Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, Princeton, NJ, 2012. MR 2850125 [7.4](#)
101. Vladimir Fock and Alexander Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. (2006), no. 103, 1–211. MR 2233852 [13.1](#), [13.3.1](#)
102. ———, *Moduli spaces of convex projective structures on surfaces*, Adv. Math. **208** (2007), no. 1, 249–273. MR 2304317 [13.1](#), [13.3.1](#)
103. Ralph H. Fox, *Free differential calculus. I. Derivation in the free group ring*, Ann. of Math. (2) **57** (1953), 547–560. MR 0053938 [7.3](#)
104. Robert Fricke and Felix Klein, *Vorlesungen über die Theorie der automorphen Funktionen. Band 1: Die gruppentheoretischen Grundlagen. Band II: Die funktionentheoretischen Ausführungen und die Anwendungen*, Bibliotheca Mathematica Teubneriana, Bände 3, vol. 4, Johnson Reprint Corp., New York; B. G. Teubner Verlagsgesellschaft, Stuttgart art, 1965. MR 0183872 (32 #1348) [7.3.1](#), [7.3.6](#)

105. David Fried, *Radiant flows without cross-sections*, Preprint. [6.5.2.3](#)
106. ———, *Closed similarity manifolds*, Comment. Math. Helv. **55** (1980), no. 4, 576–582. MR 604714 (83e:53049) ([document](#)), [11.4](#), [11.4.6](#)
107. ———, *The geometry of cross sections to flows*, Topology **21** (1982), no. 4, 353–371. MR 670741 (84d:58068) [6.1.4](#)
108. ———, *Distality, completeness, and affine structures*, J. Differential Geom. **24** (1986), no. 3, 265–273. MR 868973 (88i:53060) [6.5.2.3](#), [8.4](#), [8.4](#), [10.5.1](#), [10.6](#)
109. ———, *Affine manifolds that fiber by circles*, IHES Preprint IHES/M/92/56, (1992). [6.5.2.3](#)
110. David Fried and William M. Goldman, *Three-dimensional affine crystallographic groups*, Adv. in Math. **47** (1983), no. 1, 1–49. MR 689763 [8.5](#), [8.6](#), [8.6.2](#), [8.6.2](#), [8.6.2](#), [10.5](#), [10.5.3](#), [10.7](#), [15.1](#), [15.2](#)
111. David Fried, William M. Goldman, and Morris W. Hirsch, *Affine manifolds with nilpotent holonomy*, Comment. Math. Helv. **56** (1981), no. 4, 487–523. MR 656210 (83h:53062) [7.5](#), [8.4.1](#), [8.4](#), [11.1](#), [11.2](#), [11.2](#), [11.2.3](#), [11.2](#)
112. Daniel Gallo, Michael Kapovich, and Albert Marden, *The monodromy groups of Schwarzian equations on closed Riemann surfaces*, Ann. of Math. (2) **151** (2000), no. 2, 625–704. MR 1765706 [14](#)
113. W Goldman and Yoshinobu Kamishima, *Topological rigidity of developing maps with applications to conformally flat structures*, Contemp. Math **74** (1988), 199–203. [14.2](#)
114. William M. Goldman, *Affine manifolds and projective geometry on surfaces*, senior thesis, Princeton University, 1977. ([document](#)), [10.2](#), [13](#)
115. ———, *Discontinuous groups and the Euler class*, Ph.D. thesis, University of California, Berkeley, 1980. [5.4](#), [9.1](#)
116. ———, *Two examples of affine manifolds*, Pacific J. Math. **94** (1981), no. 2, 327–330. MR 628585 (83a:57028) [10.8.1](#), [15.4](#)
117. ———, *Conformally flat manifolds with nilpotent holonomy and the uniformization problem for 3-manifolds*, Trans. Amer. Math. Soc. **278** (1983), no. 2, 573–583. MR 701512 [11.2](#), [15.1](#)
118. ———, *The symplectic nature of fundamental groups of surfaces*, Adv. in Math. **54** (1984), no. 2, 200–225. MR 762512 [7.4](#)
119. ———, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. Math. **85** (1986), no. 2, 263–302. MR 846929 [7.4](#), [7.4](#)

120. ———, *Projective structures with Fuchsian holonomy*, J. Differential Geom. **25** (1987), no. 3, 297–326. MR 882826 [5.4.5](#), [13.4.0.1](#), [13.4.0.1](#), [14](#), [14.2](#)
121. ———, *Geometric structures on manifolds and varieties of representations*, Geometry of group representations (Boulder, CO, 1987), Contemp. Math., vol. 74, Amer. Math. Soc., Providence, RI, 1988, pp. 169–198. MR 957518 [7.2.1](#), [15.1](#)
122. ———, *Convex real projective structures on compact surfaces*, J. Differential Geom **31** (1990), no. 3, 791–845. [13.1](#), [13.2.1](#)
123. William M Goldman, *The symplectic geometry of affine connections on surfaces*, Journal für die reine und angewandte Mathematik **407** (1990), 126–159. [13.1](#), [B.2](#)
124. William M. Goldman, *Complex hyperbolic geometry*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1999, Oxford Science Publications. MR 1695450 [1](#), [3.2.2](#), [15.1](#)
125. ———, *The Margulis invariant of isometric actions on Minkowski $(2 + 1)$ -space*, Rigidity in dynamics and geometry (Cambridge, 2000), Springer, Berlin, 2002, pp. 187–201. MR 1919401 [15.3.3](#)
126. ———, *Trace coordinates on Fricke spaces of some simple hyperbolic surfaces*, Handbook of Teichmüller theory. Vol. II, IRMA Lect. Math. Theor. Phys., vol. 13, Eur. Math. Soc., Zürich, 2009, pp. 611–684. MR 2497777 (2010j:30093) [7.3.1](#), [7.3.1.1](#), [7.3.1.2](#)
127. ———, *Locally homogeneous geometric manifolds*, Proceedings of the International Congress of Mathematicians. Volume II, Hindustan Book Agency, New Delhi, 2010, pp. 717–744. MR 2827816 [7.2](#), [15.1](#)
128. ———, *Two papers which changed my life: Milnor’s seminal work on flat manifolds and bundles*, Frontiers in complex dynamics: In celebration of John Milnor’s 80th birthday, Princeton Math. Ser., vol. 51, Princeton Univ. Press, Princeton, NJ, 2014, pp. 679–703. MR 3289925 [9.1](#), [15.1](#)
129. ———, *Flat affine, projective and conformal structures on manifolds: a historical perspective*, Geometry in history, Springer, Cham, 2019, pp. 515–552. MR 3965773 ([document](#))
130. ———, *Parallelism on Lie groups and Fox’s free differential calculus*, Characters in low-dimensional topology, Contemp. Math., vol. 760, Amer. Math. Soc., Providence, RI, 2020, pp. 157–166. MR 4193925 [7.3](#)

131. William M. Goldman and Morris W. Hirsch, *Flat bundles with solvable holonomy*, Proc. Amer. Math. Soc. **82** (1981), no. 3, 491–494. MR 612747 (82e:57010) [9.2](#)
132. ———, *The radiance obstruction and parallel forms on affine manifolds*, Trans. Amer. Math. Soc. **286** (1984), no. 2, 629–649. MR 760977 (86f:57032) [11.1](#), [11.3](#), [11.3](#)
133. ———, *Affine manifolds and orbits of algebraic groups*, Trans. Amer. Math. Soc. **295** (1986), no. 1, 175–198. MR 831195 (88c:57025) [10.3.2](#), [10.3.2](#), [10.3.4](#)
134. William M. Goldman and Gregory A. Margulis, *Flat Lorentz 3-manifolds and cocompact Fuchsian groups*, Crystallographic groups and their generalizations (Kortrijk, 1999), Contemp. Math., vol. 262, Amer. Math. Soc., Providence, RI, 2000, pp. 135–145. MR 1796129 [15.3](#), [15.3.3](#)
135. Walter Helbig Gottschalk and Gustav Arnold Hedlund, *Topological dynamics*, American Mathematical Society Colloquium Publications, Vol. 36, American Mathematical Society, Providence, R. I., 1955. MR 0074810 [8.6.2](#)
136. M. Gromov, H. B. Lawson, Jr., and W. Thurston, *Hyperbolic 4-manifolds and conformally flat 3-manifolds*, Inst. Hautes Études Sci. Publ. Math. (1988), no. 68, 27–45 (1989). MR 1001446 [15.1](#)
137. Michael Gromov, *Volume and bounded cohomology*, Inst. Hautes Études Sci. Publ. Math. (1982), no. 56, 5–99 (1983). MR 686042 (84h:53053) [9.1](#)
138. Mikhael Gromov, *Partial differential relations*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 9, Springer-Verlag, Berlin, 1986. MR 864505 [7.5](#)
139. François Guéritaud, *On Lorentz spacetimes of constant curvature*, Geometry, groups and dynamics, Contemp. Math., vol. 639, Amer. Math. Soc., Providence, RI, 2015, pp. 253–269. MR 3379833 [15.1](#)
140. Olivier Guichard, *Sur la régularité Hölder des convexes divisibles*, Ergodic Theory Dynam. Systems **25** (2005), no. 6, 1857–1880. MR 2183298 [13.2](#), [13.2.1](#)
141. Olivier Guichard and Anna Wienhard, *Convex foliated projective structures and the Hitchin component for $\mathrm{PSL}_4(\mathbf{R})$* , Duke Math. J. **144** (2008), no. 3, 381–445. MR 2444302 [6.4.3](#)
142. R. C. Gunning, *Lectures on Riemann surfaces*, Princeton Mathematical Notes, Princeton University Press, Princeton, N.J., 1966. MR 0207977 [10.2](#), [14](#), [14.1.2](#)

143. ———, *Special coordinate coverings of Riemann surfaces*, Math. Ann. **170** (1967), 67–86. MR 0207978 [10.2](#), [14](#)
144. Andre Haefliger, *Lectures on the theorem of Gromov*, Proceedings of Liverpool Singularities Symposium II, Springer, 1971, pp. 128–141. [7.5](#)
145. Dennis A. Hejhal, *Monodromy groups and linearly polymorphic functions*, Discontinuous groups and Riemann surfaces (Proc. Conf., Univ. Maryland, College Park, Md., 1973), Princeton Univ. Press, Princeton, N.J., 1974, pp. 247–261. Ann. of Math. Studies, No. 79. MR 0355035 [7.2.1](#)
146. ———, *Monodromy groups and linearly polymorphic functions*, Acta Math. **135** (1975), no. 1, 1–55. MR 0463429 [7.2.1](#), [14](#)
147. Jacques Helmstetter, *Radical d'une algèbre symétrique à gauche*, Annales de l'institut Fourier **29** (1979), no. 4, 17–35. [10](#), [10.3.1](#), [10.3.2](#), [10.3.4](#), [11.1](#)
148. Robert Hermann, *Gauge fields and cartan-ehresmann connections, part a*, vol. 10, Math Science Press, 1975. [5](#)
149. Morris W Hirsch, *Immersions of manifolds*, Trans. Amer. Math. Soc **93** (1959), no. 2, 242–276. [7.5](#)
150. Morris W. Hirsch and William P. Thurston, *Foliated bundles, invariant measures and flat manifolds*, Ann. Math. (2) **101** (1975), 369–390. MR 0370615 (51 #6842) [9.2](#)
151. N. J. Hitchin, *Lie groups and Teichmüller space*, Topology **31** (1992), no. 3, 449–473. MR 1174252 [13.3.1](#)
152. Ryan Hoban, *Affine automorphisms of properly convex domains*, Masters scholarly paper, University of Maryland 2006. [4.3](#)
153. John H. Hubbard, *The monodromy of projective structures*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 257–275. MR 624819 [7.2.1](#)
154. John Hamal Hubbard, *Teichmüller theory and applications to geometry, topology, and dynamics. Vol. 1*, Matrix Editions, Ithaca, NY, 2006, Teichmüller theory, With contributions by Adrien Douady, William Dunbar, Roland Roeder, Sylvain Bonnot, David Brown, Allen Hatcher, Chris Hruska and Sudeb Mitra, With forewords by William Thurston and Clifford Earle. MR 2245223 [1](#), [7.4](#), [14](#), [14.1.2](#)
155. James E. Humphreys, *Linear algebraic groups*, Springer-Verlag, New York-Heidelberg, 1975, Graduate Texts in Mathematics, No. 21. MR 0396773 [8.4](#)

156. Paul Igodt, Herbert Abels, Yves Félix, and Fritz Grunewald (eds.), *Crystallographic groups and their generalizations*, Contemporary Mathematics, vol. 262, American Mathematical Society, Providence, RI, 2000. MR 1796123 [15.4](#), [21](#), [93](#)
157. Nathan Jacobson, *Basic algebra. I*, second ed., W. H. Freeman and Company, New York, 1985. MR 780184 (86d:00001) [A.3](#)
158. Lizhen Ji and Athanase Papadopoulos (eds.), *Sophus Lie and Felix Klein: the Erlangen program and its impact in mathematics and physics*, IRMA Lectures in Mathematics and Theoretical Physics, vol. 23, European Mathematical Society (EMS), Zürich, 2015. MR 3379703 [5](#)
159. Kyeonghee Jo, *Homogeneity, quasi-homogeneity and differentiability of domains*, Proc. Japan Acad. Ser. A Math. Sci. **79** (2003), no. 9, 150–153. MR 2022059 [4.6](#)
160. ———, *Quasi-homogeneous domains and convex affine manifolds*, Topology Appl. **134** (2003), no. 2, 123–146. MR 2009094 [4.6](#)
161. ———, *A rigidity result for domains with a locally strictly convex point*, Adv. Geom. **8** (2008), no. 3, 315–328. MR 2427461 [4.6](#)
162. ———, *Domains with flat boundary pieces*, Bull. Korean Math. Soc. **53** (2016), no. 6, 1879–1886. MR 3582484 [4.6](#)
163. Kyeonghee Jo and Inkang Kim, *Convex affine domains and Markus conjecture*, Math. Z. **248** (2004), no. 1, 173–182. MR 2092727 [11.1](#)
164. Dennis Johnson and John J. Millson, *Deformation spaces associated to compact hyperbolic manifolds*, Discrete groups in geometry and analysis (New Haven, Conn., 1984), Progr. Math., vol. 67, Birkhäuser Boston, Boston, MA, 1987, pp. 48–106. MR 900823 (88j:22010) [7.3.2](#)
165. Yoshinobu Kamishima and Ser P. Tan, *Deformation spaces on geometric structures*, Aspects of low-dimensional manifolds, Adv. Stud. Pure Math., vol. 20, Kinokuniya, Tokyo, 1992, pp. 263–299. MR 1208313 [7.3.2](#), [14](#), [15.1](#)
166. Michael Kapovich, *Deformation spaces of flat conformal structures*, Questions and Answers in General Topology Special Issue **8** (1990), 253–264. [7.2](#)
167. ———, *Hyperbolic manifolds and discrete groups*, Progress in Mathematics, vol. 183, Birkhäuser Boston, Inc., Boston, MA, 2001. MR 1792613 [7.2.1](#), [7.3](#), [7.3](#), [14](#)
168. ———, *Convex projective structures on Gromov-Thurston manifolds*, Geom. Topol. **11** (2007), 1777–1830. MR 2350468 [4.6](#)

169. Steven P. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. (2) **117** (1983), no. 2, 235–265. MR 690845 [7.3.2](#)
170. Hyuk Kim, *Complete left-invariant affine structures on nilpotent Lie groups*, J. Differential Geom. **24** (1986), no. 3, 373–394. MR 868976 [10.3.1](#), [10.3.9](#), [10.5.1](#)
171. ———, *The geometry of left-symmetric algebra*, J. Korean Math. Soc. **33** (1996), no. 4, 1047–1067. MR 1424207 [10](#)
172. Felix Klein, *Le programme d'Erlangen*, Collection “Discours de la Méthode”, Gauthier-Villars Éditeur, Paris-Brussels-Montreal, Que., 1974, Considérations comparatives sur les recherches géométriques modernes, Traduit de l'allemand par H. Padé, Préface de J. Dieudonné, Postface de François Russo. MR 0354290 [5](#)
173. Bruno Klingler, *Complétude des variétés lorentziennes à courbure constante*, Math. Ann. **306** (1996), no. 2, 353–370. MR 1411352 [10.4.3.1](#)
174. ———, *Structures affines et projectives sur les surfaces complexes*, Ann. Inst. Fourier (Grenoble) **48** (1998), no. 2, 441–477. MR 1625606 [14](#)
175. ———, *Chern's conjecture for special affine manifolds*, Ann. of Math. (2) **186** (2017), no. 1, 69–95. MR 3665000 ([document](#)), [9.2](#)
176. Shoshichi Kobayashi, *Transformation groups in differential geometry*, Springer-Verlag, New York, 1972, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 70. MR 0355886 (50 #8360) [5](#)
177. ———, *Intrinsic distances associated with flat affine and projective structures*, J. Fac. Sci. Univ. Tokyo **24** (1977), 129–135. ([document](#)), [4](#), [12](#), [12.2.1](#)
178. ———, *Invariant distances for projective structures*, Symposia Mathematica, Vol. XXVI (Rome, 1980), Academic Press, London, 1982, pp. 153–161. MR 663030 (83i:53037) [4](#), [12.2.1](#)
179. ———, *Projectively invariant distances for affine and projective structures*, Banach Center Publ., vol. 12, PWN, Warsaw, 1984. MR 961077 (89k:53043 ([document](#)), [4](#), [4.2](#), [4.2.1](#), [8.3.2](#), [12](#), [12.1.3](#), [12.2.1](#), [12.2.8](#), [12.2.2](#), [12.2.2](#), [12.2.3](#), [12.2.3](#))
180. ———, *Hyperbolic manifolds and holomorphic mappings*, second ed., World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005, An introduction. MR 2194466 [12.2.1](#), [C.2.1](#), [C.2](#)
181. Shoshichi Kobayashi and Katsumi Nomizu, *Foundations of differential geometry. Vol. I*, Wiley Classics Library, John Wiley &

- Sons, Inc., New York, 1996, Reprint of the 1963 original, A Wiley-Interscience Publication. MR 1393940 (97c:53001a) ([document](#)), [5](#), [8.1.1](#), [8.1.1](#), [B.1](#)
182. B. Kostant and D. Sullivan, *The Euler characteristic of an affine space form is zero*, Bull. Amer. Math. Soc. **81** (1975), no. 5, 937–938. MR 0375341 (51 #11536) ([document](#)), [9.2.3](#), [9.2.7](#), [9.2.8](#)
183. Jean-Louis Koszul, *Domaines bornés homogènes et orbites de groupes de transformations affines*, Bull. Soc. Math. France **89** (1961), 515–533. MR 0145559 (26 #3090) [10](#), [12.3](#)
184. ———, *Ouverts convexes homogènes des espaces affines*, Mathematische Zeitschrift **79** (1962), no. 1, 254–259. [12.3](#)
185. ———, *Lectures on groups of transformations*, Notes by R. R. Simha and R. Sridharan. Tata Institute of Fundamental Research Lectures on Mathematics, No. 32, Tata Institute of Fundamental Research, Bombay, 1965. MR 0218485 (36 #1571) [7.2.1](#), [A.3](#)
186. ———, *Variétés localement plates et convexité*, Osaka J. Math. **2** (1965), 285–290. MR 0196662 (33 #4849) [8.3.10](#), [12.3](#)
187. ———, *Déformations de connexions localement plates*, Ann. Inst. Fourier (Grenoble) **18** (1968), no. fasc. 1, 103–114. MR 0239529 (39 #886) [12.3](#)
188. Nicolaas H. Kuiper, *Compact spaces with a local structure determined by the group of similarity transformations in E^n* , Nederl. Akad. Wetensch., Proc. **53** (1950), 1178–1185 = Indagationes Math. 12, 411–418 (1950). MR 0039248 [11.4](#)
189. ———, *On compact conformally Euclidean spaces of dimension > 2* , Ann. of Math. (2) **52** (1950), 478–490. MR 0037575 [14.2](#)
190. ———, *Locally projective spaces of dimension one.*, The Michigan Mathematical Journal **2** (1953), no. 2, 95–97. [5.4](#)
191. ———, *Sur les surfaces localement affines*, Géométrie différentielle. Colloques Internationaux du Centre National de la Recherche Scientifique, Strasbourg, 1953, 1953, pp. 79–87. [8.5](#), [10.2](#)
192. ———, *On convex locally projective spaces*, Convegno Internazionale di Geometria Differenziale (Italia, 1953) (1954), 200–213. [4.5.2.3](#), [13.2](#)
193. ———, *Hyperbolic 4-manifolds and tessellations*, Inst. Hautes Études Sci. Publ. Math. (1988), no. 68, 47–76 (1989). MR 1001447 [15.1](#)
194. Ravi S. Kulkarni and Ulrich Pinkall, *Uniformization of geometric structures with applications to conformal geometry*, Differential geometry, Peñíscola 1985, Lecture Notes in Math., vol. 1209,

- Springer, Berlin, 1986, pp. 190–209. MR 863757 (88b:53036)
[2.6.4](#), [11.4](#), [14.2](#), [14.2.2](#), [15.1](#)
195. ———, *A canonical metric for Möbius structures and its applications*, Math. Z. **216** (1994), no. 1, 89–129. MR 1273468 [11.4](#), [15.1](#)
196. François Labourie, *Fuchsian affine actions of surface groups*, J. Differential Geom. **59** (2001), no. 1, 15–31. MR 1909247 [15.3](#), [15.3.3](#)
197. ———, *Flat projective structures on surfaces and cubic holomorphic differentials*, Pure Appl. Math. Q. **3** (2007), no. 4, Special Issue: In honor of Grigory Margulis. Part 1, 1057–1099. MR 2402597 [13.3.2](#), [13.3.2](#)
198. François Labourie, *Anosov flows, surface groups and curves in projective space*, Invent. Math. **165** (2006), no. 1, 51–114. MR 2221137 [13.3.1](#)
199. Sean Lawton, *Generators, relations and symmetries in pairs of 3×3 unimodular matrices*, J. Algebra **313** (2007), no. 2, 782–801. MR 2329569 [2](#)
200. John M. Lee, *Riemannian manifolds*, Graduate Texts in Mathematics, vol. 176, Springer-Verlag, New York, 1997, An introduction to curvature. MR 1468735 ([document](#)), [8.1.1](#)
201. ———, *Introduction to smooth manifolds*, second ed., Graduate Texts in Mathematics, vol. 218, Springer, New York, 2013. MR 2954043 ([document](#))
202. Walter L. Lok, *Deformations of locally homogeneous spaces and kleinian groups*, Ph.D. thesis, Columbia University, 1984. [7.2.1](#)
203. Darren Long, Alan W. Reid, and Morwen Thistlethwaite, *Zariski dense surface subgroups in $SL(3, \mathbf{Z})$* , Geom. Topol. **15** (2011), no. 1, 1–9. MR 2764111 [13.2.1](#)
204. Alexander Lubotzky and Andy R. Magid, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. **58** (1985), no. 336, xi+117. MR 818915 [7.3](#)
205. Anton Lukyanenko, *Projective deformations of triangle tilings*, Master’s thesis, University of Maryland, 2008. [13.2.1](#)
206. A. Marden, *Outer circles*, Cambridge University Press, Cambridge, 2007, An introduction to hyperbolic 3-manifolds. MR 2355387 [14](#)
207. G. A. Margulis, *Discrete groups of motions of manifolds of non-positive curvature*, Proceedings of the International Congress of Mathematicians (Vancouver, B.C., 1974), Vol. 2, Canad. Math. Congress, Montreal, Que., 1975, pp. 21–34. MR 0492072 [11.3](#)

208. ———, *Complete affine locally flat manifolds with a free fundamental group*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **134** (1984), 190–205, Automorphic functions and number theory, II. MR 741860 [15.1](#)
209. Lawrence Markus, *Cosmological models in differential geometry*, mimeographed book, Institute of Technology, Department of Mathematics, University of Minnesota, Minneapolis, Minnesota, (1963). ([document](#)), [11](#)
210. Ludovic Marquis, *Espace des modules marqués des surfaces projectives convexes de volume fini*, Geom. Topol. **14** (2010), no. 4, 2103–2149. MR 2740643 [4](#)
211. ———, *Exemples de variétés projectives strictement convexes de volume fini en dimension quelconque*, Enseign. Math. (2) **58** (2012), no. 1-2, 3–47. MR 2985008 [4](#)
212. ———, *Around groups in Hilbert geometry*, Handbook of Hilbert geometry, IRMA Lect. Math. Theor. Phys., vol. 22, Eur. Math. Soc., Zürich, 2014, pp. 207–261. MR 3329882 [4](#)
213. Jerrold Marsden, *On completeness of homogeneous pseudo-riemannian manifolds*, Indiana Univ. Math. J. **22** (1972/73), 1065–1066. MR 319128 [10.4.3.1](#)
214. Bernard Maskit, *On a class of Kleinian groups*, Ann. Acad. Sci. Fenn. Ser. A I No. **442** (1969), 8. MR 0252638 [14](#)
215. Shigenori Matsumoto, *Foundations of flat conformal structure*, Aspects of low-dimensional manifolds, Adv. Stud. Pure Math., vol. 20, Kinokuniya, Tokyo, 1992, pp. 167–261. MR 1208312 [14](#), [15.1](#)
216. Yozô Matsushima, *Affine structures on complex manifolds*, Osaka Math. J. **5** (1968), 215–222. MR 240741 [10](#)
217. Benjamin McKay, *Holomorphic geometric structures on Kähler-Einstein manifolds*, Manuscripta Math. **153** (2017), no. 1-2, 1–34. MR 3635971 [14](#)
218. Curtis T. McMullen, *Complex earthquakes and Teichmüller theory*, J. Amer. Math. Soc. **11** (1998), no. 2, 283–320. MR 1478844 [7.3.2](#), [13.2.3](#)
219. Alberto Medina Perea, *Flat left-invariant connections adapted to the automorphism structure of a Lie group*, J. Differential Geom. **16** (1981), no. 3, 445–474 (1982). MR 654637 [10](#), [10.3.1](#)
220. Geoffrey Mess, *Lorentz spacetimes of constant curvature*, Geom. Dedicata **126** (2007), 3–45. MR 2328921 [15.3](#)
221. John W. Milnor, *On the existence of a connection with curvature zero*, Commentarii Mathematici Helvetici **32** (1958), no. 1, 215–223. [9.1](#), [9.1.3](#), [9.2.1](#), [9.2.2](#)

222. ———, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963. MR 0163331 ([document](#)), [8.1.1](#)
223. ———, *On the 3-dimensional Brieskorn manifolds $M(p, q, r)$* , Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox), Princeton Univ. Press, Princeton, N. J., 1975, pp. 175–225. Ann. of Math. Studies, No. 84. MR 0418127 (54 #6169) [11.3](#)
224. ———, *Hilbert's problem 18: on crystallographic groups, fundamental domains, and on sphere packing*, Mathematical developments arising from Hilbert problems (Proc. Sympos. Pure Math., Northern Illinois Univ., De Kalb, Ill., 1974), Amer. Math. Soc., Providence, R. I., 1976, pp. 491–506. Proc. Sympos. Pure Math., Vol. XXVIII. MR 0430101 [8.6.2](#)
225. ———, *On fundamental groups of complete affinely flat manifolds*, Advances in Mathematics **25** (1977), no. 2, 178–187. [8.6](#), [15.1](#), [15.2](#)
226. John W. Milnor and James D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974, Annals of Mathematics Studies, No. 76. MR 0440554 (55 #13428) [9.2.1](#), [9.2.2](#)
227. Robert R. Miner, *Spherical CR manifolds with amenable holonomy*, Internat. J. Math. **1** (1990), no. 4, 479–501. MR 1080108 [11.4](#)
228. Tadashi Nagano and Katsumi Yagi, *The affine structures on the real two-torus. I*, Osaka Journal of Mathematics **11** (1974), no. 1, 181–210. [10.2](#)
229. Katsumi Nomizu and Takeshi Sasaki, *Affine differential geometry*, Cambridge Tracts in Mathematics, vol. 111, Cambridge University Press, Cambridge, 1994, Geometry of affine immersions. MR 1311248 [13.3.2](#)
230. Barrett O'Neill, *Semi-Riemannian geometry*, Pure and Applied Mathematics, vol. 103, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983, With applications to relativity. MR 719023 ([document](#)), [8.1.1](#), [B.1](#)
231. V. Ovsienko and S. Tabachnikov, *Projective differential geometry old and new*, Cambridge Tracts in Mathematics, vol. 165, Cambridge University Press, Cambridge, 2005, From the Schwarzian derivative to the cohomology of diffeomorphism groups. MR 2177471 [1](#), [13.3.1](#), [B.2](#)

232. Athanase Papadopoulos, *Metric spaces, convexity and non-positive curvature*, second ed., IRMA Lectures in Mathematics and Theoretical Physics, vol. 6, European Mathematical Society (EMS), Zürich, 2014. MR 3156529 [8.1.1](#)
233. Athanase Papadopoulos and Marc Troyanov (eds.), *Handbook of Hilbert geometry*, IRMA Lectures in Mathematics and Theoretical Physics, vol. 22, European Mathematical Society (EMS), Zürich, 2014. MR 3309067 [3](#)
234. R. C. Penner, *The decorated Teichmüller space of punctured surfaces*, *Comm. Math. Phys.* **113** (1987), no. 2, 299–339. MR 919235 [13.1](#)
235. Robert C. Penner, *Decorated Teichmüller theory*, QGM Master Class Series, European Mathematical Society (EMS), Zürich, 2012, With a foreword by Yuri I. Manin. MR 3052157 [13.1](#)
236. Alekseĭ Vasilevich Pogorelov, *Hilbert's fourth problem*, Halsted Press, 1979. [3.2.2](#)
237. Walter A. Poor, *Differential geometric structures*, McGraw-Hill Book Co., New York, 1981. MR 647949 [9.2.1](#)
238. M. S. Raghunathan, *Discrete subgroups of Lie groups*, Springer-Verlag, New York-Heidelberg, 1972, *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68*. MR 0507234 [7.2.1](#), [7.3](#), [7.3](#), [8.6.2](#)
239. John G. Ratcliffe, *Foundations of hyperbolic manifolds*, second ed., Graduate Texts in Mathematics, vol. 149, Springer, New York, 2006. MR 2249478 [3](#), [11.2](#)
240. Maxwell Rosenlicht, *On quotient varieties and the affine embedding of certain homogeneous spaces*, *Trans. Amer. Math. Soc.* **101** (1961), 211–223. MR 130878 [10.5](#)
241. O. S. Rothaus, *The construction of homogeneous convex cones*, *Ann. of Math. (2)* **83** (1966), 358–376. MR 202156 [10](#)
242. H. L. Royden, *Remarks on the Kobayashi metric*, *Several complex variables, II* (Proc. Internat. Conf., Univ. Maryland, College Park, Md., 1970) (John Horvath, ed.), 1971, pp. 125–137. *Lecture Notes in Math.*, Vol. 185, Springer-Verlag. MR 0304694 [12.2.2](#), [12.2.2](#)
243. Walter Rudin, *Real and complex analysis*, third ed., McGraw-Hill Book Co., New York, 1987. MR 924157 [12.2.2](#), [D.1.2](#)
244. John Scheuneman, *Translations in certain groups of affine motions*, *Proc. Amer. Math. Soc.* **47** (1975), 223–228. MR 372120 [10.5.1](#), [10.5.1](#)
245. Richard Evan Schwartz, *Spherical CR geometry and Dehn surgery*, *Annals of Mathematics Studies*, vol. 165, Princeton University Press, Princeton, NJ, 2007. MR 2286868 [15.1](#)

246. Peter Scott, *The geometries of 3-manifolds*, Bulletin of the London Mathematical Society **15** (1983), no. 5, 401–487. ([document](#))
247. Dan Segal, *The structure of complete left-symmetric algebras*, Math. Ann. **293** (1992), no. 3, 569–578. MR 1170527 [10](#), [10.3.1](#), [10.3.2](#), [10.3.2](#)
248. J. G. Semple and G. T. Kneebone, *Algebraic projective geometry*, Oxford, at the Clarendon Press, 1952. MR 0049579 [2](#), [1](#), [3](#)
249. Richard W Sharpe, *Differential geometry: Cartan's generalization of Klein's erlangen program*, vol. 166, Springer, 1997. ([document](#)), [5](#), [7.1.4](#), [14.1.2](#)
250. Hirohiko Shima, *The geometry of Hessian structures*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007. MR 2293045 (2008f:53011) [8.3.2](#), [12.3](#)
251. Adam S. Sikora, *Character varieties*, Trans. Amer. Math. Soc. **364** (2012), no. 10, 5173–5208. MR 2931326 [7.3](#)
252. John Smillie, *The Euler characteristic of flat bundles*, Preprint. [9.1](#), [9.1.3](#)
253. ———, *Affinely flat manifolds*, Ph.D. thesis, University of Chicago, 1977. [11.1](#), [11.2](#), [11.2.3](#), [11.2](#), [13](#), [13.4.0.1](#)
254. ———, *Flat manifolds with non-zero Euler characteristics*, Commentarii Mathematici Helvetici **52** (1977), no. 1, 453–455. [9.2.2](#)
255. ———, *An obstruction to the existence of affine structures*, Inventiones mathematicae **64** (1981), no. 3, 411–415. [11.3.1](#), [11.3](#), [11.3.2](#)
256. Edith Socié-Méthou, *Caractérisation des ellipsoïdes par leurs groupes d'automorphismes*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 4, 537–548. MR 1981171 [4.6](#)
257. Stephen Bruce Sontz, *Principal bundles*, Universitext, Springer, Cham, 2015, The classical case. MR 3309256 ([document](#))
258. Michael Spivak, *A comprehensive introduction to differential geometry. Vol. II*, Published by M. Spivak, Brandeis Univ., Waltham, Mass., 1970. MR 0271845 [B.2](#)
259. Norman Earl Steenrod, *The topology of fibre bundles*, vol. 14, Princeton University Press, 1951. ([document](#)), [5.3](#), [5.3.1](#), [9.2.1](#)
260. Dennis Sullivan, *La classe d'Euler réelle d'un fibré vectoriel à groupe structural $SL_n(\mathbb{Z})$ est nulle*, C. R. Acad. Sci. Paris Sér. A-B **281** (1975), no. 1, Aii, A17–A18. MR 0372873 (51 #9077) [9.2](#)
261. ———, *A generalization of Milnor's inequality concerning affine foliations and affine manifolds*, Commentarii Mathematici Helvetici **51** (1976), no. 1, 183–189. [9.1](#), [9.1.3](#)

262. Dennis Sullivan and William Thurston, *Manifolds with canonical coordinate charts: some examples*, Enseign. Math **29** (1983), 15–25. [13](#), [13.4.0.1](#), [14](#)
263. Georges Thone (ed.), *Colloque de topologie (espaces fibrés), Bruxelles, 1950*, Masson et Cie., Paris, 1951. MR 0042768 [5.2](#), [97](#)
264. William P. Thurston, *Minimal stretch maps between hyperbolic surfaces*. [15.3.4](#)
265. ———, *The geometry and topology of three-manifolds*, unpublished notes, Princeton University Mathematics Department, 1979. ([document](#)), [3.2.4](#), [7.2](#), [7.2.1](#), [15.4](#)
266. ———, *Three-dimensional geometry and topology. Vol. 1*, Princeton Mathematical Series, vol. 35, Princeton University Press, Princeton, NJ, 1997, Edited by Silvio Levy. MR 1435975 ([document](#)), [3](#), [5](#), [11.2](#)
267. D. Tischler, *On fibering certain foliated manifolds over S^1* , Topology **9** (1970), 153–154. MR 256413 [12.3](#)
268. J. Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270. MR 286898 [8.6](#)
269. ———, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270. MR 286898 [15.2](#)
270. Izu Vaisman and Corina Reischer, *Local similarity manifolds*, Ann. Mat. Pura Appl. (4) **135** (1983), 279–291 (1984). MR 750537 ([document](#)), [11.4](#)
271. Oswald Veblen and John Wesley Young, *Projective geometry. Vol. 1*, Blaisdell Publishing Co. Ginn and Co. New York-Toronto-London, 1965. MR 0179666 [2](#), [1](#), [2.5.2](#)
272. ———, *Projective geometry. Vol. 2 (by Oswald Veblen)*, Blaisdell Publishing Co. Ginn and Co. New York-Toronto-London, 1965. MR 0179667 [2](#)
273. Edoardo Vesentini, *Invariant metrics on convex cones*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **3** (1976), no. 4, 671–696. MR 0433228 [12.3](#)
274. Jacques Vey, *Une notion d'hyperbolicité sur les variétés localement plates*, C. R. Acad. Sci. Paris Sér. A-B **266** (1968), A622–A624. MR 236837 [12](#), [12.2.1](#)
275. ———, *Sur une notion d'hyperbolicité des variétés localement plates*, Ph.D. thesis, Université Joseph-Fourier-Grenoble I, 1969. ([document](#)), [1.2.2](#), [4](#), [10](#), [10.3.1](#), [12](#), [12.2.1](#), [12.2.1](#)
276. ———, *Sur les automorphismes affines des ouverts convexes dans les espaces numériques*, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A249–A251. MR 271839 [4.2](#), [4.3](#), [12](#)

277. ———, *Sur les automorphismes affines des ouverts convexes sail-lants*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) **24** (1970), 641–665. MR 283720 [4.2](#), [4.3](#), [12](#), [15.4](#)
278. È. B. Vinberg, *The theory of homogeneous convex cones*, Trudy Moskov. Mat. Obšč. **12** (1963), 303–358. MR 0158414 ([document](#)), [4](#), [4.4.3](#), [4.4.9](#), [10](#), [10.3.1](#), [10.4.4](#), [10.4.4.3](#), [12.3](#)
279. E.B. Vinberg and V.G. Kats, *Quasi-homogeneous cones*, Mathematical Notes **1** (1967), no. 3, 231–235. [13.2.1](#)
280. H. Vogt, *Sur les invariants fondamentaux des équations différentielles linéaires du second ordre*, Ann. Sci. École Norm. Sup. (3) **6** (1889), 3–71. MR 1508833 [7.3.1](#), [7.3.6](#)
281. André Weil, *On discrete subgroups of Lie groups*, Ann. of Math. (2) **72** (1960), 369–384. MR 0137792 (25 #1241) [7.2.1](#)
282. J. H. C. Whitehead, *Locally homogeneous spaces in differential geometry*, Ann. of Math. (2) **33** (1932), no. 4, 681–687. MR 1503084 ([document](#))
283. John Henry Constantine Whitehead, *The representation of projective spaces*, Annals of Mathematics (1931), 327–360. ([document](#))
284. Anna Wienhard and Tengren Zhang, *Deforming convex real projective structures*, Geom. Dedicata **192** (2018), 327–360. MR 3749433 [13.2.1](#)
285. Stephen Willard, *General topology*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970. MR 0264581 ([document](#))
286. Joseph A. Wolf, *Spaces of constant curvature*, third ed., Publish or Perish, Inc., Boston, Mass., 1974. MR 0343214 [6](#)
287. Scott Wolpert, *An elementary formula for the Fenchel-Nielsen twist*, Comment. Math. Helv. **56** (1981), no. 1, 132–135. MR 615620 [7.4](#)
288. ———, *The Fenchel-Nielsen deformation*, Ann. of Math. (2) **115** (1982), no. 3, 501–528. MR 657237 [7.4](#), [7.4](#)
289. ———, *On the symplectic geometry of deformations of a hyperbolic surface*, Ann. of Math. (2) **117** (1983), no. 2, 207–234. MR 690844 [7.4](#), [7.4](#)
290. ———, *On the Weil-Petersson geometry of the moduli space of curves*, Amer. J. Math. **107** (1985), no. 4, 969–997. MR 796909 [7.4](#), [7.4](#)
291. H. Wu, *Some theorems on projective hyperbolicity*, J. Math. Soc. Japan **33** (1981), no. 1, 79–104. MR 597482 (82j:53061) [12](#), [12.2.2](#), [12.2.2](#)

292. Tengren Zhang, *The degeneration of convex \mathbb{RP}^2 structures on surfaces*, Proc. Lond. Math. Soc. (3) **111** (2015), no. 5, 967–1012.
MR 3477227 [13.2.1](#)

List of Figures

2.1 A perspectivity between two lines l, l' in the plane. The undefined set of $\{O\}$ of the projection is the <i>center</i> of the perspectivity.	50
2.2 The four points on the horizontal line form a harmonic quadruple. Such a quadruple is characterized by having cross-ratio -1 .	53
2.3 Non-Euclidean tessellations by equilateral triangles	55
3.1 The <i>Klein-Beltrami</i> projective model of hyperbolic space. The Hilbert metric on the convex domain bounded by a quadric in projective space is defined in terms of cross-ratio. This metric is a Riemannian metric of constant negative curvature.	75
3.2 Projective model of a $(3, 3, 4)$ -triangle tessellation of \mathbf{H}^2	79
3.3 Projective deformation of hyperbolic $(3, 3, 4)$ -triangle tessellation	79
4.1 Projection does not increase Hilbert distance	88
4.2 c_t is an isometry of the cone	89
4.3 $s(\Omega)$ is convex.	90
4.4 A sequence of projectively equivalent convex domains with a corner converging to a triangle with the corner as a vertex.	101
4.5 A sequence of projectively equivalent convex domains with a flat part of the boundary converging to a triangle having the flat part as a side.	101
5.1 Extending a coordinate chart to a developing map	119
5.2 The graph of a developing map is an \mathcal{F} -transverse section	125
5.3 The isotopy between nearby \mathcal{F} -transverse sections	126
5.4 Developing section for the canonical projective closed 1-manifold $\mathbb{R}P^1$	129
5.5 Developing section for other projective closed 1-manifolds with trivial holonomy	129

5.6 Developing sections for projective closed 1-manifolds with elliptic holonomy	130
5.7 Developing sections for projective closed 1-manifolds with hyperbolic holonomy	130
5.8 Developing sections for projective closed 1-manifolds with parabolic holonomy	131
5.9 Some incomplete complex-affine structures on T^2	138
5.10 Some hyperbolic affine structures on T^2	139
5.11 Radiant affine structures on T^2 developing to a halfplane	140
5.12 Nonradiant affine structures on T^2 developing to a halfplane	141
7.1 Immersions of a one-holed torus into the plane and the sphere.	180
7.2 Development of one-holed torus with holonomy generated by translation	180
8.1 Tilings corresponding to some complete affine structures on the 2-torus. The second picture depicts a complete non-Riemannian deformation where the affine holonomy contains no nontrivial horizontal translation. The corresponding torus contains no closed geodesics.	197
9.1 Decomposing a genus $g = 2$ surface along $2g$ curves into a $4g$ -gon. The single common intersection of the curves is a single point which decomposes into the $4g$ vertices of the polygon.	207
9.2 Identifying the edges of a $4g$ -gon into a closed surface of genus g . The sides are paired into $2g$ curves, which meet at the single vertex.	207
9.3 Cell-division of torus where all but one angle at the vertex is 0.	211
9.4 Doubly periodic tiling of the Euclidean plane with all but one angle at vertex 0.	211
13.1A projective transformation leaving invariant a closed convex curve.	295
13.2 Conics tangent to a triangle	299
13.3 Deforming a conic	300
13.4A piecewise conic	300
13.5 Bulging data	301
13.6 The deformed conic	301
13.7 The conic with its deformation	301

LIST OF FIGURES

379

13. Pathological development for $\mathbb{R}P^2$ -torus

306

14. Octagonal fundamental domain for a $\mathbb{C}P^1$ -surface of genus two

309

D.1 Continuous approximation of indicator function

350

List of Tables

1	Multiplication tables for complete (Euclidean and non-Riemannian) structures	226
2	Multiplication tables for incomplete (radiant halfplane, nonradiant halfplane and hyperbolic) structures	226
3	A commutative non-associative 2-dimensional algebra	229
4	Left-invariant vector fields on $\text{Aff}_+(1, \mathbb{R})$ form an associative algebra $\mathfrak{a}_{\mathfrak{L}}$	234
5	Covariant derivatives of left-invariant vector fields with respect to the Levi-Civita connection of the Poincaré metric.	235
6	Right-invariant vector fields on $\text{Aff}_+(1, \mathbb{R})$ form an associative algebra $\mathfrak{a}_{\mathfrak{R}}$ opposite to $\mathfrak{a}_{\mathfrak{L}}$	236
7	Left-invariant vector fields for the complete left-invariant structures on $\text{Aff}_+(1, \mathbb{R})$	237
8	Left-invariant vector fields of deformation of bi-invariant structure	238
9	Left-invariant vector fields on the halfspace family structures define an algebra $\mathfrak{a}_{\mathfrak{L}}^{\beta}$ depending on a parameter β	239
10	Left-invariant vector fields on the parabolic deformation $\mathfrak{a}_{\mathfrak{L}}^2(\alpha)$	241
11	Nonradiant deformation of $\mathfrak{a}_{\mathfrak{R}}$ depending on a parameter $\eta \in \mathbb{R}$	243
12	Radiant deformation of $\mathfrak{a}_{\mathfrak{R}}$ depending on an eigenvalue parameter $\mu \in \mathbb{R}$	243
13	Deformations of $\mathfrak{a}_{\mathfrak{R}}$ depending on an eigenvalue parameter $\delta \in \mathbb{R}$ containing the complete structure at $\delta = 0$	244
14	Multiplication tables for $\dim(\mathfrak{3}) = 3$ and $\dim(\mathfrak{3}) = 2$, respectively	247
15	When $\dim(\mathfrak{3}) = 1$ and $G/\mathfrak{3}$ Euclidean, the structure is defined by bilinear form $G/\mathfrak{3} \times G/\mathfrak{3} \xrightarrow{\mathfrak{a}} \mathfrak{3}$	247
16	When $\dim(\mathfrak{3}) = 1$ and $G/\mathfrak{3}$ non-Riemannian, the structure is defined by a pair $(b, c) \in \mathbb{R}^2$	247
17	Complete affine structure on 4-dimensional filiform algebra	249

18	Fried's counterexample to Auslander's conjecture on central translations	250
19	Algebra corresponding to parabolic 3-dimensional halfspaces	253
20	A radiant suspension	254
21	Nonradiant Deformation	255

Index

- 1-manifolds
 - affine, 132
 - Euclidean, 131
 - projective, 131, 134
- h -principle, 179, 180
- action
 - proper, 333
 - simply transitive
 - locally, 223
 - syndetic, 333
 - wandering, 334
- affine, 23
 - connection, 186
 - geometry, 27
 - map, 26
 - patch, 51
 - structure
 - incomplete, 179
 - left-invariant, 221
 - subspace, 36
 - vector field, 33
- affine connection
 - projective equivalence of, 337
- affine parameter, 36
- Affine Schwarzian, 310
- algebra
 - commutative associative
 - 2-dimensional, 224
 - left-symmetric, 227
 - nilpotent associative
 - 3-dimensional, 248
- algebraicization
 - of classical geometries, 7
 - of geometric structures, 143
- anti-de Sitter geometry, 319
- Auslander-Milnor question, 198
 - in dimension 3, 319
- Benzécri's theorem
 - compactness, 100
 - on surfaces, 205
- Bieberbach theorems, 198
- Bonahon-Dreyer coordinates, 300
- boosts
 - affine, 322
- Bruhat decomposition, 310
- Cartesian product, 146
- centroid, 37, 38
- Chain rule, 17
- characteristic function
 - of left-symmetric algebra, 235
 - of convex cone, 91
- Chern-Weil theory, 213
- Choi's convex decomposition
 - theorem, 304
- Clans, 240
- collineation, 46
- complete affine structures, 198
 - on solvmanifolds, 199
 - on the two-torus, 194
- completeness
 - geodesic, 189
- complex affine structure, 154
- cone
 - dual, 90
- connections
 - affine, 187
 - geodesically complete, 188
- contact projective structure, 71, 145, 156
- convex, 82
- convex body, 38

- convex domain
 - divisible, 108
 - quasi-homogeneous, 108
- convexity
 - in projective space, 84
- corners, 102
- correlation, 69
- covariant differentiation and matrix
 - multiplication, 35
- Coxeter extension, 174
- crooked planes, 322
- cross-ratio, 57
- cross-section to flow, 147
- curvature tensor, 335

- developing map, 117
 - tameness of, 183
- dual projective space, 14

- earthquake deformation, 176
- Einstein Universe, 319
- ellipsoid, 105
 - of inertia, 105
- Elliptic geometry, 70
- elliptic polarity, 71
- elliptic space
 - projective model of, 70
- embedding of geometries, 145
- equivalence of categories, 333
- Erlanger Program, 1, 113
- Euclidean rational homology
 - 3-sphere, 149, 258
- Euler class, 214

- Fenchel-Nielsen coordinates, 178
- Fenchel-Nielsen section, 178
- Fenchel-Nielsen twist deformation, 176
- fibration of geometries, 146
- Fitting subspace, 245
- flat conformal structures, 317
- flat tori, 132
 - moduli of, 166
- flats in the boundary, 102
- Fock-Goncharov coordinates, 300
- Fricke space of three-holed sphere, 174
- Fubini-Study metric, 70
- functor
 - fully faithful, 333
- fundamental group, 16
- Fundamental Theorem on Flows, 19

- geodesic completeness, 181
- geodesic spray, 336
- geometric atlas, 114
- geometric structure transverse to
 - foliation, 156
- geometry
 - affine, 1, 27
 - Euclidean, 1
 - extending, 122
 - non-Euclidean, 2
 - projective, 1
 - similarity, 1
- grafting, 308
 - $\mathbb{R}P^2$ -structures, 304
 - in dimension one, 142

- Hölder exponent of limit set of
 - convex $\mathbb{R}P^2$ -structure, 295
- harmonic
 - homology, 53
 - set, 53
- harmonic net, 59
- harmonic subdivision, 59
- Hessian metrics, 290
- hex-metric, 86
- Hilbert distance, 75
- Hilbert metric, 84
- Hill's equation, 312
- holonomy representation, 117
- holonomy sequence, 264, 268
- Holonomy-invariant subdomains, 123
- homogeneous coordinates, 43
- homography, 46
- homothety, 34
- Hopf
 - circle, 133
 - manifold, 133
- Hopf manifolds, 151
 - geodesics on, 152
 - in dimension one, 133
- Hopf-Rinow theorem, 184
- hyperbolic geometry
 - projective model of, 74
- hyperbolic polarity, 72

- incidence, 43
- internal parameters, 297
- invisible, 188
- involutions, 46
- Jacobson product, 229
- Kac-Vinberg example, 80, 295
- Klein-Beltrami model of hyperbolic geometry, 75
- Kobayashi metric, 275
 - infinitesimal, 281
- Kostant-Sullivan theorem, 218
- Koszul formula, 338
- Labourie-Loftin coordinates, 303
- Lebesgue number, 344
- left-symmetric algebras
 - central translations, 246
 - simple, 241
- Levi-Civita connection, 234, 338
- Locally homogeneous Riemannian manifolds, 183
- Margulis spacetime, 319
- Margulis superrigidity, 263
- marked geometric structure, 165
- marking, 165
- Markus conjecture, 257
 - infinitesimal, 230
- maximal ball, 266
- Milnor-Wood inequality, 212
- moment of inertia, 105
- Murphy's Law, 171
- Naturality of Flows, 19
- net of rationality, 59
- non-Riemannian affine torus, 196
- nonradiant deformation, 253
- normal family, 65
- normality domain, 65
- null polarity, 71
- open manifolds, 179
- osculating conic, 102
- osculating Möbius transformation, 312
- Pappus's Theorem, 67
- parabolic convex region, 82
- parabolic cylinders, 252
- parallel
 - Riemannian structure, 187
- parallel structures, 31
- parallel transport, 29
- parallel vector field, 29
- parameter
 - affine, 36
- pathological developing maps, 304
- perspectivity, 49
- Poincaré metric, 234
- point of normality, 65
- point-symmetry in H^3 , 78
- polarity, 69
 - conjugate points of, 71
 - null, 67
- pre-Schwarzian, 309
- projection, 49
- projective chain
 - in a domain, 277
 - in a projective manifold, 280
- projective equivalence, 14
- projective reflection, 47
- projective vector fields, 52
- pullback
 - of a vector field, 19
- quadric, 72
- quakebend deformation, 176
- Quaternions, 76
- quotient structure, 183
- reflection in H^3 , 78
- representation
 - étale, 223
- Riemann moduli space, 165
- Riemannian structures
 - parallel, 31
- scale factor homomorphism, 32, 265
- Schwarzian differential, 309
- Seifert 3-manifold, 263
- semicontinuous function, 189, 282, 347
- similarity manifolds, 263
 - complete, 264
 - radiant, 160
- similarity transformations, 32
- simple left-symmetric algebra, 241

- singular projective transformation,
 - 60
 - range, 60
 - undefined set, 60
- Smale-Hirsch immersion theorem, 180
- Smillie-Thurston example, 304, 305
- spherical CR-structures, 318
- stably parallelizable manifolds, 216
- stably trivial vector bundle, 216
- Sturm-Liouville equation, 312
- suspension
 - parallel, 147
 - radiant, 160
- symmetry about a point, 76
- syndetic, 100

- Teichmüller space, 165
- thin subgroup, 296
- topological transformation
 - groupoids, 333
- tori
 - flat, 132
- torsion tensor, 335
- torsor, 26
- transpose, 69
- Triple ratio, 302
- tube domain, 92
- turning number, 208
- Twist flows, 178

- unipotent holonomy, 192
- Unique Extension Property, 50, 115
- unit ball
 - Euclidean, 74
 - in Hilbert metric, 85
- universal covering space, 16
- Upper halfspace model of hyperbolic
 - 3-space, 76

- vector field
 - affine, 33
 - Euler, 34
 - parallel, 33
 - pullback of, 19
 - radiant, 34
- Vey semisimplicity theorem, 86
- visible, 188
- Vogt-Fricke theorem, 173
- volume obstruction, 257
- Weil-Petersson symplectic structure, 178
- Whitney-Graustein theorem, 208