## AMSC/CMSC 460 Problem set 5

1. Write a MATLAB program to implement the trapezoid rule. Make it a function trap(f,a,b,n) where f is the function to be integrated over the interval [a,b] and n is the number of subintervals. The function f(x)should be written as inline function. The function mfile trap.m will use the instruction feval. This instruction passes the variable x to the named function with the command feval(f,x). For more info, enter help feval. The call for trap will be trap(f,a,b,n). Output should be the trapezoid estimate for the integral.

Apply the trapezoid rule to estimate each of the following integrals with n = 2, 4, 8, 16, 32, 64.

(i) 
$$\frac{2}{\sqrt{\pi}} \int_0^1 e^{-x^2} dx = \operatorname{erf}(1)$$
 (ii)  $\int_0^1 x^{5/2} dx$   
(iii)  $\int_{-4}^4 \frac{dx}{1+x^2}$  (iv)  $\int_0^{2\pi} \frac{dx}{2+\cos(x)}$ .

(a) In cases (i), (ii), and (iii), compare your result with the exact answer (for case (i) this means compare with the MATLAB value for the error function). What power law governs the decrease of the error? Does this agree with the error behavior predicted by the error formula?

(b) In case (iv), we do not have an exact answer, but you will see that the results converge very quickly. Why?

2. Write a function mfile to implement Simpson's rule and apply it to the integrals of problem 1, same questions.

**3.** Now write a trapezoid program, myquad, with input a function f(x) (given as an inline function), the interval [a, b] and a tolerance  $\varepsilon$ . The call should be myquad(f,a,b,tol). The program should apply the trapezoid rule, doubling the number n of intervals each time, until the estimated error

$$(4/3)|T_{2n}(f) - T_n(f)| < \varepsilon.$$

Use a while loop. The program should be very efficient, never having to compute a value of f(x) more than once. Output should be the last value  $T_{2n}$ , the estimated error and the number n. Apply this function to each of

the integrals (i), (ii), and (iii) of problem 1 with a tolerance of  $\varepsilon = 10^{-6}$ . Compare the estimated error with the actual error.

4. The weights and nodes for 4 point Gaussian quadrature on [-1, 1] are

Nodes t1 = .339981043584856;t2 = .861136311594052;

Weights A1 = .652145154862546;A2 = .347854845137454;

a) Transform to the interval [a, b].

b) Write a function mfile gauss4.m, with call gauss4(f,a,b), that implements Gaussian quadrature on the interval [a, b] for the function f(x), given as an inline function.

c) Apply your Gaussian quadrature rule to the integral (i) of problem 1. Compare your results with trapezoid and Simpson's rule for n = 4.

**5.** Let  $f(x,c) = 1/\sqrt{c + e^x - x}$  and let

$$F(c) = \sqrt{2} - \int_0^3 f(x, c) dx.$$

Use the MATLAB code quad1 to compute F(c) where c is a parameter. You should see that F changes sign between c = 1 and c = 1.5. Use the MATLAB code fzero to find the value of c such that F(c) = 0 with a tolerance of  $10^{-6}$ .

6. Let  $f(x) = \frac{\cos x}{\sqrt{x}}$ . Use the MATLAB routine quad to estimate

$$\int_0^1 f(x) dx.$$

For info on quad, enter help quad. You will see that there are various options. You can specify the tolerance and you can request the routine to display how many functions evaluations are made, and where. The default tolerance is  $10^{-6}$ . If f is given as an inline function, the call is quad(f, a,

b, tol, trace) where trace is any nonzero number. To use the default tolerance, the call is quad(f, a,b,[], trace).

a) Use quad on this integral with the default tolerance. How many function evaluations are needed altogether? How many are needed in the interval [0, b] where  $b = 2.52928585 \times 10^{-10}$ ? How many are needed for the last interval [.724842, 1]?

b) Now make the change of variable  $x = t^2$  in the integral and repeat your calculation with quad. What is the difference? You have "softened the singularity" at x = 0 with the change of variable.