

Lecture 8: The Change of Basis Formula for the Matrix of a Linear Transformation

The Second Change of Basis Formula

Recall that $P_{\mathcal{C} \leftarrow \mathcal{B}}$, the change of basis matrix from the basis \mathcal{B} to the basis \mathcal{C} (so read from right to left) is the matrix whose columns are the basis vectors of \mathcal{B} written out in terms of \mathcal{C} .

Theorem (The second change of basis formula)

Suppose $T : V \rightarrow V$ is a linear transformation and \mathcal{B} and \mathcal{C} are bases of V .

Then

$${}_{\mathcal{C}}[T]_{\mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{B}} {}_{\mathcal{B}}[T]_{\mathcal{B}} P_{\mathcal{B} \leftarrow \mathcal{C}} \quad (*)$$

The whole point of the notation is to make this formula easy to remember. Mnemonic—keep the \mathcal{B} 's together.

Proof. By Proposition (1) of Lecture 7, we have

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = {}_{\mathcal{B}}[I_V]_{\mathcal{C}}$$

and

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = {}_{\mathcal{C}}[I_V]_{\mathcal{B}}.$$

Hence, the right-hand side of (*) becomes

$$\text{RHS} = {}_{\mathcal{C}}[I_V]_{\mathcal{B}} \bullet {}_{\mathcal{B}}[T]_{\mathcal{B}} \bullet {}_{\mathcal{B}}[I_V]_{\mathcal{C}}.$$

Here \bullet is matrix multiplication.

But by Proposition (1) of Lecture 6 (applied twice) we have

$$\begin{aligned}\text{RHS} &= \mathcal{C}[I_V \circ T \circ I_V]_{\mathcal{C}} \\ &= \mathcal{C}[T]_{\mathcal{C}}.\end{aligned}$$



We now do two examples.

Two Solved Change of Basis Problems for Matrices

Problem 1

Let T be the linear transformation for \mathbb{R}^2 to \mathbb{R}^2 whose matrix ${}_{\mathcal{E}}[T]_{\mathcal{E}}$ relative to the standard basis $\mathcal{E} = \{(1, 0), (0, 1)\}$ is

$${}_{\mathcal{E}}[T]_{\mathcal{E}} = A = \begin{pmatrix} a & b \\ c & c \end{pmatrix}$$

Let $\mathcal{C} = \{f_1, f_2\}$ be the new basis for \mathbb{R}^2 given by $f_1 = \frac{e_1 + e_2}{\sqrt{2}}$,

$f_2 = \frac{-e_1 + e_2}{\sqrt{2}}$ (so the old basis e_1, e_2 rotated by 45°).

Find the matrix C of T relative to f_1, f_2 . So we want

$$C = {}_{\mathcal{C}}[T]_{\mathcal{C}}.$$

Problem 1

Solution

There are two steps

1. Compute the change of basis matrices $P_{\mathcal{C} \leftarrow \mathcal{B}}$ and $P_{\mathcal{B} \leftarrow \mathcal{C}}$.
2. Apply the Second Change of Basis Formula from Theorem 2 (pg 13).

Step 1

Put $\mathcal{E} = \text{standard basis} = \{e_1, e_2\}$ (we will use \mathcal{E} instead of \mathcal{B} so you have to replace \mathcal{B} by \mathcal{E} in the formulas) and $\mathcal{C} = \{f_1, f_2\}$.

Basic Principle

It is easy to compute $P_{\mathcal{E} \leftarrow \mathcal{C}}$ for any basis $\mathcal{C} \in \mathbb{R}^n$. Then you compute $P_{\mathcal{C} \leftarrow \mathcal{E}}$ by inverting $P_{\mathcal{E} \leftarrow \mathcal{C}}$.

Problem 1

Computation of $P_{\mathcal{E} \leftarrow \mathcal{C}}$

The change of basis matrix from \mathcal{C} to \mathcal{E} is the matrix whose i -th column is the coordinates of f_i relative to e_1, e_2 so

$$P_{\mathcal{E} \leftarrow \mathcal{C}} = \begin{matrix} e_1 \\ e_2 \end{matrix} \begin{pmatrix} f_1 & f_2 \\ \downarrow & \downarrow \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

So “writing \mathcal{C} in terms of \mathcal{E} ”.

Computation of $P_{\mathcal{C} \leftarrow \mathcal{E}}$

$$P_{\mathcal{C} \leftarrow \mathcal{E}} = (P_{\mathcal{E} \leftarrow \mathcal{C}})^{-1} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

Problem 1

Step 2 Apply the second change of basis formula.

$${}_{\mathcal{C}}[T]_{\mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{E}} {}_{\mathcal{E}}[T]_{\mathcal{E}} P_{\mathcal{E} \leftarrow \mathcal{C}}$$

So,

$$\begin{aligned} C &= P^{-1}AP \\ &= \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \\ &= \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \frac{a+b}{\sqrt{2}} & \frac{-a+b}{\sqrt{2}} \\ \frac{c+d}{\sqrt{2}} & \frac{-c+d}{\sqrt{2}} \end{pmatrix} \\ &= \begin{pmatrix} \frac{a+b+c+d}{2} & \frac{-a-c+b+d}{2} \\ \frac{-a-b+c+d}{2} & \frac{-b-c+a+d}{2} \end{pmatrix} \end{aligned}$$

Problem 2

Problem 2

Let L be the (oriented) line that makes an angle of θ radians with the x -axis. Let r be reflection in the line L . Find the matrix of r relative to the standard basis \mathcal{E} .

Solution

The unit vector $t = (\cos \theta, \sin \theta)$ lies in L and has correct orientation. Use trigonometry.

Problem 2

The vector $n = (-\sin \theta, \cos \theta)$ perpendicular to the line L (see the picture). I use t for “tangent” and n for “normal”.

The reflection r leave the line L fixed and carries the normal vector $(-\sin \theta, \cos \theta)$ to its negative. Think of L as the mirror for r .

Hence

$$r(t) = t$$

and

$$r(n) = -n.$$

Put $\mathcal{C} = \{t, n\}$ (t and n are orthogonal so they are independent).

Problem 2

Hence

$${}_{\mathcal{C}}[r]_{\mathcal{C}} = \begin{matrix} t & n \\ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \end{matrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

We now apply the Second Change of Basis Formula to compute ${}_{\mathcal{E}}[r]_{\mathcal{E}}$.

Step 1 Compute the change of basis matrices $P_{\mathcal{E} \leftarrow \mathcal{C}}$ and $P_{\mathcal{C} \leftarrow \mathcal{E}}$.

$$P_{\mathcal{E} \leftarrow \mathcal{C}} = \begin{matrix} t & n \\ \begin{pmatrix} e_1 & \cos \theta & -\sin \theta \\ e_2 & \sin \theta & \cos \theta \end{pmatrix} \end{matrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

Hence

$$P_{\mathcal{C} \leftarrow \mathcal{E}} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

This matrix has determinant $\det = 1$.

Problem 2

Step 2 By the Second Change of Basis Formula we have:

$$\begin{aligned}[r]_{\mathcal{E}} &= P_{\mathcal{E} \leftarrow \mathcal{C}} \mathcal{C} [r]_{\mathcal{C}} P_{\mathcal{C} \leftarrow \mathcal{E}} \\ &= \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos^2 \theta - \sin^2 \theta & 2 \cos \theta \sin \theta \\ 2 \cos \theta \sin \theta & -(\cos^2 \theta - \sin^2 \theta) \end{pmatrix} \\ &= \begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix} \\ &= \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\end{aligned}$$

In the last line, the first matrix is a rotation matrix and the second matrix is a reflection across the x -matrix.