

# Lecture 6

## Discrete Random Variables and Probability Distributions

Go to "BACKGROUND COURSE NOTES" at the end of my web page and download the file distributions.

Today we say goodbye to the elementary theory of probability and start Chapter 3. We will open the door to the application of algebra to probability theory by introducing the concept of "random variable".

27  
What you will need to get from it (at a minimum) is the ability to do the following "Good Citizen Problems".

I will give you a probability mass function  $p(x)$ . You will be asked to compute

- (i) The expected value (or mean)  $E(X)$ .
- (ii) The variance  $V(X)$ .
- (iii) The cumulative distribution function  $F(x)$ .

You will learn what these words mean shortly.

# Mathematical Definition

3

Let  $S$  be the sample space of some experiment (mathematically a set  $S$  with a probability measure  $P$ ). A random variable  $X$  is a real-valued function on  $S$ .

## Intuitive Idea

A random variable is a function whose values have probabilities attached.

## Remark

To go from the mathematical definition to the "intuitive idea" is tricky and not really that important at this stage.

# The Basic Example

Flip a fair coin three times so

$$S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

Let  $X$  be function on  $X$  given by

$$X = \text{number of heads}$$

so  $X$  is the function given by

$$\begin{array}{cccccccc} \{ & HHH & , & HHT & , & HTH & , & HTT & , & THH & , & THT & , & TTH & , & TTT \} \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ & 3 & & 2 & & 2 & & 1 & & 2 & & 1 & & 1 & & 0 \end{array}$$

What is

$$P(X=0), P(X=3), P(X=1), P(X=2)$$

AnswersNote  $\#(S) = 8$ 

$$P(X=0) = P(TTTT) = \frac{1}{8}$$

$$P(X=1) = P(HTTT) + P(THTT) \\ + P(TTHT) = \frac{3}{8}$$

$$P(X=2) = P(HHTT) + P(HTHT) \\ + P(THTH) = \frac{3}{8}$$

$$P(X=3) = P(HHHH) = \frac{1}{8}$$

We will tabulate these

Value	$x$	0	1	2	3
probability of the value	$P(X=x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Get used to such tabular presentations.

# Rolling a Die

6

Roll a fair die, let

$X =$  the number that comes up

So  $X$  takes values 1, 2, 3, 4, 5, 6 each with probability  $\frac{1}{6}$ .

$x$	1	2	3	4	5	6
$P(X=x)$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

This is a special case of the discrete uniform distribution where  $X$  takes values 1, 2, 3, ...,  $n$  each with probability  $\frac{1}{n}$  (so "roll a fair die with  $n$  faces").

# Bernoulli Random Variable

Usually random variables are introduced to make things numerical. We illustrate this by an important example - page 8. First meet some random variables -

## Definition (The simplest random variable(s))

The actual simplest random variable is a random variable in the technical sense but isn't really random. It takes one value (let's suppose it is 0) with probability one

x	0
$P(X=0)$	1

Nobody ever mentions this because it is too simple - it is deterministic.

8  
The simplest random variable that actually is random takes Two values, let's suppose they are 1 and 0 with probabilities  $p$  and  $q$ . Since  $X$  has to be either 1 or 0 we must have

$$p+q=1.$$

So we get

$x$	1	0
$P(X=x)$	$p$	$q$

This called the Bernoulli random variable with parameter

$p$ . So a Bernoulli random variable is a random variable that takes only two values 0 and 1.



Where do Bernoulli

random variables come from?

We go back to elementary probability.

Definition A Bernoulli

experiment is an experiment which has two outcomes which we call (by convention) "success"  $S$  and failure  $F$ .

Example: Flipping a coin.

We will call a head a success and a tail a failure.

2 Often we call a "success" something that is in fact far from an actual success - e.g. a machine breaking down

By convention we let

$$P(S) = p \text{ and } P(F) = q$$

So again  $p + q = 1$ .

Thus the sample space  $\mathcal{S}$   
of a Bernoulli experiment

is given by

$$\mathcal{S} = \{S, F\}.$$

To join up pages 7 and 9  
We define a random variable

$$X \text{ on } \mathcal{S} \text{ by } X(S) = 1 \text{ and}$$

$$X(F) = 0 \text{ so}$$


$$P(X=1) = P(S) = p \text{ and}$$

$$P(X=0) = P(F) = q$$

# Discrete Random Variables 11

## Definition

A subset  $\mathcal{S}$  of the real line  $\mathbb{R}$  is said to be discrete if for every whole number  $n$  there are only finitely many elements of  $\mathcal{S}$  in the interval  $[-n, n]$



So a finite subset of  $\mathbb{R}$  is discrete but so is the set of integers  $\mathbb{Z}$ .

## Remark

12

The definition in the text is wrong. The set of rational numbers  $\mathbb{Q}$  is countably infinite but is not discrete. This is not important for this course.

## Definition

A random variable is said to be discrete if its set of possible values is a discrete set.

A possible value means a value  $x_0$  so that  $P(X=x_0) \neq 0$ .

## Definition

The probability mass function (abbreviated pmf) of a discrete random variable  $X$  is the function  $p_X$  defined by

$$p_X(x) = P(X=x)$$

We will often write  $p(x)$  instead of  $p_X(x)$ .

## Note

- (i)  $p(x) \geq 0$
- (ii)  $\sum_{\text{all possible } x} p(x) = 1$
- (iii)  $p(x) = 0$  for all  $x$  outside a countable set.


# Graphical Representations of pmf's

There are two kinds of graphical representations of pmf's, the "line graph" and the "probability histogram".


We will illustrate them with the Bernoulli distribution with parameter  $p$ .

$x$	1	0
$P(X=x)$	$p$	$q$

table



line graph





histogram

We also illustrate these  
for the basic example (pg. 5)

$x$	0	1	2	3
$P(X=x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

table



# The Cumulative Distribution Function <sup>16</sup>

The cumulative distribution function  $F_X$  (abbreviated cdf) of a discrete random variable  $X$  is defined by

$$F_X(x) = P(X \leq x)$$

We will often write  $F(x)$  instead of  $F_X(x)$ .

## Bank account analogy

Suppose you deposit \$1000 at the beginning of every month






The "line graph" of your deposits is on the previous page. We will use  $t$  (time as your variable). Let

$F(t)$  = the amount you have accumulated at time  $t$

What does the graph of  $F$  look like?



It is critical to observe that whereas the deposit function on page 15 is zero for all real numbers except 12 the cumulation function is never zero between 1 and  $\infty$ .

You would be very upset if you walked into the bank on July 5<sup>th</sup> and they told you your balance was zero - you never took any money out.

Once your balance was non zero it was never zero thereafter.

# Back to Probability

19

The cumulative distribution  $F(x)$  is "the total probability you have accumulated when you get to  $x$ ". Once it is nonzero it is never zero again

( $p(x) \geq 0$  means "you never take any probability out").

To write out  $F(x)$  in formulas you will need


several (many) formulas. There

should never be EQUALITIES

in your formulas only INEQUALITIES

# The cdf for the Basic Example 20


We have



line graph  
of  $p$

So we start accumulation probability  
at  $x=0$

## Ordinary Graph of F



## Formulas for F

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{1}{8} & 0 \leq x < 1 \\ \frac{4}{8} & 1 \leq x < 2 \\ \frac{7}{8} & 2 \leq x < 3 \\ 1 & 3 \leq x \end{cases}$$

be  
careful

You can see you have to  
be careful about the inequalities  
on the right-hand side.

20

## Expected Value

### Definition

Let  $X$  be a discrete random variable with set of possible values  $D$  and pmf  $p(x)$ . The expected value or mean value of  $X$  denoted  $E(X)$  or  $\mu$  is defined by

↓  
Greek letter mu

$$E(X) = \sum_{x \in D} x P(X=x) = \sum_{x \in D} x p(x)$$

## Remark

$E(X)$  is the whole point for monetary games of chance e.g. lotteries, blackjack, slot machines.

If  $X =$  your payoff, the operators of these games make sure  $E(X) < 0$ . Thorp's card-counting strategy in blackjack changed  $E(X) < 0$  (because ties went to the dealer) to  $E(X) > 0$  to the dismay of the casinos. See "How to Beat the Dealer" by Edward Thorp (a math professor at UC Irvine)

# Examples

The expected value of the Bernoulli distribution

$$E(X) = \sum_x x P(X=x) = (0)(q) + (1)(p)$$

$$= p$$

The expected value for the basic example (so the expected number of heads)

$$E(X) = (0)\left(\frac{1}{8}\right) + (1)\left(\frac{3}{8}\right) + (2)\left(\frac{3}{8}\right)$$

$$+ (3)\left(\frac{1}{8}\right)$$

$$= \frac{3}{2}$$

∑ The expected value is NOT the most probable value

For the basic example the possible values of  $X$  were 0, 1, 2, 3 so  $\frac{3}{2}$  was not even a possible value

$$P(X = \frac{3}{2}) = 0$$

The most probable values were 1 and 2 (tied) each with probability  $\frac{3}{8}$ .

## Rolling a Die

$$\begin{aligned} E(X) &= (1)\left(\frac{1}{6}\right) + (2)\left(\frac{1}{6}\right) + (3)\left(\frac{1}{6}\right) \\ &\quad + (4)\left(\frac{1}{6}\right) + (5)\left(\frac{1}{6}\right) + (6)\left(\frac{1}{6}\right) \\ &= \frac{1}{6} [1+2+3+4+5+6] = \frac{1}{6} \frac{(7)(6)}{2} \\ &= \frac{7}{2}. \end{aligned}$$



# Variance

The expected value does not tell you everything you want to know about a random variable (how could it, it is just one number).

Suppose you and a friend play the following game of chance. Flip a coin. If a head comes up you get \$1 if a tail comes up you pay your friend \$1. So

If  $X =$  your payoff

$$X(H) = +1 \quad , \quad X(T) = -1$$

$$E(X) = (+1)\left(\frac{1}{2}\right) + (-1)\left(\frac{1}{2}\right) = 0$$

so this is a fair game.

Now suppose you play the game changing \$1 to \$1000. It is still a fair game

$$E(X) = (1000)\left(\frac{1}{2}\right) + (-1000)\left(\frac{1}{2}\right) \\ = 0$$

but I personally would be very reluctant to play this game.

The notion of variance is designed to capture the difference between the two games.

## Definition

Let  $X$  be a discrete random variable with set of possible values  $D$  and expected value  $\mu$ . Then the variance of  $X$ , denoted  $V(X)$  or  $\sigma^2$  (sigma squared), is defined by

$$V(X) = \sum_{x \in D} (x - \mu)^2 P(X=x)$$

$$= \sum_{x \in D} (x - \mu)^2 p(x) \quad (*)$$

The standard deviation  $\sigma$  of  $X$  is defined to be the square-root of the variance

$$\sigma = \sqrt{V(X)} = \sqrt{\sigma^2}$$

Check that for the two games above (with your friend)

$$\sigma = 1 \text{ for the } \$1 \text{ game}$$

$$\sigma = 1000 \text{ for the } \$1000 \text{ game.}$$

### The Shortcut Formula for $V(X)$

The number of arithmetic operations (subtractions) necessary to compute  $\sigma^2$  can be greatly reduced by using

### Proposition

$$(i) \quad V(X) = E(X^2) - E(X)^2$$

or

$$(ii) \quad V(X) = \sum_{x \in D} x^2 p(x) - \mu^2$$

In the formula (\*) you need  $\#(D)$  subtractions (for each  $x \in D$  you have to subtract  $\mu$  then square ...). For the shortcut formula you need only one. Always use the shortcut formula.

Remark Logically, version (i) of the shortcut formula is not correct because we haven't yet defined the random variable  $X^2$ . We will do this soon - "change of random variable".

# Example (The fair die)

$X$  = outcome of rolling a die

We have seen (pg 24)

$$E(X) = \mu = \frac{7}{2}$$

$$E(X^2) = (1)^2 \left(\frac{1}{6}\right) + (2)^2 \left(\frac{1}{6}\right) + (3)^2 \left(\frac{1}{6}\right) \\ + (4)^2 \left(\frac{1}{6}\right) + (5)^2 \left(\frac{1}{6}\right) + (6)^2 \left(\frac{1}{6}\right)$$

$$= \frac{1}{6} [1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2]$$

$$= \frac{1}{6} [91] \quad \swarrow \text{later}$$

so

$$E(X^2) = \frac{91}{6}$$

don't forget  
to square  $\mu$

Hence

$$V(X) = \frac{91}{6} - \left(\frac{7}{2}\right)^2 = \frac{91}{6} - \frac{49}{4}$$

i) How did I know

$$1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 = 91$$

This because

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

Now plug in  $n=6$ .

ii) In the formula for

$E(X^2)$  don't square

the probabilities

NOT squared

$$E(X^2) = (1^2)(\frac{1}{6}) + (2^2)(\frac{1}{6}) + \dots$$

first value squared

second value squared