

Lecture 18 : Pairs of Continuous Random Variables


Definition

Let X and Y be continuous random variables defined on the same sample space S . Then the joint probability density function, joint pdf, $f_{X,Y}(x, y)$ is the function such that

$$P((X, Y) \in A) = \underbrace{\iint_A f_{X,Y}(x, y) dx dy}_{\text{double integral}} \quad (*)$$

for any region A in the plane.

Again the geometric interpretation of (*) is very important



$P((X, Y) \in A) =$ the volume under the graph of f and above the region A .

For $f(x, y)$ to be a joint *pdf* for some pair of random variables X and Y it is necessary and sufficient that

$$f(x, y) \geq 0, \quad \text{all } x, y$$

and

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$$

or geometrically, the total volume under the graph of f has to be 1.

Example 5.3 (from text)


A bank operates a drive-up window and a walkup window. On a randomly selected day, let

X = proportion of time the
drive-up facility is in use.

Y = proportion of time the
walk-up facility is in use.

The set of possible outcomes for the pair (X, Y) is the square

$$R = \{(x, y), 0 \leq x \leq 1, 0 \leq y \leq 1\}$$



Suppose the joint *pdf* of (X, Y) is given by

$$f_{x,y}(x, y) = \begin{cases} 6/5(x + y^2), & 0 \leq x \leq 1 \\ & 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find the probability that neither facility is in use more than $1/4$ of the time.

Solution

Neither facility is in use more than $\frac{1}{4}$ of the time when re-expressed in terms of X and Y is

$$X \leq \frac{1}{4} \left(\text{the drive-up facility is in use} \leq \frac{1}{4} \text{ of the time} \right)$$

and

$$Y \leq \frac{1}{4} \left(\text{the walk-up facility is in use} \leq \frac{1}{4} \text{ of the time} \right)$$

Solution (Cont.)


The author formulated the problem in a confusing fashion, don't worry about it.
So we want

$$P\left(0 \leq X \leq \frac{1}{4}, 0 \leq Y \leq \frac{1}{4}\right)$$

or

$$P((X, Y) \in S)$$

where S is the small square



This probability is given by

$$\int_0^{\frac{1}{4}} \int_0^{\frac{1}{4}} \frac{6}{5}(x + y^2) dx dy$$
$$\iint_S \frac{6}{5}(x + y^2) dx dy \quad (\#)$$

Remark

For general (X, Y) we have

$$\begin{aligned} P(a \leq X \leq b, c \leq Y \leq d) \\ = \int_a^b \int_c^d f_{X,Y}(x, y) dx dy \end{aligned}$$

Let's do the integral (#). We will do the x -integration first. So

$$\begin{aligned} P\left(0 \leq X \leq \frac{1}{4}, 0 \leq Y \leq \frac{1}{4}\right) \\ = \int_0^{\frac{1}{4}} \left(\int_0^{\frac{1}{4}} \frac{6}{5} (x + y^2) dx \right) dy \\ = \frac{6}{5} \int_0^{\frac{1}{4}} \left(\frac{X^2}{2} + xy^2 \right) \Big|_{x=0}^{x=\frac{1}{4}} dy \end{aligned}$$

Remark (Cont.)

$$\begin{aligned} &= \frac{6}{5} \int_0^{\frac{1}{4}} \left(\frac{1}{32} + \frac{y}{4} \right) dy \\ &= \frac{6}{5} \left[\left(\frac{y}{32} + \frac{y^2}{8} \right) \Big|_{y=0}^{y=\frac{1}{4}} \right] \\ &= \frac{6}{5} \left[\frac{1}{128} + \frac{1}{(64)(12)} \right] \\ &= \left(\frac{6}{5} \right) \left(\frac{1}{64} \right) \left(\frac{1}{2} + \frac{1}{12} \right) \\ &= \left(\frac{6}{5} \right) \left(\frac{1}{64} \right) \left(\frac{7}{12} \right) \\ &= \frac{7}{640} \end{aligned}$$

An exercise in the forgotten art of fractions- more of the same later.

More Theory Marginal Distributions in the Continuous Case

Problem

Suppose you know the joint pdf $f_{X,Y}(x, y)$ of (X, Y) . How do you find the individual pdf's $f_X(x)$ of X and $f_Y(y)$. The answer is

Proposition

$$\begin{aligned} \text{(i)} \quad f_X(x) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy \\ \text{(ii)} \quad f_Y(y) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx \end{aligned} \quad (*)$$

Proposition (Cont.)

The formula () is the continuous analogue of the formula for the discrete case. Namely*

Discrete Case

$$f_X(x) = \sum_{\text{all } y} f_{X,Y}(y)$$

Continuous Case

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$

In the first case we sum away the “extra variable” y and in the second case we integrate it away.

By analogy once again we call $f_X(x)$ and $f_Y(y)$ (obtained via (*)) the marginal densities or marginal *pdf*'s.

Note the $f_X(x)$ and $f_Y(y)$ are the two partial definite integrals of $f_{X,Y}(x, y)$ - see Lecture 16.

Example 5.4

We compute the two marginal *pdf*'s for the bank problem, Example 5.3.


$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$
$$= \begin{cases} \int_0^1 \frac{6}{5}(x + y^2) dy, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

↑
this is a little tricky.

The formula for $f_X(x)$ says you integrate $f_{X,Y}(x, y)$ over the vertical

line passing through x .

If x does not satisfy $0 \leq x \leq 1$ then the vertical line does not pass through the square R where $f_{X,Y}(x, y)$ is non zero



You get $f_X(2)$ by integrating over the line $x = 2$ above which $f_{X,Y}(x, y) = 0$.
Equivalently (without geometry)

$$f_X(2) = \int_{-\infty}^{\infty} f_{X,Y}(2, y) dy = \int_{-\infty}^{\infty} 0 dy = 0$$

Now we finish the job

$$\begin{aligned}\int_0^1 \frac{6}{5}(x + y^2)dy &= \frac{6}{5} \int_0^1 (x + y^2)dy \\ &= \frac{6}{5} \left(xy + \frac{y^3}{3} \right) \Big|_{y=0}^{y=1} = \frac{6}{5} \left(x + \frac{1}{3} \right)\end{aligned}$$

Similarly

$$\begin{aligned}f_Y(y) &= \begin{cases} \frac{6}{5} \int_0^1 \frac{6}{5}(x + y^2)dx, & 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases} \\ &= \begin{cases} \frac{6}{5}y^2 + \frac{3}{5}, & 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}\end{aligned}$$

Independence of Two Continuous Random Variables

Definition

Two continuous random variables X and Y are independent if their joint pdf $f_{X,Y}(x, y)$ is the product of the two marginal pdf's $f_X(x)$ and $f_Y(y)$ so

$$f_{X,Y}(x, y) = f_X(x)f_Y(y)$$

This not true for the bank example pg. 5.

$$f_{X,Y}(x, y) = \frac{6}{5}(x + y^2)$$

↑
not a product

Covariance and Correlation of Pairs of Continuous Random Variables

We continue with a pair of continuous random variables X and Y as before.

Again we define

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$$

and

$$\rho_{X,Y} = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y}$$

But now

$$E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy$$

We will now compute the $\text{Cov}(X, Y)$ and $\text{Corr}(X, Y)$ for the bank problem. So

$$f_{X,Y}(x, y) = \begin{cases} \frac{6}{5}(x + y^2), & 0 \leq x \leq 1 \\ & 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$
$$f_X(x) = \begin{cases} \frac{6}{5}\left(x + \frac{1}{3}\right), & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$
$$f_Y(y) = \begin{cases} \frac{6}{5}y^2 + \frac{3}{5}, & 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Let's first do the calculations for X and Y - we need

$$E(X), E(Y), \sigma_X = \sqrt{V(X)} \quad \text{and} \quad \sigma_Y = \sqrt{V(Y)}$$

$$\begin{aligned}
 E(X) &= \int_0^1 x \frac{6}{5} \left(x + \frac{1}{3} \right) dx \\
 &= \frac{6}{5} \int_0^1 \left(x^2 + \frac{x}{3} \right) dx = \frac{6}{5} \left(\frac{x^3}{3} + \frac{x^2}{6} \right) \Big|_{x=0}^{x=1} \\
 &= \frac{6}{5} \left(\frac{1}{3} + \frac{1}{6} \right) = \frac{6}{5} \left(\frac{3}{6} \right) = \frac{3}{5}
 \end{aligned}$$

$$\begin{aligned}
 E(X^2) &= \int_0^1 x^2 \frac{6}{5} \left(x + \frac{1}{3} \right) dx \\
 &= \frac{6}{5} \int_0^1 \left(x^3 + \frac{x^2}{3} \right) dx = \frac{6}{5} \left(\frac{x^4}{4} + \frac{x^3}{9} \right) \Big|_{x=0}^{x=1} \\
 &= \frac{6}{5} \left(\frac{1}{4} + \frac{1}{9} \right) = \frac{6}{5} \left(\frac{13}{36} \right) = \frac{13}{30}
 \end{aligned}$$

$$V(X) = \frac{13}{30} - \left(\frac{3}{5} \right)^2 = \frac{13}{30} - \frac{9}{25} = \frac{65 - 54}{150} = \frac{11}{150}$$

$$\sigma_X = \sqrt{\frac{11}{150}} = \frac{1}{5} \sqrt{\frac{11}{6}}$$

$$\begin{aligned}
 E(Y) &= \int_y^1 \left(\frac{6}{5}y^2 + \frac{3}{5} \right) dy \\
 &= \frac{6}{5} \int_0^1 y^3 dy + \frac{3}{5} \int_0^1 y dy \\
 &= \left(\frac{6}{5} \right) \left(\frac{1}{4} \right) + \left(\frac{3}{5} \right) \left(\frac{1}{2} \right) = \frac{6}{20} + \frac{3}{10} = \frac{12}{20}
 \end{aligned}$$

$$\begin{aligned}
 E(Y^2) &= \int_0^1 y^2 \left(\frac{6}{5}y^2 + \frac{3}{5} \right) dy \\
 &= \frac{6}{5} \int_0^1 y^4 dy + \frac{3}{5} \int_0^1 y^2 dy \\
 &= \left(\frac{6}{5} \right) \left(\frac{1}{5} \right) + \left(\frac{3}{5} \right) \left(\frac{1}{3} \right) = \frac{6}{25} + \frac{1}{5} = \frac{11}{25}
 \end{aligned}$$

$$V(Y) = \frac{11}{25} - \frac{144}{400} = \frac{176}{400} - \frac{144}{400} = \frac{32}{400} = \frac{2}{25}$$

$$\sigma_Y = \sqrt{\frac{2}{25}} = \frac{1}{5} \sqrt{2}$$

Finally we need

$$\begin{aligned} E(XY) &= \int_0^1 \int_0^1 (xy) \frac{6}{5} (x + y^2) dx dy \\ &= \int_0^1 \int_0^1 \underbrace{xy \frac{6}{5} x}_{\text{product function}} dx dy + \int_0^1 \int_0^1 \underbrace{xy \frac{6}{5} y^2}_{\text{product function}} dx dy \\ &= \frac{6}{5} \left(\int_0^1 x^2 dx \right) \left(\int_0^1 y dy \right) + \frac{6}{5} \left(\int_0^1 x dx \right) \left(\int_0^1 y^3 dy \right) \\ &= \left(\frac{6}{5} \right) \left(\frac{1}{3} \right) \left(\frac{1}{2} \right) + \left(\frac{6}{5} \right) \left(\frac{1}{2} \right) \left(\frac{1}{4} \right) \\ &= \left(\frac{6}{5} \right) \left(\frac{1}{2} \right) \left(\frac{1}{3} + \frac{1}{4} \right) = \left(\frac{6}{5} \right) \left(\frac{1}{2} \right) \left(\frac{7}{12} \right) = \frac{7}{20} \end{aligned}$$

Now we can mop the fruits of our labours.

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$$

$$= \frac{7}{20} - \left(\frac{3}{5}\right)\left(\frac{12}{20}\right)$$

$$= \frac{7}{20} - \frac{36}{100} = \frac{35}{100} - \frac{36}{100}$$

$$\text{Cov}(X, Y) = \frac{-1}{100}$$

$$\text{Cov}(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} = \frac{\frac{-1}{100}}{\left(\frac{1}{5} \sqrt{\frac{11}{6}}\right) \left(\frac{1}{5} \sqrt{2}\right)}$$

$$= \left(\frac{-1}{100}\right) \left(\frac{\cancel{5}}{\sqrt{\frac{11}{\cancel{6}}}}\right) \left(\frac{\cancel{5}}{\sqrt{2}}\right) = -\frac{1}{4} \left(\frac{1}{\sqrt{\frac{11}{3}}}\right) = -\frac{\sqrt{3}}{4\sqrt{11}}$$

Independence of Continuous Random Variables

Definition

Two continuous random variables X and Y are independent if the joint pdf is the product of the two marginal pdf's

$$f_{X,Y}(x, y) = f_X(x)f_Y(y)$$

(so \iff the joint pdf is a product function)

So in Example 5.3, page 4, X and Y are NOT independent.