# Critical thresholds in pressureless Euler–Poisson equations with background states

Young-Pil Choi, Dong-ha Kim, Dowan Koo, and Eitan Tadmor

**Abstract.** We investigate the critical threshold phenomena in a large class of one-dimensional pressureless Euler–Poisson (EP) equations, with non-vanishing background states. First, we establish local-in-time well-posedness in proper regularity spaces specifically involving  $\dot{W}^{-1,p}$ , which are adapted for a *neutrality condition* to hold. This negative homogeneous Sobolev regularity is shown to be necessary: we prove an ill-posedness result in classical Sobolev spaces  $H^{s}(\mathbb{R})$  in the absence of this negative Sobolev regularity. Next we study the critical threshold phenomena in the neutrality condition can further restrict the sub-critical region into its borderline, namely, the sub-critical region is reduced to a single line in the phase plane. We then turn to providing a rather definitive answer for the critical thresholds in the case of repulsive EP systems with variable backgrounds. As an application, we analyze the critical thresholds for the damped EP system for cold-plasma-ion dynamics, where the density of electrons is given by the *Maxwell–Boltzmann relation*.

# 1. Introduction and statement of main results

# 1.1. Systems and notions

In this paper we are interested in the analysis of critical threshold phenomena in a large class of pressureless Euler–Poisson (EP) equations in one dimension. More precisely, the main purpose of our work is to propose a new method based on Lyapunov functions to investigate the super-critical region with finite-time breakdown and the sub-critical region with global regularity of  $C^1$  solutions to the following equations:

$$\begin{cases} \partial_t \rho + \partial_x (\rho u) = 0, \\ \partial_t u + u \partial_x u = -\nu u + kE, \\ \partial_x E = \rho - c, \ E(t, -\infty) = 0, \end{cases} (t, x) \in (0, T) \times \mathbb{R}, \tag{1.1}$$

subject to initial data

$$(\rho, u)(0, x) = (\rho_0, u_0)(x), \quad x \in \mathbb{R}.$$
 (1.2)

Mathematics Subject Classification 2020: 35Q35 (primary); 35B30, 76N10 (secondary).

*Keywords:* pressureless Euler–Poisson system, non-vanishing background states, neutrality condition, critical thresholds, well-posedness, non-existence.

Here,  $\rho = \rho(t, x)$ , u = u(t, x), and E = E(t, x) are respectively, the density, velocity, and forcing of the flow. The parameter  $v \ge 0$  indicates the strength of damping. The parameter k on the right of  $(1.1)_2$  is a physical constant, signifying the underlying forcing of the system – either an *attractive* or *repulsive* forcing depending on whether k < 0 or, respectively, k > 0; for simplicity, we use the rescaled parameter  $k = \pm 1$  throughout the paper.

Equation (1.1)<sub>3</sub> involves a positive background state, c = c(t, x), which is assumed to be uniformly bounded away from vacuum,

$$0 < c_{-} \leq c(t, x) \leq c_{+}, \quad \forall (t, x) \in \mathbb{R}_{+} \times \mathbb{R}$$

to vary smoothly in a manner specified in Theorem 1.1 below. Finally, we augment (1.1) with the following *neutrality condition*:

$$\int_{\mathbb{R}} (\rho(t, x) - c(t, x)) \, \mathrm{d}x = 0.$$
(1.3)

This neutrality condition plays a central role in our discussion. Accordingly, we define a *classical solution* to be a pair,  $(\rho, u)$ , satisfying both the system (1.1) pointwise, and the neutrality condition (1.3). Theorem 1.1 below establishes a proper regularity class in which the neutrality condition (1.3) is guaranteed to hold for local classical solutions. We note that the vanishing condition for E in  $(1.1)_3$  is equivalent to  $E(t, \infty) = 0$  under the neutrality condition. Consequently, the EP system (1.1), equipped with the neutrality (1.3), imposes the requirement that the forcing term E decays at both spatial ends.

#### 1.2. History and review

The system (1.1) is one of the fundamental fluid models describing the dynamics of many important physical flows, including charge transport, plasma waves, and cold ions (k > 0), as well as collapse of stars due to self-gravitation (k < 0); see [2, 3, 8, 12, 13, 16, 26] and references therein.

A mathematical study of finite-time loss of regularity (or singularity formation as shocks) and global-in-time regularity of solutions to Euler equations with non-local interaction forces are by now well established. Since the literature-related results are too vast to be mentioned here, we focus on results for the EP systems. The critical thresholds in the one-dimensional pressureless EP system (1.1) with  $v \ge 0$  and  $c = \bar{c} \ge 0$  are studied in [13]. The damped EP system or EP system with the quadratic confinement is also taken into account in [5, 10]. In the multi-dimensional case, the critical thresholds are investigated for the restricted EP system in [23, 24] and EP system with radial symmetry in [6, 29, 30].

The case with pressure, i.e., the pressure term  $p(\rho) = \rho^{\gamma}$  with  $\gamma \ge 1$ , is added to the EP system, but with c = 0 and  $\nu = 0$ , is also studied in [28]. The critical thresholds for the EP system with pressure and non-zero background state are a challenging open problem. We refer to [14, 17, 19] for the global-in-time existence of solutions with small amplitude to the EP system for electrons. The critical thresholds for the one-dimensional Euler-alignment

or Euler–Poisson-alignment systems are analyzed in [7, 9, 27]. The finite-time singularity formation for the one-dimensional EP system for cold ions is also discussed in [1, 22].

Beyond those developments, there is little literature on critical thresholds in EP systems with *variable* background. In this context, we mention the recent work [4] in which critical threshold conditions in the system (1.1) with the attractive forces are investigated. In order to handle new difficulties arising from the consideration of the variable background, a combination of phase plane analysis and a comparison principle is used in [4]. However, as mentioned in [4], that strategy would not work in the presence of repulsive forces.

#### 1.3. Main results

In this section, we present our main results on the Euler–Poisson equation with background states. The purpose of this work is fourfold:

- (i) We present a new framework for the well-posedness theory for the EP system with the background, which includes a rigorous treatment of the neutrality condition (1.3), while justifying this framework by presenting concrete examples;
- (ii) we point out that in the attractive case, the set of relevant configurations that yield global smooth solutions – with variable and even with constant background – is, in fact, restricted to the borderline case due to the neutrality condition;
- (iii) we present an in-depth threshold analysis for global solutions of the repulsive EP system with variable background; and
- (iv) as an application of the repulsive case, we study the case of cold plasma corresponding to exponential background  $c = e^{\phi}$ , where  $\phi$  is the potential of the flow.

Below, we present the main results in more detail.

**1.3.1. Local well-posedness and neutrality condition.** We first develop a local-in-time well-posedness theory for the EP system with a non-vanishing background state (1.1). The EP system with zero background has been studied by many authors, and in particular, we refer to the  $H^{s}(\mathbb{R})$  well-posedness in [20, appendix]. However, we cannot specify an appropriate reference for the local-in-time well-posedness theory for the EP system with *non-vanishing* background, (1.1).

Here, the neutrality condition, (1.3), implies the non-integrability  $\rho(t, \cdot) \notin L^1(\mathbb{R})$ , in order to match the infinite mass that "enters" the system from infinity,  $\int_{\mathbb{R}} c(t, \cdot) dx = \infty$ . Consequently, the local well-posedness of (1.1) requires a careful study of proper regularity space, beyond  $H^s(\mathbb{R})$ .

To this end, we define the following function space:

$$X^{p,s} := \{ (f,g) : f \in \dot{W}^{-1,p} \cap H^s(\mathbb{R}), g \in L^p(\mathbb{R}), \ \partial_x g \in H^s(\mathbb{R}) \}$$

with the norm

$$\|\{f,g\}\|_{X^{p,s}} := \|f\|_{\dot{W}^{-1,p}} + \|f\|_{H^s} + \|g\|_{L^p} + \|\partial_x g\|_{H^s}.$$

For p = 2, this simplifies to

$$\|\{f,g\}\|_{X^{2,s}} = \|f\|_{\dot{H}^{-1}\cap H^s} + \|g\|_{H^{s+1}}.$$

We assume  $1 to provide generality. Throughout the paper, we fix <math>s \ge 2$ .

**Theorem 1.1** (Local well-posedness). Let  $1 . Suppose that the initial data <math>(\rho_0, u_0)$  satisfy

$$(\rho_0 - c_0, u_0) \in X^{p,s},$$

where c = c(t, x) is given as

$$c(t,x) = \bar{c} + \eta(t,x) \quad \text{with } \eta \in \mathcal{C}^0([0,\infty); H^s \cap \dot{W}^{-1,p}(\mathbb{R}))$$
(1.4)

for some  $\bar{c} > 0$ . Then there exists a positive constant T > 0 such that the system (1.1)–(1.2) admits a unique classical solution ( $\rho$ , u) satisfying

$$(\rho(t, \cdot) - c(t, \cdot), u(t, \cdot)) \in \mathcal{C}^0([0, T]; X^{p,s})$$

and the neutrality condition (1.3). Moreover, if the solutions  $\rho$  and  $\partial_x u$  are bounded over the time interval [0, T], i.e.,

$$|||\{\rho, u\}||_{[0,T]} := \sup_{0 \le t \le T} (||\rho(t)||_{L^{\infty}} + ||\partial_x u(t)||_{L^{\infty}}) < \infty,$$

then there is propagation of higher-order regularity of  $(\rho, u)$ ,

$$\sup_{0 \le t \le T} \|\{\rho(t, \cdot) - c(t, \cdot), u(t, \cdot)\}\|_{X^{p,s}} \le (\|\{\rho_0 - c_0, u_0\}\|_{X^{p,s}} + (T+1)\|\eta\|_{L^{\infty}(0,T;H^s \cap \dot{W}^{-1,p})})e^{MT},$$
(1.5)

where  $M \sim ||| \{\rho, u\} ||_{[0,T]} + \bar{c} + 1$ .

**Remark 1.2.** The inclusion of  $\dot{W}^{-1,p}$  regularity is crucial for ensuring the propagation of the neutrality condition. Indeed, solutions constructed in Theorem 1.1 satisfy the neutrality condition (1.3). By Lemma A.1, any  $f \in L^2 \cap \dot{W}^{-1,p}(\mathbb{R})$  satisfies

$$\int_{\mathbb{R}} f(x) \, \mathrm{d}x = 0$$

in the sense of an improper integral. Thus,  $\rho - c \in \mathcal{C}^0([0, T]; H^s \cap \dot{W}^{-1, p}(\mathbb{R}))$  ensures the neutrality condition (1.3) is satisfied for  $t \in [0, T]$ . Moreover, the condition on  $\eta$  in (1.4) preserves *mass fluctuation* of c(t, x),

$$\int_{\mathbb{R}} (c(t,x) - c_0(x)) \, \mathrm{d}x = 0, \quad \forall t \in [0,\infty),$$

so that the neutrality condition is compatible with mass conservation of  $\rho$ .

**Remark 1.3.** The family of  $X^{p,s}$  spaces provides an (almost) optimal framework for the well-posedness theory of EP system (1.1), while satisfying the neutrality condition (1.3). By interpolation, we note that

$$p_1 < p_2 \implies X^{p_1,s} \subset X^{p_2,s}$$

However, the endpoint space  $X^{\infty,s}$  cannot guarantee the neutrality condition to hold. For instance,

$$\rho_0 - c_0 := e^{-x^2} \in \dot{W}^{-1,\infty} \cap H^s(\mathbb{R}), \quad \int_{\mathbb{R}} (\rho_0 - c_0) \, \mathrm{d}x > 0.$$

The  $\dot{W}^{-1,p}$  regularity of  $\rho - c$  must correspond to the  $L^p$  regularity of u for the wellposedness defined in  $X^{p,s}$  space. This correspondence is crucial for the theory to hold. To highlight the necessity of negative homogeneous Sobolev regularity, we present the following ill-posedness result.

**Theorem 1.4** (Ill-posedness). For the EP system (1.1)–(1.2) with  $c = \bar{c} > 0$ , there exists  $(\rho_0, u_0)$  satisfying

$$(\rho_0 - \overline{c}, u_0) \in H^s(\mathbb{R}) \times H^{s+1}(\mathbb{R})$$

and

$$\int_{\mathbb{R}} (\rho_0(x) - \bar{c}) \, \mathrm{d}x = 0$$

such that the classical solution  $(\rho, u)$  to (1.1)–(1.2) with initial data  $(\rho_0, u_0)$  satisfies

$$u(t) \notin L^2(\mathbb{R}), \quad \forall t \in (0, \delta]$$

for some  $\delta > 0$ . In particular, the EP system (1.1)–(1.2) is ill posed in  $H^s \times H^{s+1}(\mathbb{R})$ .

The intuition behind this ill-posedness result lies in the behavior of functions in  $L^p$  spaces with large p. Such functions exhibit slower decay at the far field. By choosing  $\rho_0 - \bar{c} \in \dot{W}^{-1,p}(\mathbb{R})$  with p > 2, we can only expect  $u(t) \in L^p(\mathbb{R})$ , which excludes  $L^2(\mathbb{R})$  as soon as t > 0.

**1.3.2. The case of attractive forcing.** In this part, we discuss how the neutrality condition affects the critical threshold phenomenon of the attractive EP system, which is considered to be irrelevant. The threshold conditions for the attractive EP system were derived earlier in [4, 10, 13]. The sub-critical condition for the existence of a global classical solution of the attractive EP system is given by

$$\left(\frac{-\nu + \sqrt{\nu^2 + 4c_-}}{2}\right) u'_0(x) \ge (\rho_0(x) - c_-), \quad \forall x \in \mathbb{R},$$
(1.6)

where  $c_- := c > 0$  for the constant [13, Theorem 3.2], [4, Theorem 2.5] and  $c_- := \min c(\cdot, \cdot) > 0$  for the variable background case [4, Theorem 2.2]. In particular, this threshold is sharp for the constant background case; if there exists  $x \in \mathbb{R}$  which fails

to satisfy the inequality in (1.6), then the solution forms a singularity in a finite time. In the following theorem, however, we find that the sub-critical region should be further restricted to its borderline zone – initial configurations satisfying *equality* in (1.6), in order for the neutrality condition to be fulfilled.

**Theorem 1.5** (Reduction to the borderline). Suppose that  $(\rho, u)$  is a global-in-time classical solution of the attractive EP system (1.1)–(1.2)–(1.3), subject to initial data satisfying (1.6). Then the set of admissible  $c(\cdot, \cdot)$ 's is necessarily restricted to the constant background, namely  $-c(\cdot, \cdot) = c_{-}$  and (1.6) is reduced the borderline case

$$\left(\frac{-\nu + \sqrt{\nu^2 + 4c_-}}{2}\right) u_0'(x) = (\rho_0(x) - c_-), \quad \forall x \in \mathbb{R}.$$
 (1.7)

**Remark 1.6.** Theorem 1.5 shows that in order to entertain the neutrality condition, the sub-critical criteria for variable backgrounds, (1.6), are too restrictive in the sense that they will rule out all but the borderline cases. In particular, for the constant background case  $c = c_{-}$ , the solution ( $\rho$ , u) of a neutrality-satisfying attractive EP system has global-in-time regularity if and only if its initial data ( $\rho_0$ ,  $u_0$ ) satisfies (1.7).

This reduction to the borderline is a main theme in the attractive case. We demonstrate this in the context of variable background, c = c(t, x). Now assume that  $|u_0(x)| \xrightarrow{|x| \to \infty} 0.^1$  We integrate (1.6) to find

$$\left(\frac{-\nu + \sqrt{\nu^2 + 4c_-}}{2}\right) u_0(x) \Big|_{-R}^R \ge \int_{-R}^R (\rho_0(x) - c_0(x)) \,\mathrm{d}x + \int_{-R}^R (c_0(x) - c_-) \,\mathrm{d}x.$$
(1.8)

By assumption,  $u_0(x)|_{-R}^R$  vanishes as  $R \uparrow \infty$  (in fact, equals zero in the periodic case) and the neutrality condition yields  $\int_{\mathbb{R}} (c_0(x) - c_-) dx = 0$ , which implies  $c_0(x) = c_-$  and we get equality at  $(1.8)_{R\to\infty}$ . By assumption, both sides tend to zero with  $R \uparrow \infty$ , hence the equality (1.6) takes place almost everywhere. Indeed, for initial data satisfying (1.6), we have

$$\left(\frac{-\nu + \sqrt{\nu^2 + 4c_-}}{2}\right)\partial_x u(t, x) \ge (\rho(t, x) - c_-), \quad \forall x \in \mathbb{R},$$

so that we can repeat the previous argument for each t > 0; in particular, we get  $c(\cdot, \cdot) = c_-$ . In Section 3 we recover the critical threshold results for attractive EP systems with constant/variable backgrounds via the Lyapunov function method. It captures the solution trajectories in a geometric way so that it can be more intuitively understood than the previous approaches. More importantly, a notable feature of the Lyapunov-based approach is that it goes beyond the attractive forcing case; it can be employed in obtaining the critical thresholds for the repulsive EP systems, in particular, with variable backgrounds.

<sup>&</sup>lt;sup>1</sup>This is a reasonable assumption, as even without imposing integrability on *u* itself, the continuity equation and the neutrality condition imply that  $|u_0(x) - u_0(-x)| \to 0$  as  $x \to \infty$ .

**1.3.3.** The case of repulsive forcing. We now turn our attention to the study of critical thresholds for the repulsive EP systems. Unlike the attractive interaction case, there is no literature on the sub-critical criteria for the repulsive EP system in the variable backgrounds. We provide the first sub-critical results in this case. In order to present our results more precisely, we must introduce the method of phase plane analysis. To this end, we consider the auxiliary variables (*s*, *w*) (introduced by Liu, Engelberg, and the last author [13] in their study of critical thresholds of EP systems with constant backgrounds),

$$s := \frac{1}{\rho}$$
 and  $w := \frac{\partial_x u}{\rho}$ 

Let  $\Box' := \dot{\Box}(t, x(t))$  denote differentiation along particle path  $x(t) = x(t; \alpha)$ ,

$$\dot{x}(t;\alpha) = u(t, x(t;\alpha)), \quad x(0;\alpha) = \alpha.$$
(1.9)

We find that (abusing notation) w(t) = w(t, x(t)) and s(t) = s(t, x(t)) satisfy

$$\begin{cases} w' = -vw + k(1 - cs), \\ s' = w. \end{cases}$$
(1.10)

We observe that the question of global-in-time regularity vs. finite-time breakdown is reduced to whether s(t) attains zero  $-\rho(t, \cdot)$  blows up - in a finite time. There are two classes of initial configurations  $(w_0, s_0)$  of our interest. We say  $\Sigma^b$  is a sub-critical region of (1.10) if it is an *invariant* set, i.e.,  $(w_0, s_0) \in \Sigma^b$  implies  $(w(t), s(t)) \in \Sigma^b$  for all t > 0. On the other hand, we say  $\Sigma^{\sharp}$  is a super-critical region of (1.10) if for each  $(w_0, s_0) \in \Sigma^{\sharp}$ , there exists  $T_* > 0$  such that  $s(T_*) = 0$ . Hence, the set of sub-/super-critical regions gives rise to the criteria of initial data  $(\rho_0, u_0)$  for which the local-in-time classical solution of (1.1)–(1.2) becomes global or forms a singularity in a finite time. Lastly, we say the critical threshold is *sharp* if

$$\mathbb{R} imes \mathbb{R}_+ = \Sigma^{\flat} \cup \Sigma^{\sharp}.$$

For the constant backgrounds, where the solutions to (1.10) can be obtained explicitly, the sharp critical threshold was obtained in [13] for the undamped case, precisely given as

$$\Sigma^{\flat} = \left\{ -\sqrt{s(2-\bar{c}s)} < w < \sqrt{s(2-\bar{c}s)} \right\}, \quad \Sigma^{\sharp} = \mathbb{R} \times \mathbb{R}_{+} \setminus \Sigma^{\flat}, \tag{1.11}$$

where  $c = \bar{c} > 0$  denotes the background. Recently, Bhatnagar and Liu obtained the sharp critical thresholds for the damped case [5].

For the variable backgrounds, however, we note that  $c = c(t, x(t, \alpha))$  in  $(1.10)_2$  varies along the characteristic and the system (1.10) is not closed. We overcome this non-locality by exploiting the uniform bounds of c. We compare the system (1.10) to the one associated with its extremes  $c = c_{\pm}$ , and we establish the critical thresholds for the variable background by the solutions originating from the constant backgrounds. This idea – a Lyapunov-based approach – is widely used for attractive and repulsive force

in Sections 3 and 4. By nature of this comparison method, all our results in the sequel are sharp in the sense that we can recover the sharp critical threshold results for the constant backgrounds once we set  $c_{-} = c_{+}$ .

Throughout this paper, we use

$$\mathcal{L}(w,s) = w + f(s)$$

as a general form of the Lyapunov function, where f will be chosen so as to make the *sign* of  $\mathcal{L}$  determine the critical threshold regions. In particular,  $\mathcal{L}(w, s) = 0$  designates the boundary of sub-/super-critical regions – often, we will call this the threshold line.

To obtain the threshold lines of the (damped) repulsive EP system, we make use of the following Lyapunov functions:

$$\mathscr{L}_P(w,s) := w + \sqrt{2P(s)}, \quad \mathscr{L}_N(w,s) := w - \sqrt{2N(s)}, \tag{1.12}$$

where P and N are the maximal solutions of the following ODEs:

$$\frac{\mathrm{d}P}{\mathrm{d}s} = \nu \sqrt{2P(s)} + 1 - c_1 s, \qquad P(0) = 0, \tag{1.13}$$

$$\frac{\mathrm{d}N}{\mathrm{d}s} = -\nu\sqrt{2N(s)} + 1 - c_2 s, \quad N(s_*) = 0, \tag{1.14}$$

with constants  $c_1, c_2 > 0, \nu \ge 0$ , and  $s_* > \frac{1}{c_2}$ . The choice of  $c_1$  and  $c_2$  will depend on the context, either  $\bar{c}$  for the constant background or  $c_{\pm}$  for the variable background case.

To exploit these Lyapunov functions, we characterize the domains of  $\sqrt{2P(\cdot)}$  and  $\sqrt{2N(\cdot)}$ . By abusing notation, we denote the domain of  $\sqrt{2P(\cdot)}$  by Dom(P). We apply the parallel statement to N. Indeed, the domains are characterized as

$$Dom(P) = \begin{cases} [0, \infty), & \nu \ge 2\sqrt{c_1}, \\ [0, \tilde{s}], & 0 \le \nu < 2\sqrt{c_1}, \end{cases}$$

$$Dom(N) = \begin{cases} (-\infty, s_*], & \nu \ge 2\sqrt{c_2}, \\ [s_{**}, s_*], & 0 \le \nu < 2\sqrt{c_2}, \end{cases}$$
(1.15)

where

$$\tilde{s} = \frac{1 + e^{\gamma_1}}{c_1}, \quad s_{**} = \frac{1}{c_2} - \left(s_* - \frac{1}{c_2}\right)e^{\gamma_2}, \quad \gamma_i := \frac{\pi\nu}{\sqrt{4c_i - \nu^2}}, \quad i \in \{1, 2\}.$$
(1.16)

In the context of constructing the sub-/super-critical regions, we always choose  $s_* := \tilde{s}$  so as to make these regions as large as possible. Here we mention that the formula for  $s_{**}$  plays a vital role in establishing the critical regions.

The main novelty of this approach originated with Bhatnagar and Liu [5], with the purpose of replacing an explicit *algebraic description* of the critical threshold, by a *dif-ferential description* expressed in terms of a maximal solution of the ODE in the w-s plane:

$$\frac{dQ(s)}{ds} = v + \frac{1 - \bar{c}s}{Q(s)}.$$
(1.17)

Indeed, while  $\sqrt{2P}$  and  $-\sqrt{2N}$  coincide with the maximal solution Q of [5] in the case of  $c_1 = c_2 = \bar{c}$ , here we make a distinction between  $\sqrt{2P}$  and  $-\sqrt{2N}$ , as we trace the *sign* of Q, corresponding to the positive and negative solutions. In Section 4 we provide a precise characterization of domains (1.15)–(1.16), replacing the graphical representation associated with (1.17) in [5].

**Comparison principle.** We now introduce our comparison principles, deploying the Lyapunov functions (1.12), to establish the critical thresholds. For this, we denote  $P_{\pm}$  by the solution of (1.13) with  $c_1 = c_{\pm}$  and  $N_{\pm}$  by that of (1.14) with  $c_2 = c_{\pm}$ . The *weak* comparison principle is given as follows:

$$\begin{aligned} \mathcal{L}_{P_{-}}(w(t), s(t)) &\leq 0 \quad \text{if } \mathcal{L}_{P_{-}}(w_{0}, s_{0}) \leq 0, \\ \mathcal{L}_{N_{+}}(w(t), s(t)) &\geq 0 \quad \text{if } \mathcal{L}_{N_{+}}(w_{0}, s_{0}) \geq 0, \end{aligned}$$
(1.18)

provided that

$$\forall \tau \in [0, t], s(\tau) \in \text{Dom}(P_{-}) \text{ (resp. } s(\tau) \ge 0 \text{ and } s(\tau) \in \text{Dom}(N_{+}))$$

This gives rise to the construction of super-critical regions:

$$\Sigma^{\sharp} := \mathbb{R} \times \mathbb{R}_{+} \setminus \left\{ -\sqrt{2P_{-}(s)} < w \right\} \qquad \qquad \text{for } \nu \ge 2\sqrt{c_{-}},$$

and

$$\Sigma^{\sharp} := \mathbb{R} \times \mathbb{R}_+ \setminus \left\{ -\sqrt{2P_{-}(s)} < w < \sqrt{2N_{+}(s)} \right\} \quad \text{for } 0 \le \nu < 2\sqrt{c_{-}}. \tag{1.19}$$

On the other hand, we have the *strong* comparison principle:

$$\begin{aligned} \mathcal{L}_{P_{+}}(w(t), s(t)) &> 0 \quad \text{if } \mathcal{L}_{P_{+}}(w_{0}, s_{0}) > 0, \\ \mathcal{L}_{N_{-}}(w(t), s(t)) &< 0 \quad \text{if } \mathcal{L}_{N_{-}}(w_{0}, s_{0}) < 0, \end{aligned}$$
(1.20)

as long as

$$\forall \tau \in [0, t], s(\tau) \in \text{Dom}(P_+) \text{ (resp. } s(\tau) \ge 0 \text{ and } s(\tau) \in \text{Dom}(N_-)).$$

It leads us to define

$$\Sigma^{\flat} := \left\{ -\sqrt{2P_+(s)} < w \right\} \qquad \qquad \text{for } \nu \ge 2\sqrt{c_+},$$

and

$$\Sigma^{\flat} := \left\{ -\sqrt{2P_{+}(s)} < w < \sqrt{2N_{-}(s)} \right\} \qquad \text{for } 0 \le \nu < 2\sqrt{c_{+}}.$$
(1.21)

It is not always the case that the set  $\Sigma^{\flat}$  (1.21) is sub-critical; some extra conditions should be met to secure the sub-criticality.

**Closing condition.** In order to prove that  $\Sigma^{\flat}$  (1.21) and  $\Sigma^{\sharp}$  (1.19) are *indeed* sub-/supercritical regions, we require

$$s_{**} \leqslant 0 \tag{1.22}$$

to hold. We call (1.22) the *closing condition*. The term *closing* is motivated by the geometric depiction that the  $\Sigma^{\flat}$  and  $\mathbb{R} \times \mathbb{R}_+ \setminus \Sigma^{\sharp}$  are enclosed by its boundaries, as in Figures 1 and 3.

The closing condition plays a key role in constructing a sub-/super-critical region. We illustrate this by demonstrating that  $\Sigma^{\sharp}$  in (1.19) is a super-critical set. We readily check that the closing condition is validated for  $\Sigma^{\sharp}$ . Indeed, by assigning  $(c_1, c_2) = (c_-, c_+)$  to (1.16), we obtain

$$s_* \ge \frac{2}{c_-} > \frac{1}{c_+}$$
 and  $s_{**} \le \frac{1}{c_+} - \left(\frac{2}{c_-} - \frac{1}{c_+}\right) \le 0.$ 

In particular, it implies that the  $\mathcal{L}_{N_+}(w, s) = 0$  line crosses the *w*-axis at  $\sqrt{2N_+(0)} \ge 0$  so that the boundary of  $\Sigma^{\sharp}$  can be decomposed as

$$\{\mathcal{L}_{P_{-}}(w,s)=0\} \cup \{\mathcal{L}_{N_{+}}(w,s)=0, s \ge 0\} \cup \{w \ge \sqrt{2N_{+}(0)}, s=0\} \cup \{w \le 0, s=0\}$$

By the weak comparison principle (1.18), we deduce that if  $(w_0, s_0) \in \Sigma^{\sharp}$ , then (w(t), s(t)) can escape  $\Sigma^{\sharp}$  only through the s = 0 line (with  $w \leq 0$  because  $s(t)' = w(t) \geq \sqrt{2N_+(0)} > 0$  on the right-hand part of horizontal boundary). In fact, for each  $(w_0, s_0) \in \Sigma^{\sharp}$ , there exists  $t_* < +\infty$  such that  $s(t_*) = 0$ . Consult Propositions 4.4 and 4.5. It follows that  $\Sigma^{\sharp}$  is a super-critical set. Hence, we obtain a super-critical condition for the undamped repulsive EP with variable backgrounds.

**Theorem 1.7** (Finite-time breakdown). Consider the repulsive EP system (1.1)–(1.2). For given initial data ( $\rho_0$ ,  $u_0$ ), if there exists  $x \in \mathbb{R}$ , which does not satisfy

• 
$$-\rho_0(x)\sqrt{2P_-(\frac{1}{\rho_0(x)})} < u'_0(x) \text{ when } v \ge 2\sqrt{c_-},$$
  
•  $-\rho_0(x)\sqrt{2P_-(\frac{1}{\rho_0(x)})} < u'_0(x) < \rho_0(x)\sqrt{2N_+(\frac{1}{\rho_0(x)})} \text{ when } 0 \le v < 2\sqrt{c_-},$ 

• 
$$-\rho_0(x)\sqrt{2P_-(\frac{1}{\rho_0(x)})} < u'_0(x) < \rho_0(x)\sqrt{2N_+(\frac{1}{\rho_0(x)})}$$
 when  $0 \le v < 2\sqrt{2N_+(\frac{1}{\rho_0(x)})}$ 

then the solution  $(\rho, u)$  will lose  $\mathcal{C}^1$  regularity in a finite time.

**Remark 1.8.** For the undamped  $\nu = 0$  case, we have

$$\Sigma^{\sharp} = \mathbb{R} \times \mathbb{R}_{+} \setminus \left\{ -\sqrt{s(2-c_{-}s)} < w < \sqrt{s(2-c_{+}s) + \frac{4}{c_{-}} \left(\frac{c_{+}}{c_{-}} - 1\right)} \right\}.$$
 (1.23)

Note that  $\Sigma^{\sharp}$  coincides with (1.11) if we set  $c_{-} = c_{+} = \bar{c}$ . Moreover, it improves previously known super-critical results for repulsive EP systems with variable backgrounds [1, 22]. In the phase plane coordinates, these previous super-criteria can be expressed as  $s \ge \frac{2}{c_{-}}$  and  $w + \sqrt{2s} \le 0$ , respectively. In fact, (1.23) contains these two regions. We display this in Figure 1 for the reader's convenience.



Figure 1. Illustration of the super-critical regions

We now turn our attention to the sub-critical case. The closing condition can also be exploited to show that  $\Sigma^{\flat}$  in (1.21) is a sub-critical set. Once we assume the closing condition holds for  $\Sigma^{\flat}$ , then the boundary of  $\Sigma^{\flat}$  can be decomposed as

$$\{\mathcal{L}_{P_+}(w,s) = 0\} \cup \{\mathcal{L}_{N_-}(w,s) = 0, \ s > 0\} \cup \{0 < w \le \sqrt{2N_-(0)}, \ s = 0\}.$$
(1.24)

Note that  $\Sigma^{b}$  is an open set, and we can prove that any trajectory issued inside this set cannot hit the boundary by the strong comparison principle. Consult Proposition 4.7 for the details.

Unlike the super-critical set, the closing condition is not satisfied for  $\Sigma^{b}$  in (1.21) in general. For instance, if  $\nu = 0$  with  $c_{-} < c_{+}$ , then the closing condition is violated because

$$s_{**} = \frac{2}{c_-} - \frac{2}{c_+} > 0.$$

In fact, the closing condition is not merely an artifact of technique but reflects the difficulties in obtaining sub-critical regions for the repulsive EP system. To exhibit this, we introduce the following system of ODEs:

$$\begin{cases} w' = 1 - (1 + \varepsilon \sin(t))s, \\ s' = w, \end{cases}$$
(1.25)

with  $(w_0, s_0) = (0, 1)$ . Although the deviation of *background* from 1 is assumed to be small as  $|(1 + \varepsilon \sin(t)) - 1| \le \varepsilon$ , the oscillation of (w(t), s(t)) and c(t) can be aligned so that it makes a *resonance*. In particular, s(t) attains zero in a finite time. We implemented the numerical experiment of (1.25) with  $\varepsilon = 0.05$  using the built-in ode45 MATLAB command. See Figure 2 for its depiction. This example convinces us that the closing condition is necessary to construct a sub-critical region.



Figure 2. The evolution of s (left) and the phase plane plot (right) of the solution trajectory of (1.25)

In the presence of damping ( $\nu > 0$ ), the closing condition always holds for the  $\nu \ge 2\sqrt{c_+}$  case. For the  $0 < \nu < 2\sqrt{c_+}$  case, the closing condition can be entertained by imposing suitable conditions on  $c_+$ ,  $c_-$ , and  $\nu$ . The heuristic is that the ratio of  $c_+$  over  $c_-$  is close enough to 1, and the damping effect is large enough to control the oscillation so that the system is similar to that of a constant background case.

**Theorem 1.9** (Global-in-time regularity). Consider the repulsive EP system (1.1)–(1.2). It admits a global classical solution, depending on the relative size of v and  $\sqrt{c_{\pm}}$ . Specifically, a global classical solution exists if the initial data  $(\rho_0, u_0)$  falls into one of the following two cases, so that for all  $x \in \mathbb{R}$  there holds

#1: 
$$v \ge 2\sqrt{c_+} \text{ and } -\rho_0(x)\sqrt{2P_+(\frac{1}{\rho_0(x)})} < u_0'(x);$$

$$#2: \ 0 < \nu < 2\sqrt{c_{+}} \ and -\rho_{0}(x)\sqrt{2P_{+}\left(\frac{1}{\rho_{0}(x)}\right)} < u_{0}'(x) < \rho_{0}(x)\sqrt{2N_{-}\left(\frac{1}{\rho_{0}(x)}\right)},$$
where  $s_{+} := \frac{1+e^{\gamma_{+}}}{c_{+}}, \ \gamma_{\pm} := \frac{\pi\nu}{\sqrt{4c_{\pm}-\nu^{2}}}, \ and \ one \ of \ the \ following \ two \ sub-cases \ is \ fulfilled:$ 

$$\begin{cases} #2.1: \ either \ 2\sqrt{c_{-}} \le \nu < 2\sqrt{c_{+}} \ and \ s_{+}c_{-} > 1, \\ #2.2: \ or \ 0 < \nu < 2\sqrt{c_{-}} \ and \ e^{\gamma_{-}}(s_{+}c_{-}-1) \ge 1. \end{cases}$$

$$(1.26)$$

**Remark 1.10.** In fact, the condition (1.26) is equivalent to the closing condition (1.22). We arrive at (1.26) by putting  $(c_1, c_2) = (c_+, c_-)$  in (1.16). If we assume  $c_- = c_+$ , the closing condition is satisfied without any extra condition, and we recover the sharp critical threshold result for the damped EP system in [5].

We display the critical threshold results for the repulsive interaction case in Figure 3.



Figure 3. Illustration of the sub-/super-critical regions for the repulsive EP system

**1.3.4.** Application to the EP system for cold-plasma-ion dynamics. Our final result is finite-time singularity formation and the global regularity of the damped EP system for cold-plasma-ion dynamics. Consider a plasma consisting of electrons and ions on  $\mathbb{R}$ , where both electrons and ions have constant temperature. We assume that the temperatures of electrons and ions equal 1 and 0, respectively. In this case, our main system is given by

$$\begin{cases} \partial_t \rho + \partial_x (\rho u) = 0, \\ \partial_t u + u \partial_x u = -\nu u - \partial_x \phi, \\ -\partial_{xx} \phi = \rho - e^{\phi}, \end{cases}$$
(1.27)

where the density of electrons  $\rho_e$  is given by the *Maxwell–Boltzmann relation* [11], i.e.,  $\rho_e = e^{\phi}$ , where  $\phi = \phi(t, x)$  is the electric potential.

Note that the system (1.27) corresponds to (1.1) with  $(k, c) = (1, e^{\phi})$ . We consider the system (1.27) around a constant state (1, 0), i.e.,

$$(\rho, u)(t, x) \to (1, 0) \quad \text{as } |x| \to \infty.$$
 (1.28)

We notice from the above that  $\phi(t, x) \to 0$  as  $|x| \to \infty$ . For the system (1.27)–(1.28) with  $\nu = 0$ , the local-in-time existence and uniqueness of smooth solutions are studied in [18] for multi-dimensional cases. Additionally, global smooth irrotational solutions with small amplitude are constructed in [15] in three dimensions.

In order to apply the framework developed in Section 1.3.3, we need to secure lower and upper bounds of  $e^{\phi}$ . To this end, we introduce the total energy function H = H(t) for (1.1)–(1.2) as

$$H = H(t) := \int_{\mathbb{R}} \left( \frac{1}{2} \rho u^2 + \frac{1}{2} |\partial_x \phi|^2 + U(\phi) \right) \mathrm{d}x, \tag{1.29}$$

where  $U(r) := (r-1)e^r + 1$ , which is non-negative for all  $r \in \mathbb{R}$ .

Then, as long as there exist classical solutions, one can readily check that the total energy is not increasing in time, i.e.,  $H(t) \leq H_0 := H(0)$  for all  $t \geq 0$ , and conserved in time when  $\nu = 0$ . It was investigated in [1] that a uniform bound of the potential function,  $\phi$ , can be obtained by  $H_0$ . Hence, we can apply the results in the previous section to this case.

**Theorem 1.11** (Finite-time breakdown). Consider the repulsive EP system (1.27)–(1.28) with v = 0. For a given initial data  $(\rho_0, u_0)$ , if there exists  $x \in \mathbb{R}$  which does not satisfy

$$-\sqrt{2\rho_0(x)-c_-} < u_0'(x) < \sqrt{2\rho_0(x)-c_+ + \frac{4}{c_-} \left(\frac{c_+}{c_-} - 1\right)\rho_0(x)^2},$$

then the classical solution  $(\rho, u)$  will lose  $\mathcal{C}^1$  regularity in a finite time. Here,  $c_- := \exp(V_-^{-1}(H_0))$  and  $c_+ := \exp(V_+^{-1}(H_0))$  denote positive constants, where  $V_{\pm}^{-1}$  are inverses of  $V_{\pm}$  defined in (4.7).

For the damped Euler–Poisson system for a cold plasma, the global existence and uniqueness of  $H^s$  solutions are obtained in [21] under the smallness assumptions on the initial data ( $\rho_0, u_0$ ) near the constant equilibrium state (1, 0) in  $H^s$  norm. However, the following theorem shows that the global-in-time existence of classical solutions to the damped Euler–Poisson system for a cold plasma can be obtained *only* under the smallness assumption on the initial energy  $H_0$  compared to the strength of damping.

**Theorem 1.12** (Global-in-time regularity). For a class of initial data ( $\rho_0$ ,  $u_0$ ), whose initial energy  $H_0$  (1.29) is small enough, we assume that

$$\nu \geqslant C\sqrt{H_0},\tag{1.30}$$

for some absolute constant C > 0 and

$$\|\rho_0 - 1\|_{L^{\infty}} + \|u_0'\|_{L^{\infty}} \le \Lambda_0, \tag{1.31}$$

where  $\Lambda_0 > 0$  is a small absolute constant. Then the damped cold-ion system (1.27)–(1.28) admits a global classical solution with initial data ( $\rho_0, u_0$ ).

**Remark 1.13.** We note that (1.31) restricts  $(\rho_0, u_0)$  to fall into the sub-critical regions in Theorem 1.9. We can take  $\Lambda_0 = \frac{1}{2}$ , for instance. Thus, we are only concerned with the smallness of  $H_0$  and the requirement on the strength of damping (1.30).

### 1.4. Outline of the paper

This paper is organized as follows. In Section 2 we establish the local-in-time wellposedness of the EP system ensuring the neutrality condition. We then prove the illposedness result of Theorem 1.4. We also explore the propagation of integrability for the density perturbation around the background, which serves as an independent area of interest in this study. In Section 3 we present our Lyapunov-based approach to establish the critical thresholds in the attractive case. In Section 4 we discuss the repulsive interaction case, completing the proofs of Theorems 1.7 and 1.9. As a consequence, we prove Theorem 1.12, which concerns the global regularity of damped cold-plasma-ion dynamics.

### 2. Local well-posedness and the neutrality condition

In this section, we present a rigorous mathematical treatment of the local well-posedness and the neutrality condition of the EP system introduced in Section 1.3.1. We begin by proving the local well-posedness theory.

### 2.1. Local well-posedness and non-existence

*Proof of Theorem* 1.1. We only provide a priori estimates of solutions in the desired regularity spaces. The local well-posedness can be obtained by the standard arguments developed for the types of conservation laws; see [4,5,25] for instances. Indeed, we obtain

$$\frac{d}{dt} (\|\rho - \bar{c}\|_{H^s \cap \dot{W}^{-1,p}} + \|u\|_{L^p} + \|\partial_x u\|_{H^s}) 
\leq C(\|\partial_x u\|_{L^{\infty}} + \|\rho - \bar{c}\|_{L^{\infty}} + \bar{c} + 1) 
\times (\|\rho - \bar{c}\|_{H^s \cap \dot{W}^{-1,p}} + \|u\|_{L^p} + \|\partial_x u\|_{H^s}) + \|\eta\|_{H^s \cap \dot{W}^{-1,p}},$$
(2.1)

where C > 0 is an absolute constant; see Appendix B for the details of (2.1). We apply Grönwall's lemma and  $\|\rho - c\|_{H^s \cap \dot{W}^{-1,p}} \leq \|\rho - \bar{c}\|_{H^s \cap \dot{W}^{-1,p}} + \|\eta\|_{H^s \cap \dot{W}^{-1,p}}$  to obtain (1.5), thereby concluding the proof.

In fact, the proofs of Theorem 1.4 and Proposition 2.1 below are the main themes in this section. They restrict our attention to the case of a constant background state. For the sake of simplicity in presenting our results, we only focus on the undamped case. The damped case can be treated in a similar way. Our analysis relies on the characteristic method and, especially, the explicit formulation of the Jacobian of the characteristic flow. We briefly explain the main tools and the strategy.

For a  $\mathcal{C}^1$  solution  $(\rho, u)$  to EP system (1.1)–(1.2) with  $c = \bar{c}$ , we consider the characteristic flow  $x(t, \alpha)$  defined in (1.9). Write

$$\Gamma(t,\alpha) := \partial_{\alpha} x(t,\alpha),$$

signifying the Jacobian of the map  $x(t, \cdot): \mathbb{R} \to \mathbb{R}$ . We note that  $\Gamma(0, \alpha) = 1$  and  $x(t, \cdot)$  is a diffeomorphism as long as  $\Gamma(t, \cdot) > 0$ . In particular, we have

$$\lim_{\alpha \to \pm \infty} x(t, \alpha) = \pm \infty.$$
 (2.2)

Taking  $\partial_{\alpha}$  of (1.9), we obtain

$$\partial_x u(t, x(t, \alpha)) = \frac{\partial_t \Gamma(t, \alpha)}{\Gamma(t, \alpha)}.$$
 (2.3)

In view of the mass conservation  $\rho' = -\rho \partial_x u$ , we find that

$$\rho(t, x(t, \alpha)) = \frac{\rho_0(\alpha)}{\Gamma(t, \alpha)}.$$
(2.4)

Finally, we recall the explicit formulas of  $\Gamma(t, \alpha)$  for the undamped ( $\nu = 0$ ) case [13]. They will lay the foundation of the proofs throughout this section:

• Repulsive case:

$$\Gamma(t,\alpha) = 1 + \left(\frac{\rho_0(\alpha)}{\bar{c}} - 1\right) (1 - \cos(\sqrt{\bar{c}}t)) + u_0'(\alpha) \frac{\sin(\sqrt{\bar{c}}t)}{\sqrt{\bar{c}}}.$$
 (2.5)

• Attractive case:

$$\Gamma(t,\alpha) = 1 + \left(\frac{\rho_0(\alpha)}{\bar{c}} - 1\right)(1 - \cosh(\sqrt{\bar{c}}t)) + u'_0(\alpha)\frac{\sinh(\sqrt{\bar{c}}t)}{\sqrt{\bar{c}}}.$$
 (2.6)

We now prove Theorem 1.4.

*Proof of Theorem* 1.4. For simplicity, we assume  $\bar{c} = 1$  and  $\nu = 0$ . We set initial data  $(\rho_0, u_0)$  by assigning  $u_0 = 0$  and

$$\rho_0(x) = 1 + F'(x), \quad F(x) := \frac{1}{(1 + x^2)^{1/4}}$$

We note that

$$F \notin L^2(\mathbb{R}), \quad F \in \bigcap_{p>2} L^p(\mathbb{R}), \quad \rho_0 - 1 = F' \in \bigcap_{k=0}^{\infty} H^k(\mathbb{R}),$$

where  $H^0 := L^2$  by convention. From the definition of  $\rho_0$ , we find

$$\int_{\mathbb{R}} \rho_0 - 1 \, \mathrm{d}x = \lim_{R \to \infty} (F(R) - F(0)) - \lim_{L \to \infty} (F(0) - F(-L)) = 0.$$

Thus the neutrality condition is initially satisfied.

Since  $(\rho_0 - 1, u_0) \in X^{p,s}$  for any p > 2, by Theorem 1.1, there exists a unique localin-time classical solution  $(\rho, u)$  to EP system (1.1)–(1.2) satisfying

$$(\rho - 1, u) \in L^{\infty}([0, T]; X^{p,s})$$

for some T > 0. We note that this, in particular, requires u to decay at the far field,

$$\lim_{\alpha \to \pm \infty} u(t, \alpha) = 0, \quad \forall t \in [0, T]$$
(2.7)

and be bounded in  $[0, T] \times \mathbb{R}$  due to (A.1):

$$\|u(t)\|_{L^{\infty}(\mathbb{R})} \leq \|u(t)\|_{L^{p}(\mathbb{R})} + \|\partial_{x}u(t)\|_{L^{2}(\mathbb{R})} \leq C, \quad \forall t \in [0, T].$$
(2.8)

Similarly, we observe that  $\|\partial_x u\|_{L^{\infty}}$  is uniformly bounded in  $t \in [0, T]$ . Hence, we can employ the forward characteristic  $x(t, \cdot)$  defined in (1.9). For each  $t \in [0, \delta]$  and R, L > 0, we infer from (2.3) that

$$u(t, x(t, R)) - u(t, x(t, -L)) = \int_{x(t, -L)}^{x(t, R)} (\partial_x u)(t, x) dx$$
$$= \int_{-L}^{R} (\partial_x u)(t, x(t, \alpha)) dx(t, \alpha)$$
$$= \int_{-L}^{R} \partial_t \Gamma(t, \alpha) d\alpha.$$

Hence, an application of the explicit formula (2.5), for the case of repulsive forcing, supplies that

$$u(t, x(t, R)) - u(t, x(t, -L)) = (F(R) - F(-L)) \cdot \sin t.$$

By taking  $L \to \infty$ , thanks to (2.2) and (2.7), we find that, for all small t > 0,

$$u(t, x(t, R)) = F(R) \sin t.$$

Furthermore, we observe that

$$|x(t,R) - R| \le CT, \quad \forall t \in [0,T]$$

due to (2.8). This implies that for all sufficiently large R > 0,

$$u(t, R) \sim F(R) \sin t = \frac{\sin t}{(1+R^2)^{1/4}} \notin L^2(\mathbb{R}), \quad \forall t \in (0, \delta],$$

for  $\delta := \frac{1}{2} \min\{T, \pi\} > 0$ . The attractive forcing case can be tackled in the same manner by recalling (2.6). The conclusion of the theorem is now immediate.

### 2.2. Integrability and its failure in the neutralized EP system

The neutrality condition (1.3) is formulated as an *improper integral*, meaning that it does not require the integrability of  $\rho - c$ . As an independent area of interest, we investigate whether *integrability* of  $\rho - c$  can be propagated over time. Specifically, we ask whether

$$\rho(t, \cdot) - c(t, \cdot) \in L^1(\mathbb{R}) \tag{2.9}$$

propagates over time as well.

In the following, we construct an *anomalous solution*, exhibiting the non-propagation of the integrability in time even when starting with the neutralized and integrable initial data. For simplicity, we treat the p = 2 case, noting that parallel results hold for 1 .

**Proposition 2.1** (Failure of integrability: anomalous solution). *Consider the EP system* (1.1) with  $c = \bar{c} > 0$ . There exists an initial data  $(\rho_0, u_0)$  satisfying

$$(\rho_0 - \overline{c}, u_0) \in H^s \cap \dot{H}^{-1} \cap L^1(\mathbb{R}) \times H^{s+1}(\mathbb{R})$$

such that the local solution  $(\rho, u)$  to (1.1)–(1.2) satisfies

$$\rho(t,\cdot) - \bar{c} \notin L^1(\mathbb{R})$$

as long as t > 0.

The following lemma identifies the necessary and sufficient conditions for the integrability (2.9) in the case of constant background. It plays a central role in constructing the *anomalous* solutions.

**Lemma 2.2** (Neutrality condition: constant background). *Consider the EP system* (1.1) with  $c = \overline{c} > 0$ . Let  $(\rho, u)$  be a local-in-time classical solution of (1.1)–(1.2) in  $[0, T) \times \mathbb{R}$  with neutralized and integrable initial data, i.e.,

$$\rho_0 - \bar{c} \in L^1(\mathbb{R}), \quad \text{with } \int_{\mathbb{R}} (\rho_0(x) - \bar{c}) \, \mathrm{d}x = 0.$$
(2.10)

*Then the propagation of integrability* (2.9) *holds if and only if* 

$$u_0 \in \mathrm{BV}(\mathbb{R}), \quad \text{with } \int_{\mathbb{R}} u'_0(x) \,\mathrm{d}x = 0.$$

*Proof.* Let  $x(t, \alpha)$  be the forward characteristic flow satisfying (1.9). Note that  $\Gamma(t, \cdot) > 0$  is well defined for each  $t \in [0, T)$ . We claim that

$$\rho(t, \cdot) - \overline{c} \in L^1(\mathbb{R})$$
 if and only if  $\Gamma(t, \cdot) - 1 \in L^1(\mathbb{R})$ .

In this context, we apply (2.4) to see that

$$\int_{x(t,-R)}^{x(t,R)} |\rho(t,x) - \bar{c}| \, \mathrm{d}x = \int_{-R}^{R} |\rho(t,x(t,\alpha)) - \bar{c}| \, \mathrm{d}x(t,\alpha)$$
$$= \int_{-R}^{R} \left| \frac{\rho_0(\alpha)}{\Gamma(t,\alpha)} - \bar{c} \right| \Gamma(t,\alpha) \, \mathrm{d}\alpha$$
$$= \int_{-R}^{R} |(\rho_0(\alpha) - \bar{c}) - \bar{c}(\Gamma(t,\alpha) - 1)| \, \mathrm{d}\alpha.$$

On applying the triangle inequality with (2.2), the claim follows. Hence, we deduce from the explicit formulas for  $\Gamma(t, \cdot) - 1$  recorded in (2.5) and (2.6) that  $u'_0 \in L^1(\mathbb{R})$  if and only if  $\Gamma(t, \cdot) - 1 \in L^1(\mathbb{R})$ , which is equivalent to  $\rho(t, \cdot) - \overline{c} \in L^1(\mathbb{R})$ .

We now assume that  $u'_0 \in L^1(\mathbb{R})$ . Similarly, in view of (2.4), one sees that

$$\int_{\mathbb{R}} (\rho(t,x) - \bar{c}) \, \mathrm{d}x = \int_{\mathbb{R}} (\rho_0(\alpha) - \bar{c}) - \bar{c}(\Gamma(t,\alpha) - 1) \, \mathrm{d}\alpha = -\bar{c} \int_{\mathbb{R}} (\Gamma(t,\alpha) - 1) \, \mathrm{d}\alpha.$$

By inserting (2.5), (2.6) into  $\Gamma(t, \cdot) - 1$  and integrating over  $\mathbb{R}$ , we discern that

$$\int_{\mathbb{R}} (\rho(t, x) - \bar{c}) \, \mathrm{d}x = 0 \quad \text{if and only if} \quad \int_{\mathbb{R}} u_0'(\alpha) \, \mathrm{d}\alpha = 0$$

This confirms the conclusion of the lemma.

It is worth noting that if  $u_0 \notin BV(\mathbb{R})$ , we infer that the local classical solution instantaneously fails the integrability (2.9) by taking T > 0 arbitrarily small. Hence, this lemma leads us to find *anomalous* solutions in Proposition 2.1.

*Proof of Proposition* 2.1. We fix  $\rho_0 = \bar{c} > 0$ . It satisfies (2.10), evidently. In view of Theorem 1.1 and Lemma 2.2, we aim to choose  $u_0$  satisfying

$$u_0 \in H^{s+1} \setminus BV(\mathbb{R}).$$

Consider

$$u_0(x) := \frac{\sin x}{(1+x^2)^{\frac{3}{8}}}$$

Since  $u_0 \in H^{s+1}(\mathbb{R})$ , we obtain a local smooth solution  $(\rho, u)$  via Theorem 1.1. In particular, we find by means of Morrey's embedding that this solution is of class  $\mathcal{C}^1$ . However, upon noting

$$u_0'(x) = \frac{\cos x}{(1+x^2)^{\frac{3}{8}}} - \frac{3}{4} \frac{x \sin x}{(1+x^2)^{\frac{11}{8}}} \notin L^1(\mathbb{R}).$$

we find that  $(\rho, u)$  fails the integrability (2.9) instantaneously by Lemma 2.2. This completes the proof.

To circumvent this anomaly, additional regularity must be imposed, for instance,  $u_0 \in BV(\mathbb{R})$  as detailed in the following proposition.

**Proposition 2.3** (Well-posedness with integral neutrality). Suppose that the initial data  $(\rho_0, u_0)$  satisfy

$$(\rho_0 - c_0, u_0) \in H^s \cap \dot{H}^{-1} \cap L^1(\mathbb{R}) \times H^{s+1} \cap BV(\mathbb{R})$$

where c = c(t, x) is given as

$$c = \overline{c} + \eta, \quad \eta \in \mathcal{C}^0([0,\infty); H^s \cap \dot{H}^{-1} \cap L^1(\mathbb{R}))$$

for some  $\bar{c} > 0$ . Then the system (1.1)–(1.2) admits a classical solution ( $\rho(t, \cdot), u(t, \cdot)$ ) such that

$$\sup_{0 \le t \le T} (\|(\rho - c)(t)\|_{L^1} + \|(\partial_x u)(t)\|_{L^1}) < \infty$$
(2.11)

for some T > 0. In particular, integrability condition (2.9) holds for all  $t \in [0, T]$ .

*Proof.* We estimate the evolution of  $\|(\rho - \bar{c})(t)\|_{L^1}$  and  $\|(\partial_x u)(t)\|_{L^1}$ . Straightforward computations give

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \|\rho - \bar{c}\|_{L^{1}} &\leq \|\partial_{x}((\rho - \bar{c})u)\|_{L^{1}} + \bar{c}\|\partial_{x}u\|_{L^{1}} \\ &\leq \|\rho - \bar{c}\|_{H^{1}} \|u\|_{H^{1}} + \bar{c}\|\partial_{x}u\|_{L^{1}} \\ &\leq \|\{\rho(t, \cdot) - \bar{c}, u(t, \cdot)\}\|_{X^{2,s}}^{2} + \bar{c}\|\partial_{x}u\|_{L^{1}} \end{aligned}$$

and

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \|\partial_x u\|_{L^1} &\leq \|\partial_x u\|_{L^2}^2 + \|u\|_{L^2} \|\partial_x^2 u\|_{L^2} + \|\rho - c\|_{L^1} \\ &\leq 2 \|\{\rho(t, \cdot) - \bar{c}, u(t, \cdot)\}\|_{X^{2,s}}^2 + \|\rho - \bar{c}\|_{L^1} + \|\eta\|_{L^1}. \end{aligned}$$

By adding these two inequalities, we obtain

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} (\|\rho - \bar{c}\|_{L^1} + \|\partial_x u\|_{L^1}) \\ &\leq (1 + \bar{c})(\|\rho - c\|_{L^1} + \|\partial_x u\|_{L^1}) + 3\|\{\rho(t, \cdot) - \bar{c}, u(t, \cdot)\}\|_{X^{2,s}}^2 + \|\eta\|_{L^1}. \end{aligned}$$

Applying Grönwall's lemma to the above concludes the desired result (2.11).

We, therefore, conclude from Propositions 2.1 and 2.3 that it is necessary to assume additional regularity  $u_0 \in BV(\mathbb{R})$  to secure the propagation of the integrability of  $\rho - c$ .

# 3. Attractive forcing and the restricted borderline threshold

In this section, we focus on the critical threshold phenomenon of the attractive EP system (1.1)–(1.2). To apply the method of phase plane analysis introduced in Section 1.3.3, we consider the following ODE system:

$$\begin{cases} w' = -vw - (1 - c(t)s), \\ s' = w, \end{cases}$$
(3.1)

subject to  $(w(0), s(0)) = (w_0, s_0)$ , where c(t) is a smooth function satisfying  $0 < c_- \le c(t) \le c_+$ .

### 3.1. Constant background case

To give a motivation for the Lyapunov function, we initiate the discussion for the constant background case  $-c(t) = \overline{c}$ . In this case, (3.1) can be rewritten in the matrix form as

$$\begin{pmatrix} w\\ s-\frac{1}{\bar{c}} \end{pmatrix}' = \begin{pmatrix} -\nu & \bar{c}\\ 1 & 0 \end{pmatrix} \begin{pmatrix} w\\ s-\frac{1}{\bar{c}} \end{pmatrix}.$$
(3.2)

The dynamics of solution trajectories are determined by the eigenvalues of their coefficient matrix, whose eigenvalues are given by

$$\lambda_s = \frac{-\nu - \sqrt{\nu^2 + 4\bar{c}}}{2} < 0 \quad \text{and} \quad \lambda_u = \frac{-\nu + \sqrt{\nu^2 + 4\bar{c}}}{2} > 0,$$

with corresponding eigenvectors  $X_s = \begin{pmatrix} \lambda_s \\ 1 \end{pmatrix}$  and  $X_u = \begin{pmatrix} \lambda_u \\ 1 \end{pmatrix}$ . In particular, the equilibrium point  $(0, 1/\bar{c})$  of (3.2) is unstable; the solution curves are attracted to  $(0, 1/\bar{c})$  along the  $X_s$  direction while repulsed from  $(0, 1/\bar{c})$  along the  $X_u$  direction.

This motivates us to define the following Lyapunov functions:

$$\mathcal{L}_s(w,s) := w - \lambda_s \left( s - \frac{1}{\bar{c}} \right)$$
 and  $\mathcal{L}_u(w,s) := w - \lambda_u \left( s - \frac{1}{\bar{c}} \right)$ .

In particular, the solution (w(t), s(t)) of (3.2) satisfies

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_s(w(t),s(t)) = \lambda_u \mathcal{L}_s(w(t),s(t)) \quad \text{and} \quad \frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_u(w(t),s(t)) = \lambda_s \mathcal{L}_u(w(t),s(t)),$$

obtaining

$$\mathcal{L}_s(w(t), s(t)) = \mathcal{L}_s(w_0, s_0)e^{\lambda_u t} \quad \text{and} \quad \mathcal{L}_u(w(t), s(t)) = \mathcal{L}_u(w_0, s_0)e^{\lambda_s t}.$$
 (3.3)

Note that  $\mathcal{L}_s(w_0, s_0) \ge 0$  implies  $\mathcal{L}_s(w(t), s(t)) \ge 0$  for all  $t \ge 0$ . This, combined with s' = w and that the *w*-intercept of the line  $\mathcal{L}_s(w, s) = 0$  is positive, yields that  $\{\mathcal{L}_s(w, s) \ge 0, s > 0\}$  is *invariant*. On the other hand, if  $\mathcal{L}_s(w_0, s_0) < 0$ , then (3.3) implies that

$$\mathcal{L}_s(w(t), s(t)) \to -\infty, \quad \mathcal{L}_u(w(t), s(t)) \to 0 \quad \text{as } t \to \infty.$$

By noticing that

$$\mathcal{L}_s(w,s) - \mathcal{L}_u(w,s) = (-\lambda_s + \lambda_u) \Big( s - \frac{1}{\bar{c}} \Big),$$

we conclude that s(t) attains zero in a finite time. This argument recovers the sharp critical threshold for the attractive EP with constant background [13, Theorem 3.2], [4, Theorem 2.5].

### 3.2. Variable background case

To study the critical threshold for the variable background case, we consider the following Lyapunov functions:

$$\mathscr{L}_{s}^{\pm}(w,s) := w - \lambda_{s}^{\pm}\left(s - \frac{1}{c_{\pm}}\right) \quad \text{and} \quad \mathscr{L}_{u}^{-}(w,s) := w - \lambda_{u}^{-}\left(s - \frac{1}{c_{-}}\right),$$

where

$$\lambda_s^{\pm} = \frac{-\nu - \sqrt{\nu^2 + 4c_{\pm}}}{2} < 0 \text{ and } \lambda_u^{\pm} = \frac{-\nu + \sqrt{\nu^2 + 4c_{\pm}}}{2} > 0.$$

We readily check that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_{s}^{-}(w(t),s(t)) \ge \lambda_{u}^{-}\mathcal{L}_{s}^{-}(w(t),s(t)), \quad s(t) \ge 0.$$

Similarly, this implies that  $\{\mathcal{L}_s^-(w,s) \ge 0, s > 0\}$  is a sub-critical region. On the other hand, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{X}_{s}^{+}(w(t),s(t)) \leq \lambda_{u}^{+}\mathcal{X}_{s}^{+}(w(t),s(t)),$$
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{X}_{u}^{-}(w(t),s(t)) \geq \lambda_{s}^{-}\mathcal{X}_{u}^{-}(w(t),s(t)), \quad s(t) \geq 0$$

By Grönwall's lemma, we observe that

$$(-\lambda_{s}^{+}+\lambda_{u}^{-})s(t) = \mathcal{L}_{s}^{+}(w(t),s(t)) - \mathcal{L}_{u}^{-}(w(t),s(t)) + \frac{1}{\lambda_{u}^{+}} - \frac{1}{\lambda_{s}^{-}}$$
$$\leq \mathcal{L}_{s}^{+}(w_{0},s_{0})e^{\lambda_{u}^{+}t} + \mathcal{O}(1)$$

as long as s(t) is non-negative; in particular, if  $\mathcal{L}_{s}^{+}(w_{0}, s_{0}) < 0$ , then  $s(\cdot)$  attains zero in a finite time. Hence, we obtain the sub-/super-critical thresholds for attractive EP systems with variable backgrounds, depicted in Figure 4. They coincide with [4, Theorems 2.2 and 2.3].

# 4. Repulsive forcing and threshold with variable background

In this section, we aim to establish the critical thresholds for the repulsive EP system (1.1)–(1.2). To this end, we conduct the phase plane analysis as introduced in Section 1.3.3; for repulsive interaction, we recall the transformed ODE system (1.10):

$$\begin{cases} w' = -vw + 1 - c(t)s, \\ s' = w, \end{cases}$$
(4.1)

subject to initial point  $(w_0, s_0)$ . Here we assume that  $0 < c_- \leq c(t) \leq c_+$ .



**Figure 4.** Illustration of the sub-critical  $\{\mathcal{L}_s^-(w, s) \ge 0\}$  and super-critical  $\{\mathcal{L}_s^+(w, s) < 0\}$  criteria for the attractive case

We first confirm the comparison principles. Note that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_{P}(w(t), s(t)) = -vw(t) + 1 - c(t)s(t) + \left(\frac{v\sqrt{2P(s(t))} + 1 - c_{1}s(t)}{\sqrt{2P(s(t))}}\right)w(t) \\
= \left(\frac{1 - c_{1}s(t)}{\sqrt{2P(s(t))}}\right)\mathcal{L}_{P}(w(t), s(t)) + (c_{1} - c(t))s(t).$$
(4.2)

Similarly, we observe that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_N(w(t), s(t)) = \left(\frac{c_2 s(t) - 1}{\sqrt{2N(s(t))}}\right)\mathcal{L}_N(w(t), s(t)) + (c_2 - c(t))s(t).$$
(4.3)

In particular, putting  $(c_1, c_2) = (c_-, c_+)$  yields

$$\begin{aligned} \mathcal{L}_{P_{-}}(w(t),s(t)) &\leq \mathcal{L}_{P_{-}}(w_{0},s_{0}) \exp\left(\int_{0}^{t} \frac{1-c_{-}s(\tau)}{\sqrt{2P_{-}(s(\tau))}} \,\mathrm{d}\tau\right), \\ \mathcal{L}_{N_{+}}(w(t),s(t)) &\geq \mathcal{L}_{N_{+}}(w_{0},s_{0}) \exp\left(\int_{0}^{t} \frac{c_{+}s(\tau)-1}{\sqrt{2N_{+}(s(\tau))}} \,\mathrm{d}\tau\right), \end{aligned}$$

provided that  $s(\tau)$  satisfies  $P_{-}(s(\tau)) \ge 0$  (resp.  $N_{+}(s(\tau)) \ge 0$ ) for all  $\tau \in [0, t]$ , and the weak comparison principle (1.18) follows. Similarly, by taking  $(c_1, c_2) = (c_+, c_-)$ , we obtain the strong comparison principle (1.20).

The Lyapunov-based approach explained in Section 1.3.3 – essentially for the *closing* condition (1.22) – makes use of the domains of  $\sqrt{2P}$  and  $\sqrt{2N}$ . By witnessing the

defining equations of *P* and *N* in (1.13), (1.14), we observe that the domains of  $\sqrt{2P(\cdot)}$ ,  $\sqrt{2N(\cdot)}$  correspond to the set of  $s \in \mathbb{R}$  for which  $P(s) \ge 0$ ,  $N(s) \ge 0$ , respectively. It implies that the endpoints of the domains are precisely determined by zeros of *P* and *N*. The domains (1.15) can be obtained by the explicit solutions for (4.1).

We first investigate the exact formula for  $\tilde{s}$  and  $s_{**}$  in (1.16). We take index  $i \in \{1, 2\}$ . For the  $0 \le v < 2\sqrt{c_i}$  case, the solution (w(t), s(t)) of (4.1) with  $c(t) = c_i > 0$ ,  $(w_0, s_0) = (0, a)$  is explicitly given as

$$w(t) = -\frac{1}{\mu_i}(\bar{c}a - 1)\sin(\mu_i t)e^{-\frac{\nu t}{2}}, \quad \mu_i = \sqrt{c_i - \frac{\nu^2}{4}},$$
  

$$s(t) = \frac{1}{\bar{c}} + \left(a - \frac{1}{\bar{c}}\right)\left(\cos\mu_i t + \frac{\nu}{2\mu_i}\sin\mu_i t\right)e^{-\frac{\nu t}{2}}.$$
(4.4)

To unveil the formula for  $\tilde{s}$  (1.16), we take  $a = \tilde{s}$  and index i = 1 so that

$$\mathcal{L}_P(w_0, s_0) = 0 + \sqrt{2P(\tilde{s})} = 0,$$

where P satisfies (1.13). By (4.2), we observe that

$$\mathcal{L}_{P}(w(t), s(t)) = w(t) + \sqrt{2P(s(t))} = 0, \quad t \in \left[0, \frac{\pi}{\mu_{1}}\right]$$

Note that the range of t can be obtained by the criteria  $P(s(t)) \ge 0$  and (4.4). In particular, we deduce that

$$s\left(\frac{\pi}{\mu_i}\right) = 0 \iff \tilde{s} = s_0 = \frac{1 + e^{\gamma_1}}{c_1}$$

Similarly, by taking  $a = s_*$  and i = 2, we obtain

$$\sqrt{2N(s(t))} = w(t) \ge 0 \quad \text{for } t \in \left[-\frac{\pi}{\mu_2}, 0\right].$$

By noting  $s_0 = s_*$ , we infer that

$$s_{**} = s\left(-\frac{\pi}{\mu_2}\right) = \frac{1}{c_2} - \left(s_* - \frac{1}{c_2}\right)e^{\gamma_2}$$

This confirms (1.16).

### 4.1. Finite-time breakdown

In this subsection, we prove that finite breakdown occurs for the super-critical initial data. To this end, we present the following lemma, which is crucially utilized in obtaining an upper bound of the maximal time for which a local smooth solution persists.

**Lemma 4.1.** For the solution (w(t), s(t)) of (4.1), the following identity holds:

$$t_2 - t_1 \leqslant \int_{s(t_2)}^{s(t_1)} \frac{\mathrm{d}s}{\sqrt{2P_-(s)}},$$

provided that

$$w_0 + \sqrt{2P_-(s_0)} \leqslant 0$$

and  $s(\tau) \in \text{Dom}(P_-)$  for all  $\tau \in [t_1, t_2]$ . Indeed, we have equality in the integral if  $c(t) = c_-$  and  $w_0 + \sqrt{2P_-(s_0)} = 0$ .

*Proof.* By change of coordinates s = s(t), we have

$$\int_{s(t_2)}^{s(t_1)} \frac{\mathrm{d}s}{\sqrt{2P(s)}} = \int_{t_2}^{t_1} \frac{w(t)}{\sqrt{2P(s(t))}} \,\mathrm{d}t \ge \int_{t_2}^{t_1} (-1) \,\mathrm{d}t = t_2 - t_1,$$

where the inequality comes from the weak comparison principle (1.18). Under the latter assumption, we note that (4.2) turns the inequality into equality. This completes the proof.

Several remarks are in order. Suppose that *P* satisfies (1.13) with  $c_1 = \bar{c}$ .

**Remark 4.2.** For the  $0 \le \nu < 2\sqrt{\overline{c}}$  case, we obtain an explicit integral formula:

$$\int_{\text{Dom}(P)} \frac{\mathrm{d}s}{\sqrt{2P(s)}} = \frac{\pi}{\mu_{\bar{c}}}, \quad \text{with } \mu_{\bar{c}} := \sqrt{\bar{c} - \frac{1}{4}\nu^2}.$$

The initial condition in (1.13) can be switched to P(a) = 0 for any  $a \neq 1/\overline{c}$ . In particular, for the  $\nu = 0$  case, we obtain the so-called *elliptic integral*:

$$\int_{\frac{2}{c}-a}^{a} \frac{\mathrm{d}s}{\sqrt{(a-s)(\bar{c}(a+s)-2)}} = \frac{\pi}{\sqrt{\bar{c}}}.$$

**Remark 4.3.** For the  $\nu \ge 2\sqrt{\bar{c}}$  case, we have

$$\int_0^{s_0} \frac{\mathrm{d}s}{\sqrt{2P(s)}} \leq \frac{\log\left(\lambda s_0\right) \vee 2}{\lambda}, \quad \text{with } \lambda := \frac{\nu - \sqrt{\nu^2 - 4\bar{c}}}{2} > 0.$$

for any  $s_0 > 0$ . In particular, it implies that  $Dom(P) = [0, +\infty)$ .

**Proposition 4.4.** Let  $v \ge 2\sqrt{c_-}$ . For any  $(w_0, s_0) \in \Sigma^{\sharp} = \{(w, s) : w \le -\sqrt{2P_-(s)}, s > 0\}$ , the solution to (4.1) with initial data  $(w_0, s_0)$  satisfies  $s(t_*) = 0$  for some finite  $t_* > 0$ .

*Proof.* We claim that  $s(t_*) = 0$  with  $t_* \leq T_0$  where

$$T_0 := \int_0^{s_0} \frac{\mathrm{d}s}{\sqrt{2P_-(s)}}$$

Suppose not, i.e., there exists  $(w_0, s_0) \in \Sigma^{\sharp}$  such that s(t) > 0 for all  $0 \le t \le t_*$  satisfying  $t_* > T_0$ . Then, by the weak comparison principle (1.18), we have

$$w(t) \leq -\sqrt{2P_{-}(s(t))}, \quad 0 \leq t \leq t_*.$$

Hence, by Lemma 4.1, we note in passing that

$$t_* \leq \int_{s(t_*)}^{s_0} \frac{\mathrm{d}s}{\sqrt{2P_{-}(s)}} \leq T_0 < t_*,$$

which leads to a contradiction. Hence, the conclusion of the proposition follows by recalling that  $T_0 < +\infty$  via Remark 4.3.

**Proposition 4.5.** Suppose that  $0 \le v < 2\sqrt{c_{-}}$ . For any

$$(w_0, s_0) \in \Sigma^{\sharp} = \{(w, s) : s > 0\} \setminus \{(w, s) : -\sqrt{2P_{-}(s)} < w < \sqrt{2N_{+}(s)}, s > 0\},\$$

the solution to (4.1) with initial data  $(w_0, s_0)$  satisfies  $s(t_*) = 0$  for some finite  $t_* > 0$ .

*Proof.* We decompose  $\Sigma^{\sharp}$  into three subsets:

$$\begin{split} \Sigma_1^{\sharp} &:= \{ w \leqslant -\sqrt{2P_{-}(s)}, \ 0 < s \leqslant s_* \} \cup \{ w \leqslant 0, \ s > s_* \} \\ \Sigma_2^{\sharp} &:= \{ w > 0, \ s \geqslant s_* \}, \\ \Sigma_3^{\sharp} &:= \{ w \geqslant \sqrt{2N_{+}(s)}, \ 0 < s < s_* \}. \end{split}$$

• For  $(w_0, s_0) \in \Sigma_1^{\sharp}$ , we put  $a := \max\{s_0, s_*\}$ . Consider  $P_*(\cdot)$  satisfying

$$\frac{\mathrm{d}P_*}{\mathrm{d}s} = v\sqrt{2P_*(s)} + 1 - c_{-s}, \quad P_*(a) = 0.$$

Then we have

$$\mathcal{L}_{P_*}(w_0, s_0) = w_0 + \sqrt{2P_*(s_0)} \leq 0.$$

Indeed, for  $s_0 > s_*$ , we have  $w_0 + \sqrt{2P_*(s_0)} = w_0 + \sqrt{2P_*(a)} = w_0 \le 0$ , and for  $0 < s_0 \le s_*$ , we notice that  $w_0 + \sqrt{2P_*(s_0)} = w_0 + \sqrt{2P_-(s_0)} \le 0$ . The weak comparison principle (1.18) holds for *P* as well. Hence, following the argument in the proof of Proposition 4.4 leads us to find that  $s(t_1) = 0$  with

$$t_1 \leq \int_0^a \frac{\mathrm{d}s}{\sqrt{2P(s)}} \leq \frac{\pi}{\mu_-}, \quad \mu_- := \sqrt{c_- - \frac{\nu^2}{4}},$$

where the last inequality follows from Remark 4.2.

• For  $(w_0, s_0) \in \Sigma_2^{\sharp}$ , we prove that  $w(t_2) = 0$ ,  $s(t_2) \ge s_*$  with  $t_2 \le \frac{\pi/2}{\sqrt{c_-}}$ . To this end, we temporarily introduce the following Lyapunov function, defined by

$$\theta_{-}(w,s) := \arctan\left(\frac{\sqrt{c_{-}}(s-\frac{1}{c_{-}})}{w}\right).$$

We claim that

$$\frac{\mathrm{d}}{\mathrm{d}t}\theta_{-}\big((w(t),s(t))\big) \ge \sqrt{c_{-}}$$

as long as  $s(t) \ge \frac{1}{c_{-}}$  and w(t) > 0. It can be shown by the following straightforward computation:

$$\frac{\mathrm{d}}{\mathrm{d}t} \frac{\theta_{-}(w,s)}{\sqrt{c_{-}}} = \frac{1}{1+c_{-}\left(\frac{s-\frac{1}{c_{-}}}{w}\right)^{2}} \left(\frac{w^{2}-(s-\frac{1}{c_{-}})(-vw+1-c(t)s)}{w^{2}}\right)$$
$$= \frac{1}{1+c_{-}\left(\frac{s-\frac{1}{c_{-}}}{w}\right)^{2}} \left(1+c_{-}\left(\frac{s-\frac{1}{c_{-}}}{w}\right)^{2}+v\left(\frac{s-\frac{1}{c_{-}}}{w}\right)+\frac{(s-\frac{1}{c_{-}})(c(t)-c_{-})s}{w^{2}}\right) \ge 1.$$

Hence, we find that  $\theta_{-}(w(t_2), s(t_2)) = \frac{\pi}{2}$  with

$$t_2 \leqslant \frac{\pi/2 - \theta_-(w_0, s_0)}{\sqrt{c_-}} \leqslant \frac{\pi/2}{\sqrt{c_-}}$$

In particular, we obtain  $w(t_2) = 0$ .

• For  $(w_0, s_0) \in \Sigma_3^{\sharp}$ , we argue as in the proof of Proposition 4.4 that  $s(t_3) = s_*, w(t_3) \ge 0$  with

$$t_3 \leq \int_{s_0}^{s_*} \frac{\mathrm{d}s}{\sqrt{2N_+(s)}} \leq \frac{\pi}{\mu_+}, \quad \mu_+ := \sqrt{c_+ - \frac{\nu^2}{4}}$$

Henceforth, we deduce that, for any  $(w_0, s_0) \in \Sigma^{\sharp}$ ,  $s(t_*) = 0$  with

$$t_* \leqslant \frac{\pi}{\mu_-} + \frac{\pi/2}{\sqrt{c_-}} + \frac{\pi}{\mu_+}.$$

This delivers the finite upper bound of  $t_*$  and completes the proof of the proposition.

*Proof of Theorem* 1.7. The proof is based on a combination of the weak comparison principle (1.18) with Propositions 4.4 and 4.5. For any regular initial data ( $\rho_0$ ,  $u_0$ ) in the sense of Theorem 1.1, if there exists  $x \in \mathbb{R}$  such that

$$\left(\frac{u_0'(x)}{\rho_0(x)}, \frac{1}{\rho_0(x)}\right) \in \Sigma^{\sharp},$$

then  $\rho(\cdot, t)$  blows up in a finite time, and the conclusion of the theorem follows.

### 4.2. Global-in-time regularity

To establish the global regularity of the initial data in the sub-critical regime, we prove the strong comparison principle (1.20) in the sequel. Recall from (4.2), (4.3) that if  $s(\tau)$ satisfies  $P_+(s(\tau)) \ge 0$  (resp.  $N_-(s(\tau)) \ge 0$ ) for all  $\tau \in [0, t]$ , then we get

$$\begin{aligned} \mathcal{L}_{P_{+}}(w(t), s(t)) &\geq \mathcal{L}_{P_{+}}(w_{0}, s_{0}) \exp\left(\int_{0}^{t} \frac{1 - c_{+}s(\tau)}{\sqrt{2P_{+}(s(\tau))}} \,\mathrm{d}\tau\right), \\ \mathcal{L}_{N_{-}}(w(t), s(t)) &\leq \mathcal{L}_{N_{-}}(w_{0}, s_{0}) \exp\left(\int_{0}^{t} \frac{c_{-}s(\tau) - 1}{\sqrt{2N_{-}(s(\tau))}} \,\mathrm{d}\tau\right). \end{aligned}$$
(4.5)

The proof for the strict inequalities in the strong comparison principle (1.20) requires us to control the integral in the exponential term in (4.5). We demonstrate this in detail in the following propositions.

**Proposition 4.6.** Let  $v \ge 2\sqrt{c_+}$ . Then

$$\Sigma^{\flat} = \{(w, s) : -\sqrt{2P_{+}(s)} < w, \ s > 0\}$$

is an invariant set for the system (4.1).

*Proof.* We prove that for any  $(w_0, s_0) \in \Sigma^{\flat}$ , the solution to (4.1) satisfies  $(w(t), s(t)) \in \Sigma^{\flat}$  for all  $t \ge 0$ . Suppose not, i.e., there exists  $(w_0, s_0) \in \Sigma^{\flat}$  such that

$$T_* := \inf\{t > 0 : (w(t), s(t)) \notin \Sigma^{\flat}\} < +\infty.$$

In particular, we have  $(w(T_*), s(T_*)) \in \partial \Sigma^{\flat}$ . We decompose the boundary of  $\Sigma^{\flat}$  into two subsets:

$$\partial \Sigma^{\flat} = \partial \Sigma_1^{\flat} \cup \partial \Sigma_2^{\flat} := \left\{ w + \sqrt{2P_+(s)} = 0, \ s \ge 0 \right\} \cup \left\{ w > 0, \ s = 0 \right\}.$$

We will demonstrate that  $T_* < +\infty$  leads to a contradiction. Indeed, the nature of proof is the same for the  $0 < \nu < 2\sqrt{c_+}$  case, which is slightly more complicated. Hence, we defer the details of the proof to the next proposition.

**Proposition 4.7.** Let  $0 < \nu < 2\sqrt{c_+}$ . Then the set

$$\Sigma^{\flat} = \left\{ (w, s) : -\sqrt{2P_{+}(s)} < w < \sqrt{2N_{-}(s)} \right\}$$

is an invariant set for the system (4.1) provided that (1.26) holds.

*Proof.* We prove that for any  $(w_0, s_0) \in \Sigma^{\flat}$ , the solution to (4.1) satisfies  $(w(t), s(t)) \in \Sigma^{\flat}$  for all  $t \ge 0$ . Suppose, on the contrary, that there exists  $(w_0, s_0) \in \Sigma^{\flat}$  such that

$$T_1 := \inf\{t > 0 : (w(t), s(t)) \in \partial \Sigma^{\flat}\} < +\infty$$

We decompose the boundary of  $\Sigma^{\flat}$  into three subsets in the sense of (1.24):

$$\begin{split} \partial \Sigma_1^{\flat} \cup \partial \Sigma_2^{\flat} \cup \partial \Sigma_3^{\flat} &:= \big\{ \mathcal{L}_{P_+}(w, s) = 0, \ 0 \le s < s_* \big\} \\ & \cup \big\{ \mathcal{L}_{N_-}(w, s) = 0, \ 0 < s \le s_* \big\} \cup \big\{ 0 < w \le \sqrt{2N_-(0)}, \ s = 0 \big\}. \end{split}$$

We now consider the following sub-three cases:

•  $(w(T_1), s(T_1)) \in \partial \Sigma_1^{\flat}$ . We claim that there exist a small  $\varepsilon > 0$  and  $C_{\varepsilon} > 0$  such that

$$\int_{T_1-\varepsilon}^{T_1} \frac{1-c_+s(\tau)}{\sqrt{2P_+(s(\tau))}} \,\mathrm{d}\tau \ge -C_\varepsilon. \tag{4.6}$$

If  $s(T_1) < \frac{1}{c_+}$ , then we can take small  $\varepsilon > 0$  such that the integrand becomes non-negative for all  $\tau \in [T_1 - \varepsilon, T_1]$ . On the other hand, if  $s(T_1) \ge \frac{1}{c_+}$ , then we have  $s(T_1) < s_*$  by the assumption. Since  $\sqrt{2P_+(s_*)} = 0$ , we can take small  $\varepsilon > 0$  such that  $P_+(\tau)$  is away from zero for all  $\tau \in [T_1 - \varepsilon, T_1]$ . Hence, this supplies the estimate (4.6). By combining this lower bound of integral (4.6) with the inequality (4.5), we find that

$$w(T_1) + \sqrt{2P_+(s(T_1))} \ge \left(w(T_1 - \varepsilon) + \sqrt{2P_+(s(T_1 - \varepsilon))}\right) \exp(-C_{\varepsilon}) > 0,$$

leading to a contradiction.

•  $(w(T_1), s(T_1)) \in \partial \Sigma_2^{\flat}$ . Similarly, we can prove that there exist a small  $\varepsilon > 0$  and  $C_{\varepsilon} > 0$  such that

$$\int_{T_1-\varepsilon}^{T_1} \frac{c_{-s}(\tau)-1}{\sqrt{2N_{-}(s(\tau))}} \,\mathrm{d}\tau > -C_{\varepsilon}.$$

We find by means of the previous case with (4.5) that

$$w(T_1) + \sqrt{2N_{-}(s(T_1))} \leq \left(w(T_1 - \varepsilon) + \sqrt{2N_{-}(s(T_1 - \varepsilon))}\right) \exp(-C_{\varepsilon}) < 0,$$

which is a contradiction.

•  $(w(T_1), s(T_1)) \in \partial \Sigma_3^{\flat}$ . Noting that  $s(T_1) = 0$  and  $w(T_1) > 0$  conveys us to

$$0 < w(T_1) = s'(T_1) = \lim_{t \to T_1 - \infty} \frac{s(T_1) - s(t)}{T_1 - t} \leq \limsup_{t \to T_1 - \infty} \frac{-s(t)}{T_1 - t} \leq 0,$$

so that we arrive at the contradiction. This completes the proof.

*Proof of Theorem* 1.9. In view of the conclusion that  $\Sigma^{\flat}$  is *invariant* for (4.1) from Propositions 4.6 and 4.7, for any regular initial data ( $\rho_0, u_0$ ) satisfying

$$\left(\frac{u_0'(x)}{\rho_0(x)}, \frac{1}{\rho_0(x)}\right) \in \Sigma^{\flat}, \quad \forall x \in \mathbb{R},$$

we find that the classical solution  $(\rho, u)$  is indeed global by invoking a priori estimate (1.5). Hence, we arrive at the conclusion of the theorem.

As a direct consequence of this result, we present the proof of Theorem 1.12.

*Proof of Theorem* 1.12. We begin by discussing the smallness assumption on  $H_0$ . In [1], the lower and upper bounds on the electric potential for a classical solution to the system (1.27)–(1.28) are investigated as follows:

$$0 < c_{-} \leq e^{\phi(t,x)} \leq c_{+}, \quad c_{-} := \exp(V_{-}^{-1}(H_{0})), \text{ and } c_{+} := \exp(V_{+}^{-1}(H_{0})),$$

where  $V_{\pm}^{-1}$  are inverses of  $V_{\pm}$  which are defined by

$$V(z) := \begin{cases} V_{+}(z) := \int_{0}^{z} \sqrt{2U(r)} \, \mathrm{d}r & \text{for } z \ge 0, \\ V_{-}(z) := \int_{z}^{0} \sqrt{2U(r)} \, \mathrm{d}r & \text{for } z \le 0. \end{cases}$$
(4.7)

Note that  $V_{\pm}(z)$  have the inverse functions and  $V \in \mathcal{C}^2(\mathbb{R})$ ; see [1, Section 2.1].

Indeed, careful estimates via Taylor's expansion suggest that

$$V_{\pm}(z) = \frac{1}{2}z^2 + \frac{1}{9}z^3 + \mathcal{O}(z^4),$$

so that

$$V_{\pm}^{-1}(x) = \pm \sqrt{2x} + \mathcal{O}(x), \quad 0 < x \ll 1.$$

Hence, for small enough  $H_0 \ll 1$ , we have

$$\frac{c_+}{c_-} = 1 + 2\sqrt{2H_0} + \mathcal{O}(H_0).$$

We are now ready to investigate what secures the closing condition to hold; we go through as

$$s_{**} \leq 0 \iff \left( (1 + e^{\gamma_+}) \frac{c_-}{c_+} - 1 \right) e^{\gamma_-} \geq 1$$
$$\iff 1 + e^{\gamma_+} \geq \frac{2c_+}{c_-}$$
$$\iff \gamma_+ \geq 2 \left( \frac{c_+}{c_-} - 1 \right) = 4\sqrt{2H_0} + \mathcal{O}(H_0)$$
$$\iff \nu \pi \geq \sqrt{4c_+ - \nu^2} (4\sqrt{2H_0} + \mathcal{O}(H_0)).$$

Since  $\sqrt{4c_+ - v^2} = \mathcal{O}(1)$  for small v > 0, we find that (1.30) guarantees the closing condition as desired.

# A. Auxiliary lemma on the neutrality condition

In this appendix, we aim to prove the following lemma.

**Lemma A.1.** For  $f \in L^2 \cap \dot{W}^{-1,p}(\mathbb{R})$ , the improper integral satisfies

$$\int_{\mathbb{R}} f(x) \, \mathrm{d}x = 0.$$

The proof relies on the following classical result, which is included here for completeness.

**Proposition A.2.** For  $f \in \dot{W}^{-1,p}(\mathbb{R})$ , there exists a unique  $F \in L^p(\mathbb{R})$  such that F' = f and

 $\|f\|_{\dot{W}^{-1,p}} = \|F\|_{L^p}.$ 

Additionally, if  $f \in L^2(\mathbb{R})$ , then  $|F(x)| \to 0$  as  $|x| \to \infty$ .

*Proof.* By the definition of the  $\dot{W}^{-1,p}$  norm, there exists  $F \in L^p(\mathbb{R})$  such that F' = f and

$$||F||_{L^p} \lesssim ||f||_{\dot{W}^{-1,p}} + 1 < +\infty.$$

On the other hand, if  $\tilde{F}' = f$  then, there exists a constant  $c \in \mathbb{R}$  such that

$$\widetilde{F} = F + c \quad \text{in } \mathbb{R}$$

If  $c \neq 0$ , then  $\tilde{F} \notin L^p$ . Thus, there exists a unique antiderivative of f with finite  $L^p$ -norm, and we deduce

$$\|f\|_{\dot{W}^{-1,p}} = \|F\|_{L^p}.$$

If we further assume that  $f \in L^2(\mathbb{R})$ , then F is 1/2-Hölder continuous, and one can readily prove that F is bounded as well: namely, for any  $x \in \mathbb{R}$ ,

$$2|F(x)| \leq \int_{x-1}^{x+1} |F(y)| \, \mathrm{d}y + \int_{x-1}^{x+1} |F(x) - F(y)| \, \mathrm{d}y$$
  
$$\leq 2^{1-\frac{1}{p}} \|F\|_{L^{p}([x-1,x+1])} + \int_{x-1}^{x+1} \|f\|_{L^{2}} |x-y|^{\frac{1}{2}} \, \mathrm{d}y$$
  
$$\leq 2(\|F\|_{L^{p}} + \|f\|_{L^{2}}). \tag{A.1}$$

Thus, we get  $||F||_{L^{\infty}} \leq ||f||_{L^2 \cap \dot{W}^{-1,p}}$ . We let  $G(x) := |F(x)|^p$ , then  $G \in L^1(\mathbb{R})$  and

$$|G(x) - G(y)| = p ||F||_{L^{\infty}}^{p-1} [F]_{\mathcal{C}^{0,\frac{1}{2}}} \sqrt{|x-y|} \le p ||f||_{L^{2} \cap \dot{W}^{-1,p}}^{p} \sqrt{|x-y|}.$$

Since G(x) is uniformly continuous and  $G \in L^1(\mathbb{R})$ , we have

$$|G(x)| \to 0$$
 as  $|x| \to \infty$ .

which directly implies that  $|F(x)| \to 0$  as  $|x| \to \infty$ . This completes the proof.

*Proof of Lemma* A.1. From Proposition A.2, there exists a unique  $F \in L^p(\mathbb{R})$  such that F' = f, and thus we find

$$\lim_{b \to \infty} \int_0^b f(x) \, \mathrm{d}x = \lim_{b \to \infty} (F(b) - F(0)) = -F(0).$$

Similarly, we obtain

$$\lim_{a \to -\infty} \int_{a}^{0} f(x) \, \mathrm{d}x = \lim_{a \to -\infty} (F(0) - F(a)) = F(0),$$

and thus

$$\int_{-\infty}^{\infty} f(x) \, \mathrm{d}x = \lim_{a \to -\infty} \int_{a}^{0} f(x) \, \mathrm{d}x + \lim_{b \to \infty} \int_{0}^{b} f(x) \, \mathrm{d}x = 0.$$

Hence, the improper integral satisfies the desired property, completing the proof.

# B. A priori estimates in the proof of Theorem 1.1

In this appendix, we provide the details of the a priori estimate (2.1). For the sake of simplicity, we assume k = 1 and  $\nu = 0$ .

• Estimates of  $\|\rho - \bar{c}\|_{H^s \cap \dot{W}^{-1,p}}$ : We rewrite the continuity equation (1.1)<sub>1</sub> as

$$\partial_t (\rho - \bar{c}) + \partial_x ((\rho - \bar{c})u) = -\bar{c}\partial_x u$$
 (B.1)

and employ the dual formulation of  $\dot{W}^{-1,p}$  to find that

$$\frac{\mathrm{d}}{\mathrm{d}t} \|\rho - \bar{c}\|_{\dot{W}^{-1,p}} \leq \|(\rho - \bar{c})u\|_{L^p} + \bar{c}\|u\|_{L^p} \leq \|\rho - \bar{c}\|_{L^{\infty}} \|u\|_{L^p} + \bar{c}\|u\|_{L^p}.$$
(B.2)

For the higher-order estimates of  $\rho - \bar{c}$ , we differentiate (B.1) by  $\partial_x^l$  for each  $l = 0, 1, \dots, s$ , and we test  $\partial_x^l (\rho - \bar{c})$  to find that

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int_{\mathbb{R}}|\partial_{x}^{l}(\rho-\bar{c})|^{2}\,\mathrm{d}x = -\int_{\mathbb{R}}\partial_{x}^{l}(\rho-\bar{c})u\partial_{x}^{l+1}(\rho-\bar{c})\,\mathrm{d}x$$
$$-\int_{\mathbb{R}}\partial_{x}^{l}(\rho-\bar{c})[\partial_{x}^{l+1},u](\rho-\bar{c})\,\mathrm{d}x$$
$$-\bar{c}\int_{\mathbb{R}}\partial_{x}^{l}(\rho-\bar{c})\partial_{x}^{l+1}u\,\mathrm{d}x$$
$$=:\mathrm{I}+\mathrm{II}+\mathrm{III},$$

where  $[\cdot, \cdot]$  stands for the commutator operator, i.e., [A, B] = AB - BA. Then standard commutator estimates and Sobolev inequalities yield

$$\begin{split} \mathbf{I} &\leq \|\partial_{x}u\|_{L^{\infty}} \|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}}^{2},\\ \mathbf{II} &\leq \|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}} \|[\partial_{x}^{l+1}, u](\rho - \bar{c})\|_{L^{2}}\\ &\leq C \|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}} (\|\partial_{x}u\|_{L^{\infty}} \|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}} + \|\partial_{x}^{l+1}u\|_{L^{2}} \|\rho - \bar{c}\|_{L^{\infty}}),\\ \mathbf{III} &\leq \bar{c} \|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}} \|\partial_{x}^{l+1}u\|_{L^{2}}. \end{split}$$

Thus, we have

$$\frac{\mathrm{d}}{\mathrm{d}t} \|\partial_x^l(\rho - \bar{c})\|_{L^2} \leq C(\|\rho - \bar{c}\|_{L^{\infty}} + \bar{c} + \|\partial_x u\|_{L^{\infty}})(\|\partial_x^l(\rho - \bar{c})\|_{L^2} + \|\partial_x^{l+1}u\|_{L^2}).$$
(B.3)

• Estimates of  $||u||_{L^p} + ||\partial_x u||_{H^s}$ : We first estimate

$$\frac{1}{p} \frac{\mathrm{d}}{\mathrm{d}t} \int_{\mathbb{R}} u^{p} \mathrm{d}x \leq \|\partial_{x}u\|_{L^{\infty}} \|u\|_{L^{p}}^{p} + \|E\|_{L^{p}} \|u\|_{L^{p}}^{p-1} 
= \|u\|_{L^{p}}^{p-1} (\|\partial_{x}u\|_{L^{\infty}} \|u\|_{L^{p}} + \|\rho - c\|_{\dot{W}^{-1,p}}) 
\leq \|u\|_{L^{p}}^{p-1} (\|\partial_{x}u\|_{L^{\infty}} \|u\|_{L^{p}} + \|\rho - \bar{c}\|_{\dot{W}^{-1,p}} + \|\eta\|_{\dot{W}^{-1,p}}). \quad (B.4)$$

For the higher-order estimates of u, where l = 0, 1, ..., s, we deduce

$$\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}} |\partial_{x}^{l+1}u|^{2} dx 
= -\int_{\mathbb{R}} (\partial_{x}^{l+1}u) (\partial_{x}^{l+2}u) u dx - \int_{\mathbb{R}} \partial_{x}^{l+1}u [\partial_{x}^{l+1}, u] \partial_{x}u dx 
+ \int_{\mathbb{R}} \partial_{x}^{l+1}u \partial_{x}^{l}(\rho - c) dx 
\leq C \|\partial_{x}u\|_{L^{\infty}} \|\partial_{x}^{l+1}u\|_{L^{2}}^{2} + \|\partial_{x}^{l+1}u\|_{L^{2}} \|\partial_{x}^{l}(\rho - c)\|_{L^{2}} 
\leq C \|\partial_{x}u\|_{L^{\infty}} \|\partial_{x}^{l+1}u\|_{L^{2}}^{2} + \|\partial_{x}^{l+1}u\|_{L^{2}} (\|\partial_{x}^{l}(\rho - \bar{c})\|_{L^{2}} + \|\eta\|_{\dot{H}^{l}}). \quad (B.5)$$

Combining (B.2), (B.3), (B.4), and (B.5), we conclude the estimate (2.1).

Acknowledgments. The authors thank Junsik Bae, Bongsuk Kwon, and Changhui Tan for helpful discussion on the Euler–Poisson system for cold-ion dynamics and Euler– Poisson–alignment system.

**Funding.** The research of YPC and of DK was supported by NRF grants 2022R1A2C1 002820 and RS-2024-00406821. The research of DK was supported by NRF grant RS-2023-00212227. The research of ET was supported in part by ONR grants N00014-2112773 and N00014-2412659.

# References

- J. Bae, J. Choi, and B. Kwon, Formation of singularities in plasma ion dynamics. *Nonlinearity* 37 (2024), no. 4, article no. 045011 Zbl 1541.35365 MR 4723815
- J. Bae and B. Kwon, Small amplitude limit of solitary waves for the Euler–Poisson system. J. Differential Equations 266 (2019), no. 6, 3450–3478 Zbl 1406.35256 MR 3912689
- [3] J. Bae and B. Kwon, Linear stability of solitary waves for the isothermal Euler–Poisson system. Arch. Ration. Mech. Anal. 243 (2022), no. 1, 257–327 Zbl 1507.35162 MR 4359453
- [4] M. Bhatnagar and H. Liu, Critical thresholds in 1D pressureless Euler–Poisson systems with variable background. *Phys. D* 414 (2020), article no. 132728 Zbl 1490.35256 MR 4155319
- [5] M. Bhatnagar and H. Liu, Critical thresholds in one-dimensional damped Euler-Poisson systems. *Math. Models Methods Appl. Sci.* 30 (2020), no. 5, 891–916 Zbl 1453.35145 MR 4098218
- [6] M. Bhatnagar and H. Liu, A complete characterization of sharp thresholds to spherically symmetric multidimensional pressureless Euler–Poisson systems. 2023, arXiv:2302.04428v1
- [7] M. Bhatnagar, H. Liu, and C. Tan, Critical thresholds in the Euler–Poisson–alignment system. J. Differential Equations 375 (2023), 82–119 Zbl 1527.35227 MR 4629746
- [8] M. P. Brenner and T. P. Witelski, On spherically symmetric gravitational collapse. J. Statist. Phys. 93 (1998), no. 3-4, 863–899 Zbl 0933.35160 MR 1666522
- [9] J. A. Carrillo, Y.-P. Choi, E. Tadmor, and C. Tan, Critical thresholds in 1D Euler equations with non-local forces. *Math. Models Methods Appl. Sci.* 26 (2016), no. 1, 185–206 Zbl 1356.35174 MR 3417728

- [10] J. A. Carrillo, Y.-P. Choi, and E. Zatorska, On the pressureless damped Euler–Poisson equations with quadratic confinement: Critical thresholds and large-time behavior. *Math. Models Methods Appl. Sci.* 26 (2016), no. 12, 2311–2340 Zbl 1349.35378 MR 3564592
- [11] F. Chen, Introduction to plasma physics and controlled fusion, vol. 1. Plenum Press, New York, 1984
- [12] Y. Deng, T.-P. Liu, T. Yang, and Z.-a. Yao, Solutions of Euler–Poisson equations for gaseous stars. Arch. Ration. Mech. Anal. 164 (2002), no. 3, 261–285 Zbl 1038.76036 MR 1930393
- [13] S. Engelberg, H. Liu, and E. Tadmor, Critical thresholds in Euler–Poisson equations. Indiana Univ. Math. J. 50 (2001), 109–157 Zbl 0989.35110 MR 1855666
- [14] Y. Guo, L. Han, and J. Zhang, Absence of shocks for one dimensional Euler–Poisson system. Arch. Ration. Mech. Anal. 223 (2017), no. 3, 1057–1121 Zbl 1361.35144 MR 3595365
- [15] Y. Guo and B. Pausader, Global smooth ion dynamics in the Euler–Poisson system. Comm. Math. Phys. 303 (2011), no. 1, 89–125 Zbl 1220.35129 MR 2775116
- [16] D. Holm, S. F. Johnson, and K. E. Lonngren, Expansion of a cold ion cloud. Appl. Phys. Lett. 38 (1981), 519–521
- [17] A. D. Ionescu and B. Pausader, The Euler–Poisson system in 2D: Global stability of the constant equilibrium solution. *Int. Math. Res. Not. IMRN* (2013), no. 4, 761–826 Zbl 1320.35270 MR 3024265
- [18] D. Lannes, F. Linares, and J.-C. Saut, The Cauchy problem for the Euler–Poisson system and derivation of the Zakharov–Kuznetsov equation. In *Studies in phase space analysis* with applications to PDEs, pp. 181–213, Progr. Nonlinear Differential Equations Appl. 84, Birkhäuser/Springer, New York, 2013 Zbl 1273.35263 MR 3185896
- [19] D. Li and Y. Wu, The Cauchy problem for the two dimensional Euler–Poisson system. J. Eur. Math. Soc. (JEMS) 16 (2014), no. 10, 2211–2266 Zbl 1308.35220 MR 3274788
- [20] F. Lin and P. Zhang, On the hydrodynamic limit of Ginzburg–Landau wave vortices. Comm. Pure Appl. Math. 55 (2002), no. 7, 831–856 Zbl 1033.35089 MR 1894157
- [21] C. Liu and Y.-j. Peng, Global convergence of the Euler–Poisson system for ion dynamics. Math. Methods Appl. Sci. 42 (2019), no. 4, 1236–1248 Zbl 1457.35075 MR 3912086
- [22] H. Liu, Wave breaking in a class of nonlocal dispersive wave equations. J. Nonlinear Math. Phys. 13 (2006), no. 3, 441–466 Zbl 1110.35069 MR 2249812
- [23] H. Liu and E. Tadmor, Spectral dynamics of the velocity gradient field in restricted flows. *Comm. Math. Phys.* 228 (2002), no. 3, 435–466 Zbl 1031.76006 MR 1918784
- [24] H. Liu and E. Tadmor, Critical thresholds in 2D restricted Euler–Poisson equations. SIAM J. Appl. Math. 63 (2003), no. 6, 1889–1910 Zbl 1073.35187 MR 2030849
- [25] A. Majda, Introduction to PDEs and waves for the atmosphere and ocean. Courant Lect. Notes Math. 9, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003 Zbl 1278.76004 MR 1965452
- [26] P. A. Markowich, C. A. Ringhofer, and C. Schmeiser, *Semiconductor equations*. Springer, Vienna, 1990 Zbl 0765.35001 MR 1063852
- [27] E. Tadmor and C. Tan, Critical thresholds in flocking hydrodynamics with non-local alignment. *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **372** (2014), no. 2028, article no. 20130401 Zbl 1353.76064 MR 3268063
- [28] E. Tadmor and D. Wei, On the global regularity of subcritical Euler–Poisson equations with pressure. J. Eur. Math. Soc. (JEMS) 10 (2008), no. 3, 757–769 Zbl 1155.35078 MR 2421160
- [29] C. Tan, Eulerian dynamics in multidimensions with radial symmetry. SIAM J. Math. Anal. 53 (2021), no. 3, 3040–3071 Zbl 1472.35311 MR 4263431

[30] D. Wei, E. Tadmor, and H. Bae, Critical thresholds in multi-dimensional Euler–Poisson equations with radial symmetry. *Commun. Math. Sci.* 10 (2012), no. 1, 75–86 Zbl 1272.35146 MR 2901301

Received 20 February 2024; revised 12 January 2025; accepted 24 April 2025.

### Young-Pil Choi

Department of Mathematics, Yonsei University, 03722 Seoul, South Korea; ypchoi@yonsei.ac.kr

### Dong-ha Kim

Department of Mathematics, Yonsei University, 03722 Seoul, South Korea; skyblue898@yonsei.ac.kr

### Dowan Koo

Department of Mathematics, Yonsei University, 03722 Seoul, South Korea; dowan.koo@yonsei.ac.kr

### Eitan Tadmor

Department of Mathematics and IPST, University of Maryland, College Park, MD 20742, USA; tadmor@umd.edu