


# Calculus 140, section 10.3 Conic Sections

notes by Tim Pilachowski

“The conic sections arise when a double right circular cone is cut by a plane.”




“Any second-degree equation  $Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$  is (except in degenerate cases) an equation of a parabola, an ellipse, or a hyperbola.” By using completing the square, we can determine shifts/translations  $(x - h)$  and  $(y - k)$  from “standard” position. (You’ll need this for some homework exercises.)


In the  $xy$  real number plane, a degenerate conic can be one of the three cases shown in the lower pictures above, or the null set (i.e., no points).

conic section	basic equation	reference line	focus/ foci	vertex/ vertices	standard position ( $h = k = 0$ ) symmetry	asymptote(s)
parabola						
ellipse						
hyperbola						

You know the basic quadratic function from Algebra,  $y = x^2$ , and its shape: a parabola.



If we turn that parabolic shape clockwise by 90 degrees, we get a slightly different equation:  $x = y^2$ .



Definition 10.1: “Let  $l$  be a fixed line and  $P$  a fixed point not on  $l$ . The set of all points in the plane equidistant from  $l$  and  $P$  is called a **parabola**.”

The line  $l$  is called the **directrix** and the point  $P$  is called the **focus**.

Applications:

The point midway between directrix and focus is the **vertex**. (You used this designation in Algebra.)

“Standard position” is with the  $x$ -axis or the  $y$ -axis as the axis of symmetry.

The text uses the distance formula to derive standard forms  $x^2 = 4cy$  and  $y^2 = 4cx$ .

A parabola has no asymptotes.


Other applications:

The non-standard position forms are  $(x - h)^2 = 4c(y - k)$  and  $(y - k)^2 = 4c(x - h)$ .

We can use shifts/translations (just like in Algebra) to help us graph or answer questions.

For finding derivatives, we have implicit differentiation (section 3.6).

Example A. The vertex is  $(-2, 0)$ , and the directrix is  $x = \frac{3}{2}$ . Identify the focus of the parabola, find its equation, and then sketch the graph.



Definition 10.2: “Let  $P_1$  and  $P_2$  be two points in the plane, and let  $k$  be a number greater than the distance between  $P_1$  and  $P_2$ . The set of all points  $P$  in the plane such that  $|P_1P| + |P_2P| = k$  is called an **ellipse**.”

The text uses the distance formula to derive standard forms  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and

$$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1, \text{ where } 0 < b \leq a.$$

Note that a larger denominator under  $x^2$  gives a horizontal orientation to an ellipse, and a larger denominator under  $y^2$  gives a vertical orientation to an ellipse.

The reference lines are called the **major axis** and the **minor axis**, with the major axis being the longer of the two (or in the special case of a circle, equal in length).

Geometric interpretation:  $2a$  is the length of the major axis, and  $2b$  is the length of the minor axis.

In standard position, the major axis will have equation either  $y = 0$  (horizontal orientation) or  $x = 0$  (vertical orientation).

The points  $P_1$  and  $P_2$  are called the **foci** (the plural of focus).

Given  $c = \sqrt{a^2 - b^2}$  (Pythagorean/distance formula), in standard position, the foci will have coordinates of

either  $(-c, 0)$  and  $(c, 0)$  when there is a horizontal orientation

or  $(0, -c)$  and  $(0, c)$  when there is a vertical orientation.

Geometric interpretation:  $2c$  is the distance between the two foci.

When  $a = b$ , the major axis and the minor axis have the same length, and the two foci are a single point: the center of a circle.

Applications:

The points where the major axis intersects the ellipse are the **vertices**.

In standard position, the vertices will have coordinates of

either  $(-a, 0)$  and  $(a, 0)$  when there is a horizontal orientation

or  $(0, -a)$  and  $(0, a)$  when there is a vertical orientation.



In standard position, an ellipse (either orientation) will be symmetric with respect to the  $x$ -axis, the  $y$ -axis and also the origin.

An ellipse has no asymptotes.


The non-standard position forms are  $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$  and  $\frac{(x-h)^2}{b^2} + \frac{(y-k)^2}{a^2} = 1$ .

We can use shifts/translations (just like in Algebra) to help us graph or answer questions.

For finding derivatives, we have implicit differentiation (section 3.6).



Example B. An ellipse passes through the points  $\left(1, \frac{\sqrt{27}}{2}\right)$  and  $\left(-\frac{1}{2}, \frac{\sqrt{135}}{4}\right)$  and is in standard position. Find the equation and sketch the graph.



Definition 10.3: “Let  $P_1$  and  $P_2$  be two points in the plane, and let  $k$  be a positive number less than the distance between  $P_1$  and  $P_2$ . The set of all points  $P$  in the plane such that  $||P_1P| - |P_2P|| = k$  called a **hyperbola**.”

The text uses the distance formula to derive standard forms  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and

$$\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1.$$

The points  $P_1$  and  $P_2$  are called the **foci** (the plural of focus).

Given  $c = \sqrt{a^2 + b^2}$  (Pythagorean/distance formula), in standard position, the foci will have coordinates of

either  $(-c, 0)$  and  $(c, 0)$  when there is a horizontal orientation  
or  $(0, -c)$  and  $(0, c)$  when there is a vertical orientation.

Geometric interpretation:  $2c$  is the distance between the two foci.

The point located halfway between the two foci is called the **center** of the hyperbola. In standard position, the center of the hyperbola will have coordinates  $(0, 0)$ .

The reference line, the line through the two foci, is called the **principal axis**.

In standard position, the major axis will have equation either  $y = 0$  (horizontal orientation) or  $x = 0$  (vertical orientation).



The points where the principal axis intersects the hyperbola are the **vertices**.

In standard position, the vertices will have coordinates of  
either  $(-a, 0)$  and  $(a, 0)$  when there is a horizontal orientation  
or  $(0, -a)$  and  $(0, a)$  when there is a vertical orientation.

The line segment connecting the two vertices is called the **transverse axis** of the hyperbola.

Geometric interpretation:  $2a$  is the length of the transverse axis.

In standard position, a hyperbola (either orientation) will be symmetric with respect to the  $x$ -axis, the  $y$ -axis and also the origin.



A hyperbola has two asymptotes:  $y = \frac{b}{a}x$  and  $y = -\frac{b}{a}x$  (when there is a horizontal orientation) or  $y = \frac{a}{b}x$  and  $y = -\frac{a}{b}x$  (when there is a vertical orientation).

Examples of hyperbolic applications are noted in the text.

The non-standard position forms are  $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$  and  $\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$ .

We can use shifts/translations (just like in Algebra) to help us graph or answer questions. For finding derivatives, we have implicit differentiation (section 3.6).

Example C. A hyperbola has foci  $(0, -\sqrt{52})$  and  $(0, \sqrt{52})$ , has asymptotes

$y = -\frac{2}{3}x$  and  $y = \frac{2}{3}x$ , and is in standard position. Find the equation and sketch the graph.

